Bile duct stone extraction under endoscopic ultrasound guidance without fluoroscopy or contrast injection

Fig. 1 Magnetic resonance imaging view of a distal common bile duct stone.

Fig. 2 Endoscopic ultrasound (EUS) demonstrated a stone in the distal common bile duct.

Fig. 3 A fistulotomy was performed under EUS guidance.

A 57-year-old woman with an allergy to iodinated contrast presented with intermittent right upper quadrant pain. Laboratory test results were as follows: total bilirubin 4 mg/dL, alkaline phosphatase 164 mg/dL, and normal amylase, aspartate transaminase, and alanine transaminase levels. A solitary 1-cm calculus was identified in the distal common bile duct on magnetic resonance imaging (● **Fig. 1**); linear-array endosonography confirmed the presence of a 0.9-cm calculus (● **Fig. 2**). Using a needle-knife, a 0.035-inch guide wire, and a 12-mm extraction balloon (all Boston Scientific, Natick, Massachusetts, USA), the bile duct was cannulated, the papilotomy exten-

ded, and the stone extracted, all under endoscopic ultrasound (EUS) guidance (● **Fig. 3, 4**). Standard cannulation under EUS guidance was unsuccessful, probably due to the presence of stone in the distal common bile duct/ampulla. There were no complications resulting from the endoscopic intervention. The patient underwent an uneventful laparoscopic cholecystectomy and no residual stones were noted at intraoperative cholangiography. The injection of contrast during endoscopic retrograde cholangiopancreatography (ERCP) has been linked to post-ERCP pancreatitis. In addition, in patients with contrast allergy, ERCP might have to be delayed for up to 12 hours until ster-

oids have been administered, delaying emergency therapeutic interventions in patients with serious conditions such as severe cholangitis or gallstone pancreatitis [1–3]. Fluoroscopy is also expensive and is a limited resource, given that its availability is often controlled by radiology departments. At the same time, EUS is becoming widely available and its range of indications is expanding [4]. We have demonstrated the feasibility of performing therapeutic interventions in the bile duct under EUS guidance alone, without fluoroscopy and contrast injection. This strategy has potential applications in pregnant women requiring therapeutic interventions in the bile duct, in patients with contrast allergy who require emergency ERCP, and also in patients with bile duct abnormalities identified at EUS in whom an additional ERCP could be avoided using this technique [5].

Endoscopy_UCTN_Code_TTT_1AS_2AD
Endoscopy_UCTN_Code_TTT_1AR_2AH

E. L. A. Artifon¹, A. Kumar², S. Marques¹, S. Ishioka¹, P. Sakai³, M. S. Bhutani⁴

Fig. 4 EUS image of the needle-knife.

Bibliography

DOI 10.1055/s-2007-966551

Endoscopy 2008; 40: 13 – 14

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

A. Kumar, MD

Northport VA Medical Center

79 Middleville Road

Northport

New York 11790

Fax: +1-631-486-6113

atul.kumar2@va.gov

¹ GI Endoscopy Division, University of Sao Paulo Medical School, Sao Paulo, Brazil

² Department of Gastroenterology and Hepatology, Northport VA Medical Center and Stony Brook University, Northport, New York, USA

³ GI Endoscopy Unit, Hospital Das Clinicas, Sao Paulo, Brazil

⁴ MD Anderson, Gastroenterology and Hepatology, Houston, Texas, USA

References

- 1 Moon JH, Cho YD, Cha SW *et al.* The detection of bile duct stones in suspected biliary pancreatitis: comparison of MRCP, ERCP, and intraductal US. *Am J Gastroenterol* 2005; 100: 1051 – 1057
- 2 Kahaleh M, Hernandez AJ, Tokar J *et al.* Interventional EUS-guided cholangiography: evaluation of a technique in evolution. *Gastrointest Endosc* 2006; 64: 52 – 59
- 3 Draganov P, Cotton PB. Iodinated contrast sensitivity in ERCP. *Am J Gastroenterol* 2000; 95: 1398 – 1401
- 4 Scheiman JM, Carlos RC, Barnett JL *et al.* Can endoscopic ultrasound or magnetic resonance cholangiopancreatography replace ERCP in patients with suspected biliary disease? A prospective trial and cost analysis. *Am J Gastroenterol* 2001; 96: 2900 – 2904
- 5 Wamsteker EJ. Updates in biliary endoscopy. *Curr Opin Gastroenterol* 2006; 22: 300 – 304