Replacing and relocating percutaneous endoscopic gastrostomy tube without esophagogastroscope

Figure 1 After the PEG tube was gently pulled out by external traction from the infected gastrostomy site, a standard-size esophagogastroscope was inserted via the same gastrotocutaneous tract to perform gastroscopy. A new site was selected well away from the infected previous stoma and a normal pull-back PEG was carried out, except that the string was pulled out and the new PEG feeding tube was inserted through the old gastrostomy opening. A “second-look” gastroscopy was also carried out easily, causing no discomfort for the patient.

Percutaneous endoscopic gastrostomy (PEG) [1] is widely used for the enteral nutrition of patients with head and neck cancer. While very safe and well tolerated, it is not without its complications. Long-term use of PEG can lead to complications requiring replacement of the gastrostomy tube and relocation of the feeding tube to a new site [2, 3]. Replacing the

Figure 2 PEG feeding tube positioned well away from the infected previous gastrostomy site.

feeding tube is an easy, straightforward procedure if done shortly after the removal of the old one, while relocating it to a new site requires repeat esophagogastroscope. However, standard peroral or transnasal introduction of the esophagogastroscope is not always possible in patients with head and neck cancers. Narrowness of the pharyngeal space in consequence of major head and neck surgery and radiotherapy can impede peroral feeding, which makes the patient dependent on feeding by gastrostomy tube for life. Replacing and relocating the gastrostomy tube for such patients to a new site in the case of any complication cannot be done endoscopically in the usual way. The technique illustrated in **Figure 1** and **2** allows endoscopically controlled replacement and relocation of the PEG feeding tube without the performance of a traditional peroral esophagogastroscope.

No general anesthesia was required. Both the gastroscope and the PEG feeding tube was passed through the former gastrostomy opening without any difficulty. I recommend the described technique of replacing and relocating the PEG feeding tube for patients with head and neck cancer, if peroral esophagogastroscope cannot be carried out.

Endoscopy_UCTN_Code_TTT_1AO_2AK

L. Lujber

Department of Oto-Rhino-Laryngology, Head and Neck Surgery, Pécs University, Medical School, Pécs, Hungary

References

- 1 Gauderer MW, Ponsky JL, Izant RJ. Gastrostomy without laparotomy: a percutaneous endoscopic technique. *J Pediatr Surg* 1980; 15: 872–875
- 2 Bumpers HL, Collure DW, Best IM et al. Unusual complications of long-term percutaneous gastrostomy tubes. *J Gastrointest Surg* 2003; 7: 917–920
- 3 Anwander T, Berge S, Appel T et al. Percutaneous endoscopic gastrostomy for long-term feeding of patients with oropharyngeal tumors. *Nutr Cancer* 2004; 50: 40–45

Bibliography

DOI 10.1055/s-2007-966567

Endoscopy 2007; 39: E174

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

L. Lujber, MD, PhD

Department of Oto-Rhino-Laryngology, Head and Neck Surgery
Pécs University Medical School
Munkácsy M. str. 2
7621 Pécs
Hungary
Fax: +36-72-228119
Lujber@yahoo.com