Giant subcutaneous abscess after percutaneous endoscopic gastrostomy

Fig. 1 The gastrostomy tube was placed in the stomach.

A 69-year-old man was admitted to our hospital with dysphagia, which had resulted from multiple cerebral infarctions. A 20-Fr gastrostomy tube was inserted under endoscopic guidance. The procedure was uneventful, and antibiotics were not administered (● **Fig. 1**). Although the skin around the fistula showed no clinical signs of abnormality, the patient's abdominal pain increased. Computed tomography showed a low-density area in the left subcutaneous space (● **Fig. 2**), and *Enterococcus faecium* was identified in cultures of pus from the stoma. The subcutaneous abscess was treated by surgical drainage and he was given sulbactam/cefoperazone. The abscess responded well to treatment and disappeared 2 months after insertion of the percutaneous endoscopic gastrostomy.

Peristomal wound infection after insertion of a percutaneous endoscopic gastrostomy is usually caused by oral bacteria. In the present case, however, a giant subcutaneous abscess was caused by *Enterococcus faecium*. The patient had undergone a Billroth II distal gastrectomy for the treatment of a peptic ulcer. The first explanation for the formation of this giant abscess in this patient could be that the resection of the pylorus and the lack of gastric acid might have enabled enterobacteria to flow back into the stomach. Secondly, because the gastrectomy scar was known to be hypovascular, the patient's subcutaneous tissue could have been more susceptible to infection.

Fig. 2 Computed tomographic views showing a large low-density area in the left subcutaneous space.

Endoscopy_UCTN_Code_CPL_1AH_2AI

**H. Sakai¹, M. Inamori¹, T. Sato¹,
K. Akimoto¹, T. Akiyama¹, K. Fujita¹,
M. Yoneda¹, H. Takahashi¹, A. Goto¹,
A. Kusakabe¹, Y. Abe¹, H. Kirikoshi¹,
K. Kubota¹, N. Ueno¹, Y. Rino²,
A. Nakajima¹**

¹ Gastroenterology Division, Yokohama City University School of Medicine, Yokohama, Japan

² Department of Surgery, Yokohama City University School of Medicine, Yokohama, Japan

Bibliography

DOI 10.1055/s-2007-966598

Endoscopy 2007; 39: E264

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

M. Inamori, MD

Gastroenterology Division
Yokohama City University School of Medicine
3-9 Fukuura
Kanazawa-ku
Yokohama 236-0004
Japan
Fax: +81-45-784-3546
inamorim@med.yokohama-cu.ac.jp