An endoscopic salvage procedure for the treatment of regurgitation occurring after an unsuccessful esophagectomy with gastric tube reconstruction and subsequent colonic bypass surgery

Fig. 1 A water-soluble contrast study and a schematic illustration key showing the gastric tube without filling of the colonic bypass (A, tracheal stent; B, esophagus proximal to the stenotic collar anastomosis; C, the gastric tube; D, location of the colonic bypass).

Fig. 3 A water-soluble contrast study and a schematic illustration key showing the appearance after endoscopic de-epithelialization of the stenotic gastric tube and sealing with bucrylate and histoacrylate. An adequate efflux of the contrast medium without filling of the sealed gastric tube was achieved (A, tracheal stent; B, esophagus proximal to the stenotic collar anastomosis; D, the colonic bypass).

Anastomotic leaks and strictures after esophagectomy are associated with high morbidity and mortality [1,2]. We present here a patient who had a subtotal esophagectomy with gastric interposition because of esophageal adenocarcinoma. Anastomotic leakage occurred and was treated by endoscopic applications of fibrin glue. One month after discharge she presented with a large (2.5 cm × 5 cm) iatrogenic esophagotracheal fistula. Because of the extensive defect in the pars membranacea and the inflammation we decided against surgical intervention and treated the defect endoscopically using a

Fig. 2 a Upper gastrointestinal endoscopic image showing a sufficiently wide collar anastomosis of the side-to-side esophagocolostomy. b A stenotic collar anastomosis with a siphon-like anatomical configuration at the entrance to the gastric interposition.

combination of a Vicryl plug and fibrin glue, as described previously [3]. The tracheal fistula was treated with a covered self-expanding tracheal stent [4].

Despite repeated endoscopic dilations, the stenosis recurred due to ischemia. Additional stenting of the anastomotic stenosis was impossible because there was only a thin tissue bridge between the pars membranacea and the (neo-)esophagus and because a stent-to-stent arrangement would most probably have led to further necrosis. The presence of the nasojunal tube caused considerable psychological stress and impaired the patient's quality of life. To enable her to recommence oral feeding, ascending colonic bypass surgery with collar side-to-side esophagocolostomy and intra-abdominal side-to-side colojejunosomy was performed 11 months after the initial operation, leaving the gastric bypass in situ. The endoscopic control showed a wide anastomosis of the esophagocolostomy, but a siphon-like reservoir at the stenotic entrance of the gastric interposition led to regurgitation (● Fig. 1, 2). We therefore performed an endoscopic closure of the esophagogastric anastomosis, with de-epithelialization of the stenotic gastric tube and sealing with bucrylate and histoacrylate. Control radiography showed adequate efflux without filling of the sealed gastric tube (● Fig. 3). She was able to resume normal oral feeding and her body weight stabilized. This endoscopic approach has not been described in the published literature before. It represents a useful alternative for the treatment of this serious clinical situation.

Endoscopy_UCTN_Code_TTT_1AO_2AI

R. Rosch, S. Truong, M. Jansen, V. Schumpelick

Department of Surgery, University Hospital Aachen, Aachen, Germany

References

- 1 Urschel JD. Esophagogastric anastomotic leaks complicating esophagectomy: a review. *Am J Surg* 1995; 169: 634–640
- 2 Atkins BZ, Shah AS, Hutcheson KA et al. Reducing hospital morbidity and mortality following esophagectomy. *Ann Thorac Surg* 2004; 78: 1170–1176
- 3 Truong S, Bohm G, Klinge U et al. Results after endoscopic treatment of postoperative upper gastrointestinal fistulas and leaks using combined Vicryl plug and fibrin glue. *Surg Endosc* 2004; 18: 1105–1108
- 4 Chauhan SS, Long JD. Management of tracheoesophageal fistulas in adults. *Curr Treat Options Gastroenterol* 2004; 7: 31–40

Bibliography

DOI 10.1055/s-2007-966609

Endoscopy 2007; 39: E270

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

R. Rosch, MD

Department of Surgery
RWTH-Aachen University
Pauwelsstraße 30
52074 Aachen
Germany

Fax: +49-241-80-82417

r.rosch@chir.rwth-aachen.de