Pyogenic granuloma of the common bile duct in a patient with choledochoduodenostomy

Fig. 1 **a** Duodenoscopic findings of a wide stoma in the side-to-side choledochoduodenostomy in the duodenum (arrow). **b** Endoscopic retrograde cholangiogram showing dilated distal common bile duct with a nonmovable, small, round filling defect (arrow).

Fig. 2 **a** Endoscopic view of the polypoid lesion in the distal common bile duct. **b** No residual lesion in the distal common bile duct after endoscopic snare polypectomy.

Pyogenic granuloma is a polypoid form of capillary hemangioma that occurs mostly on the skin and the mucous membranes of the nasal and oral cavities [1]. Al-

though pyogenic granuloma in the digestive tract excluding the oral cavity is a rarity, it can cause significant gastrointestinal bleeding, and curative resection is re-

quired for treatment in most cases [2]. We describe a case of pyogenic granuloma of the common bile duct (CBD) that was observed on endoscopy and was successfully treated by endoscopic snare polypectomy.

A 69-year-old woman was admitted to our hospital with a 1-day history of right upper quadrant pain. She had a history of cholecystectomy and CBD exploration with side-to-side choledochoduodenostomy and recurrent attacks of cholangitis. On endoscopic retrograde cholangiography (ERC) using an Olympus duodenoscope (JF-240; Olympus Optical Co., Tokyo, Japan), the CBD was cannulated with a balloon catheter through the wide stoma of the side-to-side choledochoduodenostomy (● Fig. 1 a) and the CBD visualized. A nonmovable, small, round filling defect was noted in the dilated distal CBD (● Fig. 1 b). Because of the wide stoma of the choledochoduodenostomy and the dilated CBD, it was possible to insert a forward-viewing endoscope (Olympus GIF-Q260; Olympus Optical Co.) through the stoma into the distal CBD. This showed a semipedunculated, smooth polypoid lesion that was 5 mm in diameter, hyperemic, and dark red in the distal CBD, with a tinge of blood on its surface (● Fig. 2 a). The lesion was removed by snare polypectomy without any complications (● Fig. 2 b). On the basis of histopathological analysis of the resected tissue (● Fig. 3), the lesion was diagnosed as a pyogenic granuloma. To the best of our knowledge, this is the first reported case of pyogenic granuloma in the bile duct.

Endoscopy_UCTN_Code_CCL_1AZ_2AC
Endoscopy_UCTN_Code_CCL_1AZ_2AN

Fig. 3 **a** Histological section of the resected specimen showing proliferation of small capillary-sized blood vessels and surface ulceration, accompanied by lymphocytic infiltration in the intervening stroma (hematoxylin and eosin, original magnification, $\times 200$). **b** Immunohistochemical staining for CD31 shows strong positivity in endothelial cells lining the small blood vessels (original magnification, $\times 400$).

Bibliography

DOI 10.1055/s-2007-966706

Endoscopy 2007; 39: E282–E283

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

S. H. Dong, MD, PhD

Division of Gastroenterology and Hepatology

Department of Internal Medicine

Kyung Hee University Hospital

Hoegi-Dong 1

Dongdaemun-Gu

Seoul

130-702

Korea

Fax: +82-2-9681848

gidrdong@chol.com

C. K. Lee¹, S. H. Dong¹, S. H. Jung¹, J. Y. Jang¹, H. J. Kim¹, B. H. Kim¹, Y. W. Chang¹, R. Chang¹, Y. W. Kim²

¹ Division of Gastroenterology and Hepatology, Department of Internal Medicine, Kyung Hee University, School of Medicine, Seoul, Korea

² Department of Pathology, Kyung Hee University, School of Medicine, Seoul, Korea

References

- 1 Lin RL, Janniger CK. Pyogenic granuloma. *Cutis* 2004; 74: 229–233
- 2 Kusakabe A, Kato H, Hayashi K et al. Pyogenic granuloma of the stomach successfully treated by endoscopic resection after transarterial embolization of the feeding artery. *J Gastroenterol* 2005; 40: 530–535