Implantation of an esophageal squamous cell carcinoma at the site of a percutaneous endoscopic gastrostomy

Fig. 1 Macroscopic aspect showing the skin alteration at the site of the PEG, which was regarded as granulation tissue.

Fig. 2 Gastrosopic view of the tumor at the PEG site.

Fig. 3 Metastatic tumor implantation at the site of the PEG; the tumor mass extends from the gastric lumen to the skin.

Fig. 4 Esophageal squamous cell carcinoma with distinctive lymphangiosis and hemangiosis carcinomatosa (H & E staining).

A 54-year-old man presented with a stenosing, moderately differentiated squamous cell carcinoma (SCC) of the proximal esophagus. Tumor stage was cT3–4N1M0. The patient underwent radiochemotherapy with curative intent. A percutaneous endoscopic gastrostomy (PEG) was placed using the standard pull-through method.

After 5 months the patient noticed a skin alteration at the site of the PEG, which was regarded as granulation tissue (● Fig. 1). In a routine gastroscopy 2 months later, an ulcer at the gastric site of the PEG was found (● Fig. 2), and initially diagnosed as adenocarcinoma. Computed tomography demonstrated a tumor mass along the PEG-tube (● Fig. 3). Additionally, a suspicious hypodense hepatic lesion was detected. Complete local esophageal tumor control was documented. At laparotomy, a frozen section of the liver lesion showed a poorly differentiated SCC. Therefore, palliative subtotal gastrectomy with en bloc resection of the abdominal wall was carried out (● Fig. 4).

Since the first description of PEG in 1980 [1], it has become a valuable method for nutritional support. The implantation of oropharyngeal or esophageal cancer at PEG stoma sites is a rare complication with an unknown incidence [2]. The average period of time from tube placement to metastatic spread is reported to be approximately 9 months (range 3–18 months) [3]. Length of survival following this complication is rarely reported, and varies between 2 and 28 months [4]. The mechanism of tumor spread to the PEG site is controversial. Hematogenous or lymphatic spread to a susceptible site, as well as – more likely – direct mechanical implantation at the time of the PEG placement are proposed [3–5]. To avoid mechanical tumor implantation, the contact of the PEG tube with the tumor should be minimized. In patients with bulky, stenosing tumors this can be achieved by using a sheath or overtube. Alternatively, radiologic or operative placements can be carried out.

Endoscopy_UCTN_Code_CPL_1AH_2AI

K. Volkmer¹, T. Meyer¹, M. Sailer²,
M. Fein¹

¹ Department of Surgery, University Hospital of Wuerzburg, Julius-Maximilians-University Wuerzburg, Germany

² Bethesda Hospital, Hamburg-Bergedorf, Germany

References

- 1 Gauderer MWL, Ponsky JL, Izant RJ. Gastrostomy without laparotomy: a percutaneous endoscopic technique. *J Pediatr Surg* 1980; 15: 872–875
- 2 Maccabee D, Sheppard BC. Prevention of percutaneous endoscopic gastrostomy stoma metastases in patients with active oropharyngeal malignancy. *Surg Endosc* 2003; 17: 1678
- 3 Thakore JN, Mustafa M, Suryaprasad S, Agrawal S. Percutaneous endoscopic gastrostomy associated gastric metastasis. *J Clin Gastroenterol* 2003; 37: 307–311
- 4 Ananth S, Amin M. Implantation of oral squamous cell carcinoma at the site of a percutaneous endoscopic gastrostomy: a case report. *Br J Oral Maxillofac Surg* 2002; 40: 125–130
- 5 Peghini P, Guaouguau N, Salcedo J, Al-Kawas F. Implantation metastasis after PEG: case report and review. *Gastrointest Endosc* 2000; 51: 480–482

Bibliography

DOI 10.1055/s-2007-966793

Endoscopy 2007; 39: E240–E241

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

M. Fein, MD

Department of Surgery
Zentrum Operative Medizin (ZOM)
Oberduerrbacher Strasse 6

D – 97080 Wuerzburg
Germany

Fax: +49-931-201-31049

Fein_M@chirurgie.uni-wuerzburg.de