Iron-induced esophageal ulceration

Esophageal inflammation or ulceration is a well recognized side-effect of some oral medications, especially nonsteroidal anti-inflammatory or potassium chloride tablets. A similar injury can be induced by iron but endoscopically detected lesions are rarely reported [1,2].

An 82-year-old woman presented a sudden complaint of dysphagia after ingestion of a ferrous sulphate tablet. She had chronic esophageal reflux and Raynaud's syndrome. Her physical examination was normal.

An upper endoscopy revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter, 2 cm in length, well circumscribed and associated with a slight stenosis. Below the lesion the esophageal lumen was distended and atonic, with candidiasis and a grade D peptic esophagitis (► Fig. 1, ► Video 1).

Histological evaluation revealed ulcerated esophagitis, with acute inflammatory granulation tissue containing abundant brown crystalline nonbirefringent material, confirmed as iron by Perl's staining (► Fig. 2,3). Other stainings for brown pigments and infections were negative. The patient's iron tablets were stopped and a high dose proton pump inhibitor was prescribed. The dysphagia was relieved and a repeat upper endoscopy revealed progressive resolution of the necrosis and stenosis (► Fig. 4, ► Video 2). Iron-induced esophageal erosions or ulcerations are seldom described and the differential diagnoses might be striking. Tablet debris seen at endoscopy is diagnostic but rarely detected. Histological examination reveals a typical brown crystalline material positive with Perl's staining [1,3,4]. The pathogenesis is believed to be a high local iron saturation causing a concentration-dependent absorption, resulting in the formation of reactive oxygen metabolites and mucosal injury [3]. Anatomical factors such as motility disorders or previous inflammation might predispose to the iron injury [3,4].

Treatment requires stopping the offending drug or, if iron is still necessary, changing to a liquid preparation [3,5]. Iron-induced lesions are not endoscopically specific and histological investigation is essential to the diagnosis.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AZ

M. Areia¹, R. Gradiz¹, P. Souto¹, E. Camacho¹, M. R. Silva², N. Almeida¹, A. Rosa¹, M. F. Xavier da Cunha², M. C. Leitão¹

¹ Department of Gastroenterology, Coimbra University Hospital, Coimbra, Portugal

² Department of Pathology, Coimbra University Hospital, Coimbra, Portugal

References

- 1 Abbarah TR, Fredell JE, Ellenz GB. Ulceration by oral ferrous sulfate. *JAMA* 1976; 236: 2320
- 2 Zhang ST, Wong WM, Hu WH *et al*. Esophageal injury as a result of ingestion of iron tablets. *J Gastroenterol Hepatol* 2003; 18: 466–467
- 3 Abraham SC, Yardley JH, Wu TT. Erosive injury to the upper gastrointestinal tract in patients receiving iron medication: an underrecognized entity. *Am J Surg Pathol* 1999; 23: 1241–1247
- 4 Eckstein RP, Symons P. Iron tablets cause histopathologically distinctive lesions in mucosal biopsies of the stomach and esophagus. *Pathology* 1996; 28: 142–145
- 5 Haig A, Driman DK. Iron-induced mucosal injury to the upper gastrointestinal tract. *Histopathology* 2006; 48: 808–812

Bibliography

DOI 10.1055/s-2007-966820

Endoscopy 2007; 39: E326

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

M. Areia, MD

Department of Gastroenterology
Coimbra University Hospital
3000-075 Coimbra
Portugal
Fax: +351-239-701517
miguel.areia@netcabo.pt

Video 1

Upper endoscopy at day 1 revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter and a distended atonic esophagus.

Video 2

Upper endoscopy at day 8 showing the progressive resolution of the lesion with a circumferential brown-pigmented lesion with no necrosis or stenosis.

Fig. 1 Upper endoscopy at day 1 revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter.

Fig. 2 Granulation tissue from the area with ulcerated esophagitis containing abundant brown crystalline material (hematoxylin and eosin, ×400).

Fig. 3 Positive blue staining of the crystalline material with Perl's iron stain (×400).

Fig. 4 Upper endoscopy at day 8 showing the progressive resolution of the lesion, with a brown pigmentation and no necrosis.