Colonic taeniasis

Fig. 1 Flat, white, mobile worm emerging from ileocecal valve.

Fig. 2 Extension of worm through transverse colon.

Fig. 3 *Taenia* spp. eggs. Original magnification $\times 40$.

A 52-year-old Tibetan male was referred for evaluation of rectal bleeding and intermittent mid abdominal pain. He reported frequent loose stools but no associated fever or weight loss. Results from physical exam were within normal limits. Hemoglobin level was 13.1 g/dL (normal range 14–18 g/dL), and white-cell count was 5100/mm³ (normal 4800–10800/mm³) with 13% eosinophils (normal 1%–4%). Colonoscopy revealed small

nonbleeding internal hemorrhoids, and a white ribbon-like structure protruding from the ileocecal valve (● **Fig. 1**) and extending throughout the entire colon (● **Fig. 2**). A segment of the mobile worm was successfully removed using biopsy forceps, and histologic evaluation demonstrated parasite eggs morphologically consistent with *Taenia* spp. (● **Fig. 3**, ● **Video 1**).

Taenia solium and *T. saginata* are the most common species of tapeworms infecting humans. Although the worm normally resides in the small intestine it can grow up to 25 m and extend into the large intestine. The patient was treated successfully with one dose of praziquantel.

Endoscopy_UCTN_Code_CCL_1AD_2AZ

F. Francois, I. Cho

New York University School of Medicine,
New York, USA

Bibliography

DOI 10.1055/s-2007-966965

Endoscopy 2008; 40: E28

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

F. Francois, MD

Division of Gastroenterology (11132N)
VA New York Harbor Healthcare System
423 East 23rd Street
New York
NY 10010
USA

Fax: +1-212-951-3481

fritz.francois@med.nyu.edu

Video 1

Mobile tapeworm moving in colon and grabbed with forceps.