Embolic complications associated with endoscopic injection of cyanoacrylate for bleeding duodenal ulcer

Fig. 1 Ulcer on the anterosuperior bulb wall, on 24-hour second-look endoscopy: **a** with visible vessel; **b** after second NB2C treatment.

An 87-year-old man with severe cardiac disease, on treatment with an antiplatelet agent, was admitted to our department with acute melena, in a hemodynamically unstable state, and with a hemoglobin level of 6 mg/dL. Emergency endoscopy revealed an extensive ulcer in the anterosuperior bulb wall with pulsatile bleeding. This was initially injected with epinephrine and fibrin glue (as it was a deep ulcerated lesion). When this failed to stop the bleeding, we used a 1:0.6 mixture of *N*-butyl-2-cyanoacrylate (NB2C; Histoacryl) and lipiodol, which did stop it. At second-look endoscopy, a large pulsatile vessel was still present (● Fig. 1 a), which was permanently occluded after a second NB2C application (● Fig. 1 b). Five days later, the patient developed febrile peaks (40°C) without complaints but with leukocytosis and a five-fold increase in levels of aminotransferases, amylase, and lipase. A thoracoabdominal computed tomography (CT) scan showed linear opacification of the common hepatic artery

Fig. 2 Abdominal CT scan showing radiopaque material in **a** the common hepatic artery and **b** its right branch and some splenic branches. Multiple areas of splenic infarctions are visible in **b**.

(● Fig. 2 a), its right branch, and some splenic branches (● Fig. 2 b), with a heterogeneous area in the spleen (● Fig. 2 b) and in the pancreatic head (● Fig. 3) highly suggestive of infarction lesions. The patient started treatment with an intravenous broad-spectrum antibiotic, along with nutritional support measures, and the liver test parameters improved considerably. Blood cultures failed to isolate any bacterial strain. The patient was discharged on day 15. Six-month imaging follow-up showed remarkable improvement. Bleeding peptic ulcer is still the main cause of upper gastrointestinal hemorrhage [1]. Several endoscopic hemostatic methods of similar efficacy are currently available [1]. The use of NB2C, a successful and well-established substance used in variceal hemorrhage, is still controversial in the context of bleeding peptic ulcer

Fig. 3 Abdominal CT scan showing a heterogeneous area in the pancreatic head.

[2,3]. Encouraging results have shown it to have good hemostatic efficacy when conventional endoscopic techniques have failed to control bleeding [2,3]. However, it has been associated with severe embolization with infarction [2,4,5]. The present case highlights a potential adverse effect of cyanoacrylate use.

Endoscopy_UCTN_Code_CPL_1AH_2AC

P. Peixoto, P. Ministro, A. Sadio, A. Castanheira, E. Cancela, R. Araújo, A. Silva, A. Caldas
Gastroenterology Department,
São Teotónio Hospital, Viseu, Portugal

References

- 1 Huang CS, Lichtenstein DR. Nonvariceal upper gastrointestinal bleeding. *Gastroenterol Clin North Am* 2003; 32: 1053–1078
- 2 Repici A, et al. Adrenaline plus cyanoacrylate injection for treatment of bleeding peptic ulcers after failure of conventional endoscopic haemostasis. *Dig Liver Dis* 2002; 34: 349–355
- 3 Kok KY, Kum CK, Goh PM et al. Endoscopic hemostasis of upper gastrointestinal bleeding with histoacryl: last resort before surgery. *Endoscopy* 1996; 28: 256–258
- 4 Lee GH, Kim JH, Lee KJ et al. Life threatening intraabdominal arterial embolization after histoacryl injection for bleeding gastric ulcer. *Endoscopy* 2000; 32: 422–424
- 5 Vallieres E, Jamieson C, Haber GB et al. Pancreatoduodenal necrosis after endoscopic injection of cyanoacrylate to treat a bleeding duodenal ulcer: a case report. *Surgery* 1989; 106: 901–903

Bibliography

DOI 10.1055/s-2007-995695

Endoscopy 2008; 40: E126

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

P. C. P. Peixoto, MD

Gastroenterology Department, São Teotónio Hospital, Av. Rei D. Duarte, 3504 509 Viseu, Portugal
Fax: +351-232-420591
paulacristinapeixoto@iol.pt