Intestinal graft-versus-host-disease staging by video capsule endoscopy

Fig. 1 Endoscopic appearance of severe diffuse hemorrhagic inflammation of the sigmoid colon.

Fig. 2 Histological section (H&E staining) showing autolytic mucosa of the small bowel with acute fibrinous and chronic inflammation. Pronounced submucosal edema and dilated capillary vessels due to graft-versus-host disease.

Graft-versus-host disease (GvHD) is a leading cause of morbidity and mortality in patients who have undergone allogeneic bone marrow or peripheral blood stem cell transplantation (BMT/PBSCT) [1]. About 20–50% of patients who have received BMT/PBSCT develop symptoms of acute GvHD [2]. GvHD most often affects the skin, liver, and gastrointestinal tract. In severe intestinal GvHD broad ulcerative lesions develop, leading to diarrhea, malabsorption, intestinal hemorrhage, and sepsis [3]. The gold standard in the diagnosis of intestinal GvHD is upper and lower gastrointestinal endoscopy with histological validation [4].

A 57-year-old woman with acute myeloid leukemia who had recently received a PBSCT from an HLA-identical sibling suffered severe hemorrhagic diarrhea. The symptoms started 12 days after the transplant. The diagnosis of GvHD was histologically and endoscopically confirmed by sigmoidoscopy (CF-H180AI/L; Olympus Co. Ltd., Tokyo, Japan) (● **Fig. 1**). Esophagogastroduodenoscopy did not reveal relevant pathologic findings.

Since the patient's condition rapidly deteriorated, a total colectomy was discussed as a last therapeutic option following the failure of several immunosuppressive drug regimens including corticosteroids, cyclosporine, mycophenolate mofetil, pentostatin, infliximab, and antithymocyte globulin. Video capsule endoscopy (PillCam SB; Given Imaging Ltd., Yoq-

neam, Israel) carried out to evaluate small-bowel involvement in the GvHD revealed continuous severe hemorrhagic inflammation of the entire small intestine starting from the proximal jejunum (● **Video 1**) and ending in the terminal ileum (● **Video 2**).

Video capsule endoscopy proved to be a successful minimally invasive diagnostic method, accurately visualizing the involvement of the small intestine in severe GvHD, obviating the necessity for diagnostic surgical exploration in a clinically unstable patient.

The patient died of multiorgan failure due to GvHD. Histological analysis of the intestine confirmed severe GvHD involving the entire small intestine and colon (● **Fig. 2**). We conclude that video capsule endoscopy is suitable for staging intestinal GvHD especially in those patients unable to tolerate invasive diagnostic measures such as double-balloon enteroscopy or surgery.

Video 1

Sequence of video capsule endoscopy of the proximal jejunum showing severe hemorrhagic mucosal inflammation and broad ablation of the mucosal surface.

Video 2

Sequence of video capsule endoscopy of the ileum with large amounts of intraluminal blood due to severe graft-versus-host disease.

Endoscopy_UCTN_Code_CCL_1AC_2AD

T. Meister¹, H. Heinzow¹, G. Bisping², M. Stelljes², B. Schulte³, W. E. Berdel², J. Kienast², W. Domschke¹, H. Ullerich¹

¹ Department of Medicine B, University of Münster, Münster, Germany

² Department of Medicine A, University of Münster, Münster, Germany

³ Gerhard-Domagk-Institut of Pathology, University of Münster, Münster, Germany

References

- 1 Vogelsang GB, Lee L, Bensen-Kennedy DM. Pathogenesis and treatment of graft-versus-host disease after bone marrow transplant. *Annu Rev Med* 2003; 54: 29–52
- 2 Remberger M, Persson U, Hauzenberger D et al. An association between human leucocyte antigen alleles and acute and chronic graft-versus-host disease after allogeneic haematopoietic stem cell transplantation. *Br J Haematol* 2002; 119: 751–759
- 3 Martin PJ, McDonald GB, Sanders JE et al. Increasingly frequent diagnosis of acute gastrointestinal graft-versus-host disease after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2004; 10: 320–327
- 4 Cruz-Correa M, Poonawala A, Abraham SC et al. Endoscopic findings predict the histologic diagnosis in gastrointestinal graft-versus-host disease. *Endoscopy* 2002; 34: 808–813

Bibliography

DOI 10.1055/s-2007-995767

Endoscopy 2008; 40: E144

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

H. Ullerich, MD

Department of Medicine B

University of Münster

Albert-Schweitzer-Str. 33

48149 Münster

Germany

Fax: +49-251-8347576

ullerih@mednet.uni-muenster.de