S. PROCH, Y. MEI, J. M. RIVERA VILLANUEVA, Y. LU, A. KARPOV, M. BALLAUFF,*
 R. KEMPE* (UNIVERSITÄT BAYREUTH AND BASF AG, LUDWIGSHAFEN, GERMANY)
 Suzuki- and Heck-Type Cross-Coupling with Palladium Nanoparticles Immobilized on Spherical Polyelectrolyte
 Brushes
Adv. Synth. Catal. **2008**, 350, 493-500.

Suzuki–Miyaura Coupling Reaction Using Palladium Nanoparticles Immobilized on SPB

Category

Polymer-Supported
Synthesis

Key words

palladium
 Suzuki–Miyaura
 coupling
 Heck reaction
 nanoparticles

SYNFACT
of the month

Significance: Palladium nanoparticles immobilized on spherical polyelectrolyte brushes (Pd@SPB) were synthesized and found to promote the Suzuki–Miyaura cross-coupling reaction. Thus, the spherical polyelectrolyte brushes having the (2-methylpropenyloxyethyl)trimethylammonium cation were prepared according to the reported procedures (Y. Mei, G. Sharma, Y. Lu, M. Ballauff *Langmuir* **2005**, 21, 12229). The reaction of SPB with Na₂PdCl₄ followed by reduction with NaBH₄ gave the palladium nanoparticles immobilized in the SPB matrix (Pd@SPB). The Suzuki–Miyaura coupling reaction of aryl halides with phenylboronic acid was carried out in the presence of Pd@SPB (0.09 mol% Pd) in H₂O–THF to give the corresponding biphenyls in good yield (14 examples, 50–94% yield).

Comment: Recently, the authors have reported the preparation of platinum nanoparticles immobilized on SPB (Pt@SPB), which was found to be an effective catalyst for the reduction of *p*-nitrophenol with NaBH₄. The present report described that the palladium nanoparticles (Pd@SPB) exhibited good activity for the Suzuki–Miyaura coupling reaction. This catalyst was recovered and reused for four times without any loss of catalytic activity. Heck reaction using the palladium nanoparticles (Pd@SPB) was also described.

SYNFACTS Contributors: Yasuhiro Uozumi, Toshimasa Suzuka
 Synfacts 2008, 5, 0541-0541 Published online: 23.04.2008
 DOI: 10.1055/s-2008-1072559; Reg-No.: Y03908SF