Perforation of the proximal esophagus treated with factor XIII

A 16-year-old girl presented to us with a perforated esophagus after accidental ingestion of a toothbrush. The referring physician had already tried to retrieve the brush endoscopically without success.

Esophagogastroduodenoscopy (EGD) revealed the head of the brush impacted in the rear wall of the esophagus right under the upper esophageal sphincter. Oral retrieval of the brush was not possible so it was moved to the stomach first and then retrieved using a loop around the neck of the brush and an overtube. Inspection of the esophagus showed a near total perforation of the rear wall under the upper esophageal sphincter measuring approximately 1.5×2 cm (● Fig. 1).

The patient was treated conservatively with antibiotics and nil by mouth. Two days after the primary intervention a second EGD was performed to inspect the perforation site. An area measuring 1×2 cm covered with fibrin was found. This area was treated with factor XIII in all four quadrants of the lesion up to a total of 125 IU (2 ml) (● Fig. 2).

Coagulation factor XIII was first described by Laki and Lóránd in 1948 as “fibrin-stabilizing factor” [1]. Factor XIII is the last enzyme (plasma protransglutaminase) in physiological coagulation and is also important for proliferation and immigration of fibroblasts in wound healing. Factor XIII also inhibits the migration of macrophages [2]. Factor XIII has been used in the therapy of ulceration due to pressure, large burns, sepsis, and acute liver disorders [3].

On the 9th day after primary intervention only a very small lesion barely covered with fibrin without any suspicious signs was seen (● Fig. 3). The diet was increased without complications and the patient was discharged. Eight weeks after the incident EGD showed a completely healed wound at the site of the rupture (● Fig. 4).

Endoscopy_UCTN_Code_TTT_1AO_2AI

Fig. 1 Site of the perforation of the rear esophageal wall directly underneath the upper esophageal sphincter.

Fig. 2 Injection of factor XIII in all four quadrants of the lesion.

Fig. 3 Check-up on the ninth day after primary intervention. Slight fibrinous lesions can still be seen.

Fig. 4 Last esophagogastroduodenoscopy after eight weeks. Lesion is completely healed.

C. Knorr¹, R. T. Carbon¹, S. Albrecht², M. Richter³, H. Köhler³

¹ Department of Pediatric Surgery, Children's Hospital, University Medical Center Erlangen of the Friedrich-Alexander-University Erlangen-Nürnberg, Erlangen, Germany

² Department of Anaesthesia, Children's Hospital, University Medical Center Erlangen of the Friedrich-Alexander-University Erlangen-Nürnberg, Erlangen, Germany

³ Division of Gastroenterology, Children's Hospital, University Medical Center Erlangen of the Friedrich-Alexander-University Erlangen-Nürnberg, Erlangen, Germany

References

- 1 Laki K, Lóránd L. On the solubility of fibrin clots. *Science* 1948; 108: 280
- 2 Beck E, Duckert F, Ernst M. The influence of fibrin stabilizing factor on the growth of fibroblasts in vitro and wound healing. *Thromb Diath Haemorrh* 1961; 6: 485–491
- 3 Canonico S. The use of human fibrin glue in the surgical operations. *Acta Biomed* 2003; 74 (Suppl 2): 21–25

Bibliography

DOI 10.1055/s-2008-1077422

Endoscopy 2008; 40: E197

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

C. Knorr, MD

Department of Pediatric Surgery
Universitätsklinikum Erlangen der Friedrich-Alexander-Universität Erlangen-Nürnberg
Krankenhausstraße 12
91052 Erlangen
Germany
christian.knorr@uk-erlangen.de