Capsule endoscopy with retention of the capsule in a duodenal diverticulum

The most important complication of capsule endoscopy is retention of the capsule. Among patients with obscure gastrointestinal bleeding, capsule retention was reported in 1.5%; in patients with suspected Crohn's disease or suspected stenosis of the small bowel the risk of capsule retention seems to be considerably higher (5% and 21% respectively) [1,2]. To our best knowledge this is the first case of capsule retention in a duodenal diverticulum.

A 74-year-old woman was admitted to hospital for evaluation of microcytic anemia. Gastroscopy and colonoscopy were normal. Capsule endoscopy, using the M2A capsule (Given Imaging Ltd., Yoqneam, Israel [3]), demonstrated some diverticula shortly after passage of the pylorus. Further images were not evaluable. Three weeks afterwards the patient was asymptomatic but had still not excreted the capsule. A plain film of the abdomen demonstrated the capsule superimposed on the epigastrium and was otherwise normal (● Fig. 1). A small-bowel radiograph with water-soluble contrast medium showed the capsule in a duodenal diverticulum and ruled out obstruction of the small bowel (● Fig. 2). Gastroscopy was performed and the capsule successfully extracted from a large juxtapapillary diverticulum using the Roth retrieval net (● Fig. 3).

Most patients with capsule retention are asymptomatic. However, a very few cases of symptomatic bowel obstruction requiring surgical or endoscopic removal of the impacted capsule have been reported [4,5]. This is the first case of capsule retention in a duodenal diverticulum and successful endoscopic removal of the impacted capsule. We suggest that, even in asymptomatic patients, capsules that are retained in intestinal diverticula and are not excreted within a period of about 3 weeks should be removed by gastroscopy or enteroscopy in order to prevent complications such as diverticulitis, perforation, or pancreatitis.

We conclude that capsule retention in a duodenal diverticulum is a rare complica-

Fig. 1 Plain film anteroposterior abdominal radiograph: the capsule is superimposed on the epigastrium.

Fig. 2 Small-bowel radiograph with water-soluble contrast medium: the capsule is retained in a large duodenal diverticulum.

Fig. 3 The capsule was captured in a Roth retrieval net (white arrow). The yellow arrow marks the Teflon-coated tube of the net.

tion of capsule endoscopy. In our patient endoscopic removal of the impacted capsule using a retrieval net was successful.

Endoscopy_UCTN_Code_CPL_1AI_2AB

P. Ordubadi¹, B. Blaha¹, A. Schmid²,
W. Krampla³, W. Hinterberger²,
M. Gschwantler¹

¹ Fourth Department of Internal Medicine,
Wilhelminenspital, Vienna, Austria

² Second Department of Internal
Medicine, Donauspital, Vienna, Austria

³ Department of Radiology, Donauspital,
Vienna, Austria

References

- 1 Lewis B. How to prevent endoscopic capsule retention. *Endoscopy* 2005; 37: 852 – 856
- 2 Lewis B. Capsule endoscopy – transit abnormalities. *Gastrointest Endosc Clin N Am* 2006; 16: 221 – 228
- 3 Mata A, Llach J, Bordas JM. Wireless capsule endoscopy. *World J Gastroenterol* 2008; 13: 1969 – 1971
- 4 Sears DM, Avots-Avotins A, Culp K, Gavin MW. Frequency and clinical outcome of capsule retention during capsule endoscopy for GI bleeding of obscure origin. *Gastrointest Endosc* 2004; 60: 822 – 827
- 5 Hara AK, Leighton JA, Sharma VK, Fleischer DE. Small bowel: preliminary comparison of capsule endoscopy with barium study and CT. *Radiology* 2004; 230: 260 – 265

Bibliography

DOI 10.1055/s-2008-1077563

Endoscopy 2008; 40: E247 – E248

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

M. Gschwantler, MD

Wilhelminenspital
4th Department of Internal Medicine
Montleartstraße 37
1160 Vienna
Austria

Fax: +43-1-491502409

Michael.Gschwantler@wienkav.at