Minimally invasive treatment of benign complete stenosis of colorectal anastomosis

Fig. 1 Barium enema performed through the ileostomy.

Fig. 3 Drainage catheter placed through the neo-anastomosis.

Fig. 5 Endoscopic check after removal of the self-expanding metal stent.

Fig. 2 Needle puncture under endoscopic vision.

Fig. 4 Fully covered and removable self-expanding metal stent in place.

Fig. 6 Endoscopic check after 24 months.

The most frequent complications of colorectal surgery are leaks and stenosis, which occur with a frequency of 0.3%–30% [1,2]. Strictures are commonly treated with conservative treatments such as balloon dilatation, endoscopic incision, and placement of self-expanding metal stent (SEMS) [3–5]. Two patients with complete colorectal anastomotic closure were successfully treated with an endoscopic–radiologic rendezvous procedure. A 27-year-old woman who had undergone low anterior resection due to cancer of the rectum, and a 69-year-old man who had undergone left colonic resection because of acute diverticulitis were referred to our department. Both patients had a loop ileostomy. Endoscopic and radiologic follow-up before recanalization showed complete anastomotic stric-

ture (● **Fig. 1**) at 8 cm and 16 cm from the anal verge, respectively. In both patients malignancy was excluded by means of multiple biopsies. Both patients were treated as follows. An incision of the fibrotic stenosis of the anastomosis was made through the rectum using a needle normally used for transjugular intrahepatic portosystemic shunt (TIPS). Direct vision of the puncture place was assured by an antegrade colonoscopy from the ileostomy (● **Fig. 2**). A guide wire was passed through and a drainage catheter was placed. The drainage catheter was kept in place for 3 days (● **Fig. 3**). Following two balloon dilatations, a fully covered and removable stent was placed (● **Fig. 4**). After 60 days, the stent was re-

moved (● **Fig. 5**). In one patient the stent gave mild tenesmus, which was alleviated by analgesic drugs.

In both patients the neo-anastomosis (● **Fig. 6**) was still wide open and functioning well after 24 months. Even in such difficult cases, in which the use of surgery seemed inevitable and furthermore not free of risks, the cooperation between endoscopist and radiologist made conservative treatment possible. The placement of removable SEMS in order to maintain an adequate lumen and to eliminate any possible leak was demonstrated to be a safe and efficient technique.

Endoscopy_UCTN_Code_TTT_1AQ_2AF

G. Donatelli¹, V. Ceci¹, F. Cereatti¹,
A. Bruni², F. M. Salvatori², S. Minervini¹,
F. Fiocca¹

¹ Department of General Surgery “Paride Stefanini”, Rome, Italy

² Department of Radiological Sciences, University of Rome “La Sapienza”, Rome, Italy

References

- 1 Platell C, Barwood N, Dorfmann G, Makin G. The incidence of anastomotic leaks in patients undergoing colorectal surgery. *Colorectal Dis* 2007; 9: 71 – 79
- 2 McKee R, Pricolo VE. Stapled revision of complete colorectal anastomotic obstruction. *Am J Surg* 2008; 195: 526 – 527
- 3 Forshaw MJ, Maphosa G, Sankararajah D et al. Endoscopic alternatives in managing anastomotic strictures of the colon and the rectum. *Tech Coloproctol* 2006; 10: 21 – 27
- 4 Rees JRE, Carney L, Gill TS, Dixon AR. Management of recurrent anastomotic stricture and iatrogenic stenosis by circular stapler. *Dis Colon Rectum* 2004; 47: 944 – 947
- 5 Forshaw MJ, Sankararajah D, Stewart M, Parker MC. Use of self-expanding metallic stents for benign colorectal disease. *Colorectal Dis* 2006; 8: 102 – 111

Bibliography

DOI 10.1055/s-2008-1077660

Endoscopy 2008; 40: E263 – E264

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

F. Fiocca, MD

Department of General Surgery “Paride Stefanini”
Viale del Policlinico 155

00161, Rome

Italy

Fax: +39-06-4940532

fausto.fiocca@uniroma1.it