Successful biliary stenting and sphincterotomy using an ultrathin forward-viewing endoscope

Fig. 1 Endoscopic retrograde cholangiopancreatography (ERCP) image showing common bile duct (CBD) dilatation with multiple filling defects.

Some complications of endoscopic retrograde cholangiopancreatography (ERCP) including cardiovascular stress may be related to the use of large-caliber endoscopes [1,2]. Procedures with the ultrathin endoscope have been shown to be better tolerated and require less sedation than conventional esophagogastroduodenoscopy [3]. Furthermore, transnasal ERCP utilizing an ultrathin endoscope has been shown to cause less cardiovascular stress [4]. ERCP using the ultrathin endoscope seems to be a promising alternative method, especially for those with a high risk of cardiovascular events. However, therapeutic ERCP procedures, such as biliary stenting or sphincterotomy, via an ultrathin forward-viewing endoscope have not been currently reported. We present here a first case report of successful biliary stenting and sphincterotomy using an ultrathin forward-viewing endoscope.

A 57-year-old man presented at our institution with acute cholangitis and obstructive jaundice. Cardiopulmonary assessment revealed tachypnea and inspiratory crackles over the lower lung fields. He underwent emergency ERCP for biliary decompression. A standard side-viewing duodenoscope was utilized at first. However, after repeated failed attempts at biliary cannulation, we decided to use a small-caliber forward-viewing endoscope (EG-530N; Fujinon Corporation, Saitama, Japan) transorally. The major papilla was seen after retroflexion of the endoscopic tip at the second portion

Fig. 2 Accessories used in this patient: biliary drainage tube of our own design with outer diameter 5 Fr and length 7 cm (a), needle knife (b), and retrieval basket (c).

Fig. 3 Two self-designed biliary drainage tubes were inserted subsequently to attain adequate drainage.

of the duodenum. The ERCP image showed common bile duct (CBD) dilatation with multiple filling defects (● Fig. 1). Two drainage tubes of our own design (7 cm length, outer diameter 5 Fr; ● Fig. 2) were inserted subsequently to attain adequate biliary drainage (● Fig. 3). The patient was afebrile after biliary stenting. On the fifth hospital day, a Huibregtse single-lumen needle knife (HPC-2; Cook Endoscopy, North Carolina, USA; ● Fig. 2) was used for precut

Fig. 4 a Endoscopic image showing precut sphincterotomy with a needle knife. b Removal of CBD stones with a retrieval basket.

sphincterotomy (● Fig. 4a). The drainage tubes were removed and a retrieval basket (FG-18Q-1; Olympus Medical Systems Corporation, Tokyo, Japan; ● Fig. 2) was used to remove CBD stones (● Fig. 4b). The clinical course was uneventful and the patient was discharged 3 days after sphincterotomy.

Endoscopy_UCTN_Code_TTT_1AR_2AC

Endoscopy_UCTN_Code_TTT_1AR_2AK

S.-L. Yan¹, C.-H. Chen², Y.-H. Yeh¹, S.-K. Yueh²

¹ Division of Gastroenterology, Department of Internal Medicine, Chang Bing Show Chwan Memorial Hospital, Taiwan, Republic of China

² Division of Gastroenterology, Department of Internal Medicine, Show Chwan Memorial Hospital, Taiwan, Republic of China

References

- 1 Lieberman DA, Wuerker CK, Katon RM. Cardiopulmonary risk of esophagogastroduodenoscopy. Role of endoscope diameter and systemic sedation. *Gastroenterology* 1985; 88: 468–472
- 2 Johnston SD, McKenna A, Tham TC. Silent myocardial ischaemia during endoscopic retrograde cholangiopancreatography. *Endoscopy* 2003; 35: 1039–1042
- 3 Preiss C, Charton JP, Schumacher B, Neuhaus H. A randomized trial of unsedated transnasal small-caliber esophagogastroduodenoscopy (EGD) versus peroral small-caliber EGD versus conventional EGD. *Endoscopy* 2003; 35: 641–646
- 4 Mori A, Asano T, Maruyama T et al. Transnasal ERCP/ENBD using an ultrathin esophagogastroduodenoscope. *J Gastroenterol* 2006; 41: 1237–1238

Bibliography

DOI 10.1055/s-2008-1077773

Endoscopy 2009; 41: E59–E60

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

Y.-H. Yeh, MD

Division of Gastroenterology,
Department of Internal Medicine
Chang Bing Show Chwan Memorial Hospital
No 6, Lugong Rd, Lugang Township
Changhua County 505
Taiwan, ROC
Fax: +886-4-7233190
yslesmu@hotmail.com