Letter to the Editor

Open Access

Motor grading of elbow flexion – is Medical Research Council grading good enough?

Praveen Bhardwaj*¹ and Navin Bhardwaj*²

Address: ¹Hand & Microsurgery, Ganga Hospital, Coimbatore, India and ²Consultant Physiotherapist, Badli, New Delhi- 110042, India

Email: Praveen Bhardwaj* - drpb12@yahoo.co.in; Navin Bhardwaj* - praveenbhardwaj1@rediffmail.com

* Corresponding authors

Published: 13 May 2009

Received: 1 November 2008

Accepted: 13 May 2009

Journal of Brachial Plexus and Peripheral Nerve Injury 2009, **4**:3 doi:10.1186/1749-7221-4-3

This article is available from: <http://www.jbppni.com/content/4/1/3>

© 2009 Bhardwaj and Bhardwaj; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Restoration of elbow flexion is top priority in reconstruction following brachial plexus injury. Medical Research Council (MRC) Grading is the most commonly used scale to grade muscle power. Though simple to use, it has several limitations. Each grade represents a very wide range and hence precludes accurate assessment of function and outcome following a given procedure. Wide range of Grade 4 is most worrisome. Definitely all grade 4 labeled can not equate to good functional results. With most of the nerve transfer procedures described now claiming grade 4 recoveries in more than 80% of the reported cases a need for more detailed and accurate assessment of this grade is greatly felt. A modified MRC grading system is described which is comprehensive and easy to use.

Recovery of elbow flexion is considered as top priority in reconstruction following brachial plexus injury, hence lot of procedures have been described to restore it [1-4]. Nerve transfer is the most preferred method unless the patient presents very late. To assess the recovery of elbow flexion Medical Research Council Grading has been most commonly used worldwide. Serious limitations of MRC grading system have been expressed by many authors [5,6] but it continues to be in use because of its simplicity. Many modifications have been used by various authors [5-9] but none are widely used. We believe that for any grading system to be widely acceptable it need to be a modification of the existing MRC grading system as this has been fed into at least three generations of residents and all are very used to and comfortable using this scale, may be at cost of accuracy. In addition, the grading system has to be comprehensive, easy to use and reproducible.

We have been using a modified MRC grading scale to assess the recovery of elbow flexion following nerve transfer in our patients (Table 1). This is a very simple grading system which basically is an elaborated MRC scale. The grade 0 and 1 remains same. Division of Grade 2 & 3 is influenced by the active motion scale described by Curtis et al [9]. Grade 2 has been subdivided into three subdivisions; A, B & C based on the range of motion with gravity eliminated. Grade 3 has been similarly subdivided depending on the range of motion against gravity. The subdivision of Grade 4 is based on the patient's ability to lift the weight through full range of flexion on a biceps curl machine, with weights in 0.5 Kg increments, a commonly used machine in physiotherapy departments and gymnasiums to strengthen the biceps. Grade 4 has three subdivisions; A- if the patient is able to lift less than 30% weight of the normal side; B- if he is able to lift 30–60%

Table 1: Modified Medical Research Council system of grading elbow flexion

Grade	Subdivision	Description
0	-	No contraction
1	-	Perceptible contraction in the muscle but no movement
2		Gravity Eliminated
	A	Motion less than or equal to half range
	B	Motion more than half range
	C	Full range of motion
3		Against Gravity
	A	Motion less than or equal to half range
	B	Motion more than half range
	C	Full range of motion
4		Motion Against Resistance
	A	Able to lift less than 30% weight of the normal side through full range
	B	Able to lift 30–60% weight of the normal side through full range
	C	Able to lift more than 60% weight of the normal side through full range
5		Normal strength

weight of the normal side; and C- if he is able to lift more than 60% weight of the normal side. Grade 5 will mean normal strength i.e. able to lift the same amount of weight as the normal side.

We have found this scale very easy to use and reproducible. It has several advantages; by subdividing grade 2 and 3 we are able to track the recovery better, this not only helps the treating team to assess the recovery but also gives lot of confidence to the patient by knowing that he is improving. This is an important part for any nerve injury management as the nerve recovery takes very long time, may be months before patient migrated from grade 2 to grade 3, in which period patient may be very anxious and doubtful. By further subdividing these two grades we can actually show the progressive recovery to the patient and boost his confidence. Also, it will allow comparing the rate of recovery following different nerve transfer techniques.

Grade 4 is the least defined of all the grades in MRC system because of its widespread range [5,6]. If a patient is able to lift 1 kg weight he is labeled as grade 4 and another patient who is able to lift 20 kg is also grade 4. The difference between these two is phenomenal, both from functional point of view and for assessment of the final outcome following a surgical procedure. The data of the experimental study conducted by MacAvoy and Green [5] showed that grade 4 alone represents 96% of the entire spectrum of potential strength of the particular muscle and hence demands subdivisions for more precise assessment and documentation. They suggested that gross sub-

jective estimate of strength as percentage of the normal side would be more useful than the MRC scale. But we believe that it will be too subjective and preclude standard and reproducible assessment.

Subdividing the grade 4 into three subgroups based on the percentage of weight a person could lift on a biceps curl machine is definitely useful. It allows us to objectively assess and document the recovery and the final functional outcome. With most of the nerve transfer procedures described now claiming grade 4 recoveries in more than 80% of the reported cases [2-4,10-13] it is high time we get more detailed assessment of this grade lest we shall be comparing 'apples with oranges'. Definitely all grade 4 labeled can not equate to good functional results. This subdivision shall give us clearer picture of the functional recovery and dictate the supremacy of one procedure over the other. A grading system similar to this may be applied to other muscle assessment as well.

Abbreviations

MRC Grade: Medical Research Council Grade

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PB: Conceived the idea, collected the relevant literature, designed the modified classification and wrote the article.

NB: Designed the modified classification and used it in the clinical practice.

Both the authors have read the final version of the article and agreed to its content.

References

1. Chuang DC, Epstein MD, Yeh MC, Wei FC: **Functional restoration of elbow flexion in brachial plexus injuries: Results in 167 patients.** *J Hand Surg [Am]* 1993, **18(2)**:285-291.
2. Oberlin C, Beal D, Leechavengvongs S, Salon A, Dauge MC, Sarcy JJ: **Nerve transfer to biceps muscle using a part of ulnar nerve for C5-C6 avulsion of the brachial plexus: Anatomical study and report of four cases.** *J Hand Surg [Am]* 1994, **19(2)**:232-237.
3. Mackinnon SE, Novak CB, Myckatyn TM: **Results of reinnervation of the biceps and brachialis muscles with a double fascicular transfer for elbow flexion.** *J Hand Surg [Am]* 2005, **30(5)**:978-985.
4. Sungpet A, Suphachatwong C, Kawinwonggowit V: **One fascicle median nerve transfer to biceps muscle in C5 and C6 root avulsions of brachial plexus injury.** *Microsurgery* 2003, **23**:10-13.
5. MacAvoy MC, Green DP: **Critical reappraisal of medical research council muscle testing for elbow flexion.** *J Hand Surg [Am]* 2007, **32(2)**:149-153.
6. James MA: **Use of the medical research council muscle strength grading system in the upper extremity.** *J Hand Surg [Am]* 2007, **32(2)**:154-156.
7. Waikukul S, Wongtragul S, Vanadurongwan V: **Restoration of elbow flexion in brachial plexus avulsion injury: comparing spinal accessory nerve transfer with intercostal nerve transfer.** *J Hand Surg [Am]* 1999, **24(3)**:571-577.
8. Chen L, Gu Y-D, Hu S-N, Xu J-G, Xu L, Fu Y: **Contralateral C7 transfer for the treatment of brachial plexus root avulsions in children - a report of 12 cases.** *J Hand Surg [Am]* 2007, **32(1)**:96-103.
9. Curtis C, Stephens D, Clarke HW, Andrews D: **The active movement scale: an evaluation tool for infants with obstetrical brachial plexus palsy.** *J Hand Surg [Am]* 2002, **27(3)**:470-478.
10. Leechavengvongs S, Witoonchart K, Uerpaiojkit C, Thuvasethakul P, Malungpaishrope K: **Combined nerve transfer for C5 and C6 brachial plexus avulsion injury.** *J Hand Surg [Am]* 2006, **31(2)**:183-189.
11. Bertelli JA, Ghizoni MF: **Reconstruction of C5 and C6 brachial plexus avulsion injury by multiple nerve transfers: spinal accessory to suprascapular, ulnar fascicle to biceps branch and triceps long or lateral head branch to axillary nerve.** *J Hand Surg* 2004, **29A**:131-139.
12. Phillippe A, Carlos DL, Jean-Yves B, Sebastien D, Oberlin C: **Preliminary results of double nerve transfer to restore elbow flexion in upper type of brachial plexus palsies.** *Plast Reconstr Surg* 2006, **117**:915-919.
13. Venkatramani H, Bhardwaj P, Faruquee SR, Sabapathy SR: **Functional outcome of nerve transfer for restoration of shoulder and elbow function in upper brachial plexus injury.** *J Brachial Plex Peripher Nerve Inj* 2008, **3**:15.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp