concern regarding the possibility of nosocomial infection that could arise using this technique.

We would like to bring to your notice that we use disposable electrodes and we are cutting the centre of the electrode with metal piece and the outer sticking part is used. The gel used with Doppler is wiped off after every use.

We consider that the chances of infection is highly theoretical because one of the conventional techniques of monitoring free flaps worldwide is scratching the flap which itself is not known to cause infection. We consider that the risk of infection in this technique is extremely less. While no technique is fool proof and without disadvantage, our experience of using this on 120 cases over a 3-year period has given us courage to continue its practice. We are using it only on intact normal skin. If at any time due to humidity or moisture the lead gets wet or contaminated, you can always change it on daily basis, till the monitoring goes on.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

**Hari Venkatramani, Gururaj Hosahalli,
S. Raja Sabapathy**

Department of Plastic Surgery, Hand Surgery,
Reconstructive Microsurgery and Burns, Ganga Hospital,
Coimbatore, Tamil Nadu, India

Address for correspondence:

Dr. S. Raja Sabapathy, Division of Plastic Surgery, Hand Surgery,
Reconstructive Microsurgery and Burns, Ganga Hospital, 313,
Mettupalayam Road, Coimbatore - 641 043, Tamil Nadu, India.
E-mail: rajahand@gmail.com

Author's reply to comments on 'A simple technique of marking the perforator of a free flap for post-operative monitoring'

Sir,

We read with interest the comments on our article (A simple technique of marking the perforator of a free flap for post-operative monitoring). We appreciate the

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:

Website:

www.ijps.org

DOI:

10.4103/0970-0358.182252

How to cite this article: Venkatramani H, Hosahalli G, Sabapathy SR. Author's reply to comments on 'A simple technique of marking the perforator of a free flap for post-operative monitoring'. Indian J Plast Surg 2016;49:131.