Commentary

The authors presented two rare cases of scalp malignant nodular hidradenoma (MNH).^[1] Both tumors recurred in spite of total excision and radiotherapy. The preferred treatment is an early wide surgical excision of the tumor. Recently, innovation of radiation delivery technique enabled IMRT (intensity modulated radiation therapy) for complex-shaped targets.^[2] VMAT (volumetric modulated arc therapy) or Rapid Arc^[3] therapy and helical tomotherapy^[4] give a high dose to widely spread scalp and skull tumors, sparing the underlying brain parenchyma as much as possible. IMRT as well as chemotherapy including the molecular targeting therapy is to be evaluated for adjuvant therapy after the surgical resection or for salvage therapy in case of recurrence.

Yoshimasa Mori

Department of Radiology and Radiation Oncology, Aichi Medical University, Nagakute, Aichi, Japan

Address for correspondence:

Prof. Yoshimasa Mori,
Department of Radiology and Radiation Oncology,

Aichi Medical University, Nagakute, Aichi, Japan.
E-mail: yoshim@aichi-med-u.ac.jp

References

1. Maiti T, Somanna S, Devi BI, Unchag A, Shukla D. Malignant nodular hidradenoma of scalp. *J Neurosci Rural Pract* 2014;4:423-25.
2. Samant RS, Fox GW, Gerig LH, Montgomery LA, Allan DS. Total scalp radiation using image-guided IMRT for progressive cutaneous T cell lymphoma. *Br J Radiol* 2009;82:e122-5.
3. Kelly PJ, Mannarino E, Lewis JH, Baldini EH, Hacker FL. Total dural irradiation: RapidArc versus static-field IMRT: A case study. *Med Dosim* 2012;37:175-81.
4. Gupta T, Mallik S, Master Z, Phurailatpam R, Laskar S. Brain-sparing holo-cranial radiotherapy: A unique application of helical tomotherapy. *Clin Oncol (R Coll Radiol)* 2011;23:86-94.

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com