Commentary

Giant cell arteritis (GCA) has a predilection for the extracranial branches of the carotid artery. Classic manifestation of the disease is scalp tenderness and jaw claudication and the temporal artery may appear swollen and pulseless. The disease may affect all organs but the most feared complication is sudden, irreversible blindness. Furthermore, polymyalgia rheumatica is often associated with GCA. The aetiology of both disorders is currently unknown.^[1,2]

GCA is reported to have an uneven geographical distribution.^[3] From a Danish perspective it is a relatively common disorder with more than 300 annual inpatient contacts from a background population of five million

people [Figure 1].^[4,5] However, GCA is thought to be rare in the developing countries and in patients of Asian background.^[3]

GCA is easy to diagnose and must be suspected in the patients aged over 50 years, with an increased erythrocyte sedimentation rate, and general symptoms of myalgia or headache. The gold-standard for GCA diagnosis is still the temporal artery biopsy, but clinicians need to be aware that false negative results may occur. It is of importance to diagnose GCA since the treatment with glucocorticoids often lead to rapid improvement. The standard treatment regimen is a high-dose glucocorticoid therapy (60 mg/day) which is gradually reduced over several months.^[2]

Figure 1: The age-specific rate of Inpatient contacts due to GCA in Denmark in the period 2009-2012. Source

In conclusion, the diagnosis and the treatment of GCA should be common knowledge to all neurologists, and symptoms of this disease must not be misinterpreted as tension type headache or opticneuritis.^[6,7]

References

- Weyand CM, Goronzy JJ. Medium and large-vessel vasculitis. N Engl J Med 2003;349:160-9.
- Mackie SL, Pease CT. Diagnosis and management of giant cell arteritis and polymyalgia rheumatica: Challenges, controversies and practical tips. Postgrad Med J 2013;89:284-92.
- Lee JL, Naguwa SM, Cheema GS, Gershwin ME. The geo-epidemiology of temporal (giant cell) arteritis. Clin Rev Allergy Immunol 2008;35:88-95.
- The National Registry of Patients. 2012. Ref Type: Online Source.
- Statistics Denmark. 2012. Ref Type: Online Source.
- Horwitz H, Friis T, Modvig S, Roed H, Tsakiri A, Laursen B, et al. Differential diagnoses to MS: Experiences from an optic neuritis clinic. J Neurol 2014;261:98-105.
- Rana AQ, Saeed U, Khan O, Qureshi AM, Paul D. Giant cell arteritis or tension-type headache? A differential diagnostic dilemma. J Neurosci Rural Pract 2014;4:409-411.

Henrik Horwitz

Department of Clinical Pharmacology, Bispebjerg Hospital, Copenhagen, Denmark

Address for correspondence:

Dr. Henrik Horwitz,
Department of Clinical Pharmacology, Bispebjerg Hospital,
Copenhagen, Denmark.
E-mail: henrik_horwitz@hotmail.com

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com