initially include catheter fixation and anchoring to the organ serosa from a local inflammatory reaction. Subsequently, bowel peristaltic movements, increased intraabdominal pressure, and CSF pulsations are believed to provide a continuous source of mechanical pressure that ultimately lead to necrosis at the anchoring site, perforation, and finally extrusion.^[9]

The presence of diphtheroids in the culture specimen from the ventricular catheter tip, which was kept separate from the abdominal-pelvic portion of the shunt tubing, may have contributed to the vaginal extrusion as well. Infections and hardware contamination are documented to contribute to wound erosion in multiple compartments of the body. However, the CSF analysis showed no evidence of inflammation, so it is unclear whether this germ was contributing to the vaginal erosion, or if it was perhaps a contaminant in this case.

The Raimondi catheters used in past VPS had a firm tip to avoid kinking. There was concern that this characteristic could contribute to organ perforation. However, in our case, the distal end of the Strata catheter (Medtronic, Inc. Minneapolis, MN) was soft and not reinforced. Thus, it appears that even with the modern silastic VPS catheters, vaginal extrusion can occur in an adult.

Managing such a complication requires a multi-disciplinary team, including neurosurgeons and gynecologists. Most importantly for repair of the protrusion, a cystoscopy must be performed to prove the bladder is not damaged. Finally, before pulling the catheter out, a stitch must be placed to keep the defect in site and then tied down once the catheter has been retrieved.

Behcet's syndrome is an immune-mediated small-vessel vasculitis that disrupts mucosal membranes as well as ocular, gastrointestinal, pulmonary, and genitourinary dysfunction.^[10] Wound healing is known to be generally impaired in this patient population. While it is unknown what caused the vaginal perforation in this particular case, a history of prior hysterectomy and

chronic mucosal dysfunction may have put this patient at higher risk.

Although very rare, vaginal extrusion can occur in patients other than children. To this end, consideration of alternative placement of the distal shunt catheter outside of the abdomen may be given in patients with a history of prior hysterectomy, gynecologic surgery, or chronic mucosal immunosuppression. Finally, if the peritoneum is ultimately used as the location for placement of the distal catheter, utilizing a shorter catheter length may be beneficial.

References

1. Agha FP, Amendola MA, Shirazi KK, Amendola BE, Chandler WF. Abdominal complications of ventriculoperitoneal shunts with emphasis on the role of imaging methods. *Surg Gynecol Obstet* 1983;156:473-8.
2. Mozingo JR, Cauthen JC. Vaginal perforation by a raimondi peritoneal catheter in an adult. *Surg Neurol* 1974;2:195-6.
3. Akcora B, Serarslan Y, Sangun O. Bowel perforation and transanal protrusion of a ventriculoperitoneal shunt catheter. *Pediatr Neurosurg* 2006;42:129-31.
4. Eichel L, Allende R, Mevorach RA, Hulbert WC, Rabinowitz R. Bladder calculus formation and urinary retention secondary to perforation of a normal bladder by a ventriculoperitoneal shunt. *Urology* 2002;60:344.
5. Fewel ME, Garton HJ. Migration of distal ventriculoperitoneal shunt catheter into the heart. Case report and review of the literature. *J Neurosurg* 2004;100 (Suppl Pediatrics 2):206-11.
6. Wilson CB, Bertan V. Perforation of the bowel complicating peritoneal shunt for hydrocephalus. Report of two cases. *Am Surg* 1966;32:601-3.
7. Celik A, Ergün O, Arda MS, Yurtseven T, Erşahin Y, Balık E. The incidence of inguinal complications after ventriculoperitoneal shunt for hydrocephalus. *Childs Nerv Syst* 2005;21:44-7.
8. Adeloje A. Spontaneous extrusion of the abdominal tube through the umbilicus complicating peritoneal shunt for hydrocephalus. Case report. *J Neurosurg* 1973;38:758-60.
9. de Aguiar GB, Mizrahi C, Aquino JH, Tavares CM, Telles C, Nigri F, *et al.* Urethral extrusion of a peritoneal catheter in a patient with a neobladder: A rare complication of shunt insertion. *Neuropediatrics* 2011;42:124-7.
10. Löhrer R, Eming R, Wolfrum N, Krieg T, Eming SA. Autoinflammatory diseases as cause of wound healing defects. *Hautarzt* 2011;62:524-33.

How to cite this article: Bonfield CM, Weiner GM, Bradley MS, Engh JA. Vaginal extrusion of a ventriculo-peritoneal shunt catheter in an adult. *J Neurosci Rural Pract* 2015;6:97-9.

Source of Support: Nil. **Conflict of Interest:** None declared.

Commentary

Hippocrates, 5th Century B.C. (460-377 BC), Greek physician and surgeon, the father of medicine, is thought to be the first physician to attempt and document the

treatment of hydrocephalus.^[1] Vesalius (1514-1564) clarified many of the anatomical and pathological characteristics of hydrocephalus.^[1] Hydrocephalus

is a common pediatric disorder in which there is an increase in cerebrospinal fluid (CSF) volume, which in turn causes enlargement of the ventricles, thinning of the cortical mantle, and elevation of the intracranial pressure.^[2] The use of peritoneal cavity for CSF absorption in ventriculoperitoneal (VP) shunting was introduced in 1908 by Kausch; since then, VP shunt are amongst the most frequently performed operations for the management of hydrocephalus; not only in children, but also in adults.^[1-4] Other shunting techniques are ventriculoatrial shunts, endoscopic third ventriculostomy, and lumboperitoneal shunts.^[5-7] Endoscopic third ventriculostomy is more physiological and anatomical diversion pathway for CSF and at the same time, it obviates the needs to place a foreign body such as ventricular shunt, thus avoiding shunt-related complications such as malfunction, infection, and over-drainage.^[8]

Although the risks in performing a shunt operation are low, the complications related to shunts are many and reported to occur in 24% to 47% of the cases, of which abdominal complications are reported in 25% of VP shunt operations. Many of the shunt complications need shunt revision and many of them require multiple revisions.^[3,4,9]

Extrusion/protrusion of the peritoneal catheter are an unusual but serious complication following VP shunting.^[3,9] The bowel is the most commonly and frequently involved site for perforation and extrusion by peritoneal catheter and is reported to occur in up to 2.5% of cases.^[3,9] Non-enteric viscous perforation has also been sporadically reported in literature and includes urinary bladder, vagina, uterus, urethra, etc., VP shunt catheter extrusion/protrusion has also been reported through umbilicus, gastrostomy wound, healed abdominal scar, scrotal skin, mouth, etc.^[5,9,10] In cases of bowel perforation and extrusion of peritoneal catheter through anus, the interval between shunt insertions to extrusion may range from 2 to 20 months, and most of them present within 6 months following shunt insertion.^[3,9] Many of the cases with extrusion of peritoneal catheter through non - enteric viscous may also present within 6 months following shunt insertion.^[9,10] The occurrence of a viscus perforation by the abdominal portion of a VP shunts usually does not lead to acute peritonitis.^[3,9,10] Acute viscus perforation may also occur due to peritoneal catheter following VP shunting; this is the exception rather than the rule.

The exact mechanism for the extrusion of the peritoneal catheter is not known. Multiple factors may play an important role for the occurrence of the perforation of the viscus and ultimately extrusion of the catheter and includes infection, meningitis, long length of peritoneal

catheter, foreign body reaction for shunt, pressure necrosis, age of patient, peristaltic motility of bowel, poor surgical technique, poor host immunity etc.^[3,9,10]

Management of the extruded shunt must be individualized and depends on the involved viscera, presence or absence of infection (shunt/shunt tract infection, meningitis, and peritonitis), functional status of the shunt system etc., Many options are available for the management of patients who presented with extrusion of peritoneal catheter and are (1) Mini laparotomy and revision of peritoneal catheter of the VP shunt. Most of the cases who presented with extruded catheter; they present without peritonitis and, therefore, formal abdominal exploration for localization of perforation and repair of the same are not required, (2) formal exploratory laparotomy for repair of perforation and shunt revision in selected cases having peritonitis, (3) shunt removal, external ventricular drainage, followed by delayed VP shunt/VA shunt or contra-lateral shunt, (4) removal of entire shunt system and contra lateral shunting or delayed shunting.^[3,9,10] Shunt extrusions can also be managed using laparoscope, cystoscope, sigmoidoscope etc., depending up on the site of peritoneal catheter extrusion. During shunt revisions, if there is evidence of infection (shunt, shunt tract infection, meningitis, peritonitis), then shunt removal and external ventricular drainage would be a preferred approach.

VP shunt operations done for the management of hydrocephalus are not only prone for complications, but also require shunt revisions in the follow-up period. Visceral perforation and extrusion/protrusion by peritoneal catheter are also well-known complication and may occur following VP shunt operations. A regular follow-up is a must during first 6 months after the VP shunt surgery as majority of complications are being reported within this period.

Rajendra K Ghritlaharey

Department of Pediatric Surgery, Gandhi Medical College and Associated Kamla Nehru and Hamidia Hospitals, Bhopal, Madhya Pradesh India

Address for correspondence:

Dr. Rajendra K Ghritlaharey,
Department of Pediatric Surgery,
Gandhi Medical College and Associated Kamla Nehru and
Hamidia Hospitals, Bhopal,
Madhya Pradesh - 462 001, India.
E-mail: drrajendrak1@rediffmail.com

References

1. Lifshutz JJ, Johnson WD. History of hydrocephalus and its treatments. *Neurosurg Focus* 2001;11:E1.

2. Smith JL. Management of neural tube defects, hydrocephalus, refractory epilepsy, and central nervous system infections. In: Grosfeld JL, O'Neill JA Jr, Fonkalsrud EW, Coran AG, Caldamone AA, editors. *Pediatric Surgery*. 6th ed. Vol. 2.: Mosby Yearbook Inc, USA; 2006. p. 1987-2017.
3. Ghritlaharey RK, Budhwani KS, Shrivastava DK, Gupta G, Kushwaha AS, Chanchlani R, *et al.* Trans-anal protrusion of ventriculo-peritoneal shunt catheter with silent bowel perforation: Report of ten cases in children. *Pediatr Surg Int* 2007;23:575-80.
4. Reddy GK, Bollam P, Shi R, Guthikonda B, Nanda A. Management of adult hydrocephalus with ventriculoperitoneal shunts: Long-term single-institution experience. *Neurosurgery* 2011;69:774-81.
5. Surchev J, Georgiev K, Enchev Y, Avramov R. Extremely rare complications in cerebrospinal fluid shunt operations. *J Neurosurg Sci* 2002;46:100-3.
6. Kulkarni AV, Drake JM, Mallucci CL, Sgouros S, Roth J, Constantini S, Canadian Pediatric Neurosurgery Study Group. Endoscopic third ventriculostomy in the treatment of childhood hydrocephalus. *J Pediatr* 2009;155:254-9.e1.
7. Yadav YR, Pande S, Raina VK, Singh M. Lumboperitoneal shunts: Review of 409 cases. *Neurol India* 2004;52:188-90.
8. Ray P, Jallo GI, Kim RY, Kim BS, Wilson S, Kothbauer K, *et al.* Endoscopic third ventriculostomy for tumor-related hydrocephalus in a pediatric population. *Neurosurg Focus* 2005;19:E8.
9. Ghritlaharey RK, Budhwani KS, Shrivastava DK, Shrivastava J. Ventriculoperitoneal shunt complications needing shunt revision in children: A review of 5 years of experience with 48 revisions. *Afr J Paediatr Surg* 2012;9:32-9.
10. Kumar B, Sharma SB, Singh DK. Extrusion of ventriculo-peritoneal shunt catheter. *Indian J Pediatr* 2010;77:336.
11. Bonfield CM, Weiner GM, Bradley MS, Donnellan NM, Engh JA. Vaginal extrusion of a ventriculo-peritoneal shunt catheter in an adult. *J Neurosc Rural Pract* 2015;6:10-6.

Access this article online	
Quick Response Code:	Website: www.ruralneuropractice.com