Tubercular meningitis in children: Clinical, pathological, and radiological profile and factors associated with mortality

Anil V. Israni, Divya A. Dave¹, Anirban Mandal, Amitabh Singh², Puneet K. Sahi³, Rashmi Ranjan Das⁴, Arpita Shah⁵

Department of Pediatrics, All India Institute of Medical Sciences, ²Department of Pediatrics, Chacha Nehru Bal Chikitsalaya, ³Department of Pediatrics, Kalawati Saran Children Hospital, New Delhi, ¹Department of Pediatrics, Medical College Baroda and SSG Hospital, ⁵Department of Pathology, Toprani Advanced Lab Systems, Vadodra, Gujarat, ⁴Department of Pediatrics, All India Institute of Medical Sciences, Bhubaneswar, Odisha, India

ABSTRACT

Context: Childhood tuberculosis is a major public health problem in developing countries with tubercular meningitis being a serious complication with high mortality and morbidity. **Aim:** To study the clinicopathological as well as radiological profile of childhood tuberculous meningitis (TBM) cases. **Settings and Design:** Prospective, observational study including children <14 years of age with TBM admitted in a tertiary care hospital from Western India. **Subjects and Methods:** TBM was diagnosed based on predefined criteria. Glasgow coma scale (GCS) and intracranial pressure (ICP) was recorded. Staging was done as per British Medical Council Staging System. Mantoux test, chest X-ray, cerebrospinal fluid (CSF) examination, neuroimaging, and other investigations were done to confirm TB. **Statistical Analysis Used:** STATA software (version 9.0) was used for data analysis. Various risk factors were determined using Chi-square tests, and a $P < 0.05$ was considered significant. **Results:** Forty-seven children were included, of which 11 (24.3%) died. Fever was the most common presenting symptom, and meningismus was the most common sign. Twenty-nine (62%) children presented with Stage III disease. Stage III disease, low GCS, and raised ICP were predictors of mortality. Findings on neuroimaging or CSF examination did not predict mortality. **Conclusions:** Childhood TBM presents with nonspecific clinical features. Stage III disease, low GCS, lack of Bacillus Calmette–Guérin vaccination at birth and raised ICP seem to be the most important adverse prognostic factors.

Key words: Children, developing country, meningitis, mortality, tuberculosis

Introduction

India has one of the highest burdens of tuberculosis (TB) globally, accounting for around 20% of all new TB cases annually.^[1] It is estimated that childhood TB constitutes 10–20% of all TB cases in high burden countries,^[2] accounting for 8–20% of TB-related deaths.^[3] Approximately, 25% of pediatric TB cases are extrapulmonary, with tuberculous meningitis (TBM) being the most severe form. Worldwide, TBM accounts for majority of the

deaths due to TB.^[1] A study from South Africa (a high burden country) reported TB to be the commonest cause of childhood meningitis.^[4] TBM continues to be an important cause of morbidity (especially neurologic handicap) in children from resource-poor countries. Due to suboptimal performance of the diagnostic tests, TBM diagnosis requires vital clinical information coupled with supportive investigations (biochemical, immunological, and radiological).^[5] There have been few studies looking at the clinico-patho-radiological spectrum of this disease in children from India,^[6] with fewer addressing the outcome predictors.^[7,8] This prospective, observational study was designed to address this issue.

Address for correspondence:

Dr. Rashmi Ranjan Das, Department of Pediatrics, All India Institute of Medical Sciences, Bhubaneswar, Odisha, India.
E-mail: rrdas05@gmail.com

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/0976-3147.181475

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Israni AV, Dave DA, Mandal A, Singh A, Sahi PK, Das RR, *et al.* Tubercular meningitis in children: Clinical, pathological, and radiological profile and factors associated with mortality. J Neurosci Rural Pract 2016;7:400-4.

Subjects and Methods

Children of 3 months to 14 years of age with suspected TBM admitted to a tertiary care hospital from Western India were included in the study. TBM was suspected based on following features: Fever and/or a cough for ≥ 2 weeks, neurological symptoms (irritability, refusal to feed, headache, vomiting, altered sensorium, abnormal movements or seizure) with weight loss or poor weight gain. The diagnosis of TBM was based on the clinical case definition by Doerr *et al.* [Text box 1].^[9] Ethical clearance was obtained from the Institute Ethics Committee and a written informed consent was taken from either of the parents/legal guardian before enrollment. TBM cases on treatment, those with underlying chronic illness or malignancy or HIV, and those on immunosuppressive therapy were excluded. Demographic data, immunization, contact with TB, and socioeconomic status were recorded.^[10] Detailed clinical examination, anthropometry, and the presence of Bacillus Calmette–Guérin (BCG) scar were recorded.^[11] The level of consciousness (modified Glassgow coma scale (GCS)), signs of raised intracranial pressure (ICP) and meningeal irritation was noted. Staging of tubercular meningitis was done as per British Medical Council Staging System [Text box 2].^[12]

All the recruited children underwent chest X-ray, Mantoux test, and complete hemogram. Additional testing included gastric aspirate for acid-fast *Bacilli* (AFB), ultrasonography of abdomen and fine needle aspiration cytology from enlarged lymph nodes. A standard

Text box 1: Clinical case definition of tuberculous meningitis devised by Doerr *et al.*

Abnormal neurological signs and/or symptoms, and two or more of the following

Discovery of adult source patient with contagious TB who had significant contact with child

Presence of Mantoux (5 tuberculin units) skin test reaction

Cerebrospinal fluid abnormalities without evidence of other infectious cause

Abnormalities on cranial computed tomography consistent with central nervous system TB

TB: Tuberculosis

Text box 2: British Medical Council Staging System for tubercular meningitis

Stage I: No definite neurological symptoms on admission or in the history before admission, with or without meningismus

Stage II: Signs of meningeal irritations with or without slight clouding of consciousness with focal neurological signs such as cranial nerve palsies or hemiparesis

Stage III: Severe clouding of consciousness or delirium, convulsions and serious neurological signs such as hemiplegia, paraplegia, involuntary movements

tuberculin skin test (5 TU) was used and results read after 72 h (positive: Induration of ≥ 10 mm; HIV and severe malnutrition: ≥ 5 mm was considered positive). Cerebrospinal fluid (CSF) analysis was done including the presence of AFB. All the children underwent neuroimaging (plain and contrast enhanced computed tomography or magnetic resonance imaging [MRI]). HIV testing was done where appropriate.

The data were entered into Microsoft excel sheet and analyzed using Stata 9.0 software (STATA Corp, College Station, TX, USA). Descriptive statistics was used along with analysis of risk factors by using Chi-square test or Fisher's test. A $P < 0.05$ was considered statistically significant.

Results

Of 50 TBM cases, 47 met the eligibility criteria and were included. The baseline characteristics and clinical features are presented in Table 1. Majority of children were from lower socioeconomic status (85%) and were malnourished (76.6%). Fever, altered sensorium, and seizure were the most common symptoms. Contact history of TB was positive in 49% cases, 55% of children were BCG vaccinated, Mantoux test was positive in 25% [Table 1]. CSF examination could be done in 45 children (10% had opalescent appearance, no xanthochromia or hemorrhage or cobweb formation was noted). Chest X-ray abnormality was found in 38% [Table 2]. Common abnormalities noted on neuroimaging were: Communicating hydrocephalus (70%), meningeal enhancement (64%), and infarction (45%) [Table 2]. Meningeal enhancement was most commonly observed over the basal cisterns (47%) and cerebral cortex (43%). Infarction was most common in basal ganglion (67%) followed by cortex (41%).

The mortality rate was 23.4% (11 of 47 died). Various risk factors associated with mortality or poor outcome were analyzed. Stage III disease, low GCS, and raised ICP were associated with increased mortality [Table 3], whereas biochemical and cytological parameters did not affect the outcome (data not shown). A significantly increased risk of mortality was found in children who were not vaccinated with BCG at birth.

Discussion

We summarized the clinical, pathological, and radiological features of TBM and investigated characteristics associated with mortality. Children with TBM had nonspecific clinical pictures, boys were commonly affected, Mantoux positivity was in 25%, chest X-ray

Table 1: Baseline and clinical characteristics of children with tubercular meningitis

Characteristics	n (%)
Male	28 (59.6)
Age distribution (years)	
0-3	28 (59.6)
3-5	8 (17)
>5	11 (23.4)
Socioeconomic status	
Upper-middle	1 (2.1)
Lower-middle	6 (12.8)
Lower	40 (85.1)
Positive history of contact with TB	23 (49)
BCG vaccination	26 (55.3)
Nutritional status	
Normal	11 (23.4)
Malnutrition (mild to moderate)	31 (66)
Malnutrition (severe)	5 (10.6)
Symptoms	
Fever	34 (72.3)
Altered sensorium	27 (57.4)
Seizure/abnormal movements	23 (48.9)
Abnormal posturing	13 (27.7)
Headache	12 (25.5)
Irritability	9 (19.2)
Vomiting	8 (17)
Refusal of feeds	4 (8.5)
Limping	2 (4.2)
Cough	1 (2.1)
Paucity of movement	1 (2.1)
Increasing head size	1 (2.1)
Signs	
Meningeal irritation	31 (66)
Hemiparesis	7 (14.9)
Cranial nerve palsy	6 (12.8)
Quadriparesis	4 (8.5)
Abnormal movements	3 (6.4)
Level of consciousness (GCS) (n=41)	
<7	15 (36.6)
7-10	9 (22)
>10	17 (41.4)
Raised ICP	31 (66)
Stage of TBM	
I	7 (14.9)
II	11 (23.4)
III	29 (61.7)

ICP: Intracranial pressure, TB: Tuberculosis, BCG: Bacillus Calmette-Guérin, TBM: Tuberculous meningitis, GCS: Glasgow coma scale

compatible with TB in 38% cases, and communicating hydrocephalus was the most common neuroimaging finding. Stage III disease, low GCS, and raised ICP were associated with increased mortality.

The male preponderance (1.5:1) observed in the present study was consistent with earlier pediatric studies from India.^[6,7] Most common age group affected

was <3 years (60%) followed by 3–6 years (16%). Globally, children <5 years of age have been found to be most vulnerable.^[13-15] Contact with TB could be elicited in about 50% of children in present study, this figure has been reported between 33% and 69% in various studies.^[6,14,15] This emphasizes the importance of eliciting contact history and family screening for TB in suspected cases of TBM. Despite being almost universally recommended and potentially highly effective, screening to provide preventive therapy for eligible contacts is still rarely implemented in TB endemic communities.^[16]

Though BCG vaccination is included under the universal immunization coverage, only 55% of children were vaccinated in the present study. As BCG vaccination protects against severe forms of TB (e.g., TBM, and disseminated or millitary TB), a lower vaccination coverage could explain a higher prevalence of severe form of TB (e.g., TBM) in the present study.^[17] Studies from India and Western countries have clearly documented the effect of BCG vaccine in reducing mortality from TBM.^[17-19] In a systematic review and metanalysis for treatment outcome of TBM in children, the authors emphasized on the role of BCG vaccination as an invaluable preventive measure for TBM.^[19] Our study demonstrated mantoux positivity only at mere 25% in children with TBM. TBM being an extra-pulmonary disease, a low rate of positive mantoux reaction in children with TBM is expected, though the rates vary widely.^[6-8,14]

Most of the children (62%) in this study were detected at an advanced stage (Stage III). Advanced stage has been found to be the single most important factor associated with poor outcome,^[14] and the same has been echoed in our study. Mortality was nil in Stage I or Stage II disease but was 38% in Stage III disease. Cranial nerve palsy has been recognized as an important clinical indicator in differentiating TBM from pyogenic meningitis.^[20] In this study, 13% children present with cranial nerve palsy. Around 54% of children having GCS <7 died, and GCS was found to be one of the predictors of mortality in the present study. This has been reported in previous Indian studies.^[8] Around 39% of children with raised ICP died, and this was found to be another predictors of mortality. Previous studies also report similar observation.^[8]

Abnormal chest X-ray was observed in 38% of patients in this study. A previous study found this figure as 44%.^[21] None of the radiological findings (chest X-ray, computed tomography scan, or MRI brain) predicted the mortality. While looking at the CSF findings, patients dying of TBM had higher mean CSF protein values and also CSF lymphocytic pleocytosis compared to those who survived.

The overall mortality in the present study was 23.4%. A previous systematic review reported this figure as 19.3%.^[22] However, in 10 years comparative retrospective analysis, one study reported this figure as 8% in children with TBM.^[23]

The strengths of the present study are prospective enrollment of children and careful documentation using

Table 2: Laboratory and neuroimaging findings in children with tuberculous meningitis

Investigations	n (%)
Mantoux positive	12 (25)
Abnormal chest X-ray	18 (38)
HIV positive	2 (4.3)
CSF protein (n=45) (mg/dl)	
<40	4 (8.9)
40-400	40 (88.9)
>400	1 (2.2)
CSF glucose (n=45) (mg/dl)	
<20	12 (26.7)
20-60	25 (55.6)
>60	8 (17.7)
CSF total cells (n=45)/mm ³	
<10	12 (26.7)
10-100	10 (22.2)
101-400	18 (40)
>400	5 (11.1)
CSF lymphocyte percentage (n=45)	
<20	3 (6.6)
20-80	21 (46.7)
>80	21 (46.7)
Abnormality on neuroimaging	
Meningeal enhancement	30 (63.8)
Hydrocephalus	33 (70.2)
Periventricular edema	21 (44.7)
Infarcts	27 (57.4)
Tuberculoma	5 (10.6)

CSF: Cerebrospinal fluid

a standardized predesigned data collection form. The case definition of TBM was also predefined, and the same was also used in a previous study from India, thus making it easier to compare the results. The limitations are: Being a single center study conducted at a tertiary care referral hospital, referral bias cannot be ruled out; TB culture and nucleic acid amplification tests (Genexpert) were not conducted on the CSF specimen; treatment outcome and follow-up of the patients to look for neurological sequelae could not be done.

The incidence of TBM indicates the annual risk of infection, and a robust surveillance system for documenting the occurrence of TBM in young children can improve estimates and monitoring of the TB burden and TB-related deaths in children.^[24] Newer and easier scoring systems need to be devised in the lights of ever increasing knowledge for better prediction of outcome of this dreaded disease.^[25] Furthermore, the survivors of tubercular meningitis are at a risk of long-term neurological sequelae ranging from residual paresis to subtle neurological deficit in the form of poor scholastic performance or even behavioral abnormalities. This mandates a close and regular follow-up of these children for early intervention.

Conclusion

Tubercular meningitis continues to be associated with high rate mortality in children of countries with high burden of TB. Even with the advancement of scientific knowledge and technologies, including new techniques of laboratory and imaging diagnostic aid, very little prediction can be made regarding the prognosis of children with this dreaded disease. Our study highlighted few important predictors associated with mortality in

Table 3: Factors associated with mortality from tubercular meningitis in children

Risk factors	Categories	Outcome		OR (95% CI)	P
		Survived	Expired		
Age	≥ 5 years	8	3	1.31 (0.28-6.14)	0.73
Gender	Male	21	7	1.25 (0.31-5.05)	0.75
History of contact with TB	Present	17	6	3.88 (0.69-21.7)	0.1
Socioeconomic class	Lower class	29	11	-	0.11
Nutritional status	Severe to very severe PEM	6	4	1.5 (0.33-6.82)	0.59
GCS	<7	7	8	28.6 (3.04-268.77)	<0.001
BCG vaccination	Unvaccinated	13	8	4.72 (1.06-20.96)	0.03
Mantoux test	Negative	29	6	1.03 (0.18-5.98)	0.97
Stage of TBM	III	18	11	-	0.002
Raised ICP	Present	17	11	-	0.001
Abnormality in chest X-ray	Present	13	5	3.33 (0.69-16.16)	0.12
Hydrocephalus	Present	22	9	2.86 (0.54-15.25)	0.21
Brain infarction	Present	20	7	1.58 (0.39-6.28)	0.73

OR: Odds ratio, CI: Confidence interval, ICP: Intracranial pressure, TBM: Tuberculous meningitis, GCS: Glasgow coma scale, TB: Tuberculosis, PEM: Protein-energy malnutrition

this age group coupled with a reemphasis on need for early diagnosis, though the challenge remains to devise composite index systems in predicting outcome in TBM in future years.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- World Health Organization (WHO). Global Tuberculosis Report 2013. Geneva: WHO; 2013. Available from: http://www.apps.who.int/iris/bitstream/10665/91355/1/9789241564656_eng.pdf. [Last accessed on 2015 Dec 15].
- Marais BJ, Hesselning AC, Gie RP, Schaaf HS, Beyers N. The burden of childhood tuberculosis and the accuracy of community-based surveillance data. *Int J Tuberc Lung Dis* 2006;10:259-63.
- Jain SK, Ordonez A, Kinikar A, Gupte N, Thakar M, Mave V, et al. Pediatric tuberculosis in young children in India: A prospective study. *Biomed Res Int* 2013;2013:783698.
- Wolzack NK, Cooke ML, Orth H, van Toorn R. The changing profile of pediatric meningitis at a referral centre in Cape Town, South Africa. *J Trop Pediatr* 2012;58:491-5.
- van Toorn R, Solomons R. Update on the diagnosis and management of tuberculous meningitis in children. *Semin Pediatr Neurol* 2014;21:12-8.
- Thilothammal N, Krishnamurthy PV, Banu K, Ratnam SR. Tuberculous meningitis in children – Clinical profile, mortality and morbidity of bacteriologically confirmed cases. *Indian Pediatr* 1995;32:641-7.
- Ramzan A, Nayil K, Asimi R, Wani A, Makhdoomi R, Jain A. Childhood tubercular meningitis: An institutional experience and analysis of predictors of outcome. *Pediatr Neurol* 2013;48:30-5.
- Karande S, Gupta V, Kulkarni M, Joshi A. Prognostic clinical variables in childhood tuberculous meningitis: An experience from Mumbai, India. *Neurol India* 2005;53:191-5.
- Doerr CA, Starke JR, Ong LT. Clinical and public health aspects of tuberculous meningitis in children. *J Pediatr* 1995;127:27-33.
- Kumar N, Shekhar C, Kumar P, Kundu AS. Kuppaswamy's socioeconomic status scale-updating for 2007. *Indian J Pediatr* 2007;74:1131-2.
- Khadilkar VV, Khadilkar AV, Choudhury P, Agarwal KN, Ugra D, Shah NK. IAP growth monitoring guidelines for children from birth to 18 years. *Indian Pediatr* 2007;44:187-97.
- British Medical Research Council. Streptomycin treatment of tuberculous meningitis. *Br Med J* 1948;1:582-97.
- Humphries MJ, Teoh R, Lau J, Gabriel M. Factors of prognostic significance in Chinese children with tuberculous meningitis. *Tubercle* 1990;71:161-8.
- van Well GT, Paes BF, Terwee CB, Springer P, Roord JJ, Donald PR, et al. Twenty years of pediatric tuberculous meningitis: A retrospective cohort study in the western cape of South Africa. *Pediatrics* 2009;123:e1-8.
- Lee LV. Neurotuberculosis among Filipino children: An 11 years experience at the Philippine Children's Medical Center. *Brain Dev* 2000;22:469-74.
- Hill PC, Rutherford ME, Audas R, van Crevel R, Graham SM. Closing the policy-practice gap in the management of child contacts of tuberculosis cases in developing countries. *PLoS Med* 2011;8:e1001105.
- Kelekçi S, Karabel M, Karabel D, Hamidi C, Hosoglu S, Gürkan MF, et al. *Bacillus Calmette-Guérin* is a preventive factor in mortality of childhood tuberculous meningitis. *Int J Infect Dis* 2014;21:1-4.
- Kumar R, Dwivedi A, Kumar P, Kohli N. Tuberculous meningitis in BCG vaccinated and unvaccinated children. *J Neurol Neurosurg Psychiatry* 2005;76:1550-4.
- Chiang SS, Khan FA, Milstein MB, Tolman AW, Benedetti A, Starke JR, et al. Treatment outcomes of childhood tuberculous meningitis: A systematic review and meta-analysis. *Lancet Infect Dis* 2014;14:947-57.
- Moghtaderi A, Alavi-Naini R, Rashki S. Cranial nerve palsy as a factor to differentiate tuberculous meningitis from acute bacterial meningitis. *Acta Med Iran* 2013;51:113-8.
- Solomons RS, Goussard P, Visser DH, Marais BJ, Gie RP, Schoeman JF, et al. Chest radiograph findings in children with tuberculous meningitis. *Int J Tuberc Lung Dis* 2015;19:200-4.
- Graham SM, Donald PR. Death and disability: The outcomes of tuberculous meningitis. *Lancet Infect Dis* 2014;14:902-4.
- Miftode EG, Dorneanu OS, Leca DA, Juganariu G, Teodor A, Hurmuzache M, et al. Tuberculous meningitis in children and adults: A 10-year retrospective comparative analysis. *PLoS One* 2015;10:e0133477.
- Styblo K, Sutherland I. The epidemiology of tuberculosis in children. *Bull Int Union Tuberc* 1982;57:133-9.
- Alarcón F, Moreira J, Rivera J, Salinas R, Dueñas G, Van den Ende J. Tuberculous meningitis: Do modern diagnostic tools offer better prognosis prediction? *Indian J Tuberc* 2013;60:5-14.