Importance of medial antebrachial cutaneous nerve anatomical variations in upper arm surgery

Stylianios *et al.* reported the unique case of atypical division of medial antebrachial cutaneous nerve (MACN) to four brachial cutaneous branches and presented a short literature review on MACN anatomical variations. Authors in detail described possible neurological consequences of the iatrogenic injury to the nerve and its branches.^[1] Their case report is of clinical importance in a multidisciplinary aspect. In trauma patients, an injury to minor nerves is not a critical issue. However, in patients undergoing elective surgery on the upper extremity, iatrogenic nerve injury with eventual, subsequent neurological deficits may be the source of considerable liability and litigation.

Vascular access surgeons during the formation of brachial-basilic and brachial-brachial arterio-venous fistula typically encounter the MACN, and its branches are passing over the basilic and brachial vein. Almost no reports are available in the literature how to technically deal with the MACN during upper arm vascular access surgery. Usually, the special care is advised not to injure the nerve and its branches.^[2] Specifically, the two-stage methods of brachial-basilic and brachial-brachial fistula creation require surgeon's attention. The second stage of surgery requires an exposure of the entire length of basilic or brachial vein on the arm. It is not an issue, if the surgeon decides to transect the matured vein, tunnel it on the anterior aspect of the arm, thus above the nerve and reanastomose the vein. In simplified transposition techniques, as superficialization and elevation, it is likely to encounter the anatomical conflict between the vein and the nerve. Recently, an elegant neurosurgical technique of axial splitting of the MACN nerve was described.^[3] This novel technique enables the elevation or superficialization of the vein between the separated nerve branches, without the need to transect and reanastomose the vein. The method significantly shortens the time of the second stage surgery and generally facilitates the procedure. Notably, the authors did not observe any significant neurological deficits during the follow-up.

Other vascular procedures on upper arm as thrombectomy, embolectomy of the brachial artery,

and arteriovenous fistula steal syndrome surgery (such as DRIL technique) may potentially lead to the iatrogenic injury of the MACN and its branches as well. Routine venipuncture and percutaneous endovascular procedures from brachial or axillary approach can also cause nerve injury.

Similarly to vascular surgery on the upper arm, plastic surgeons during traditional brachioplasty procedures perform a vast dissection of the upper arm. Longitudinal incision usually runs from the medial epicondyle to the axilla. As there is a possibility of iatrogenic nerve injury, an eventual anatomical variation of the upper extremity nervous system must not be neglected.^[4]

As the MACN is often used as a "cable graft" in neurosurgical reconstructions, patients undergoing surgery must be aware about possible neurological complications after harvesting.^[5]

Typical anatomy of MACN and its variations are widely described in the literature. Surgeons dealing with upper extremity surgery should be aware of this unique case of anatomical variation of MACN. Neurologists should as well consider this unusual variation during the postoperative neurological examination of patients who underwent an upper extremity surgery.

Stanisław Przywara

Department of Vascular Surgery and Angiology, Medical University of Lublin, Lublin, Poland

Address for correspondence:

Dr. Stanisław Przywara,
Department of Vascular Surgery and Angiology, Medical University of Lublin, Lublin, Poland.
E-mail: s.przywara@plusnet.pl

References

1. Stylianios K, Konstantinos G, Pavlos P, Aliki F. Brachial branches of the medial antebrachial cutaneous nerve: A case report with its clinical significance and a short review of the literature. *J Neurosci Rural Pract* 2016;7:443-6.
2. Dix FP, Khan Y, Al-Khaffaf H. The brachial artery-basilic vein arterio-venous fistula in vascular access for haemodialysis – A review paper. *Eur J Vasc Endovasc Surg* 2006;31:70-9.

3. Przywara S, Ilzecki M, Terlecki P, Zubilewicz T. Axial splitting of the medial antebrachial cutaneous nerve facilitates second-stage elevation of basilic or brachial vein in patients with arteriovenous fistula. *J Vasc Surg* 2015;62:1353-6.
4. Chowdhry S, Elston JB, Lefkowitz T, Wilhelmi BJ. Avoiding the medial brachial cutaneous nerve in brachioplasty: An anatomical study. *Eplasty* 2010;10:e16.
5. Nunley JA, Ugino MR, Goldner RD, Regan N, Urbaniak JR. Use of the anterior branch of the medial antebrachial cutaneous nerve as a graft for the repair of defects of the digital nerve. *J Bone Joint Surg Am* 1989;71:563-7.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.ruralneuropractice.com
DOI: 10.4103/0976-3147.181484

How to cite this article: Przywara S. Importance of medial antebrachial cutaneous nerve anatomical variations in upper arm surgery. *J Neurosci Rural Pract* 2016;7:337-8.

"Quick Response Code" link for full text articles

The journal issue has a unique new feature for reaching to the journal's website without typing a single letter. Each article on its first page has a "Quick Response Code". Using any mobile or other hand-held device with camera and GPRS/other internet source, one can reach to the full text of that particular article on the journal's website. Start a QR-code reading software (see list of free applications from <http://tinyurl.com/yzlh2tc>) and point the camera to the QR-code printed in the journal. It will automatically take you to the HTML full text of that article. One can also use a desktop or laptop with web camera for similar functionality. See <http://tinyurl.com/2bw7fn3> or <http://tinyurl.com/3ysr3me> for the free applications.