Cirroid aneurysm of scalp

Leo Francis Tauro, Suhith G, Prathvi Shetty, Diwakar Rao¹

Departments of Surgery and ¹Neurosurgery, Fr. Muller Medical College Hospital, Kankanady, Mangalore, India

A 40-year-old lady presented with swelling on the right side of the scalp associated with continuous headache for 1 month. Her general physical examination was normal. Local examination revealed a pulsatile swelling on the right temporal-occipital region of the scalp. Few dilated veins were noted in this region. On auscultation, a bruit was detected over the lesion. Other systemic examination was normal, including thyroid, breast, and musculoskeletal system.

Patient underwent computed tomography of the brain along with angiography. Aneurysmal dilated mass of vessels were detected in the right temporal-occipital region of scalp in the subcutaneous plane [Figure 1], which enhanced markedly. Feeding vessel was from the occipital artery posteriorly and the superficial temporal artery [Figure 2]. Large draining veins were noted running anteriorly and posterior to the occiput [Figure 3]. A diagnosis of cirroid aneurysm of scalp was made. Patient was advised to undergo excision of the aneurysm, but the patient refused surgery. She was discharged with an advice of regular follow-up.

Cirroid aneurysms of the scalp were first described in 1833 by Brecht.^[1] Cirroid aneurysms are rare arterio-venous fistulas of the scalp. They are usually congenital in etiology. However, traumatic fistulas have also been described.^[1] In 90% of the patients, the superficial temporal artery is the main supply to the fistula with only one dominant feeding artery in 71% of patients.^[2] In the remaining cases, there is usually involvement of both the superficial temporal and occipital arteries.^[1,2] Untreated patients can develop progressive scalp and

Figure 1: Unenhanced axial CT image shows a serpiginous subcutaneous mass in the right temporal-occipital region

Figure 2: Contrast-enhanced axial CT image shows marked contrast enhancement within the cirroid aneurysm

facial cosmetic deformity from the markedly tortuous subcutaneous vessels. However, this condition is not life-threatening.^[2] Surgical resection of the fistula is usually successful. Endovascular and percutaneous occlusion of the fistulas have been described. However, the results have been mixed.

Differential diagnosis of a pulsatile scalp swelling includes

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/0976-3147.91976

Address for correspondence:

Dr. Leo Francis Tauro, Department of General Surgery, Fr. Muller Medical College Hospital, Kankanady, Mangalore – 575 002, Karnataka, India. E-mail: dr1ftauro@rediffmail.com

Figure 3: CT angiogram showing the lesion

the following: superficial temporal artery aneurysm although rare should be considered when a temporal region mass is evaluated. This condition is almost always a result of blunt or penetrating head trauma. Clinical examination with imaging confirms the diagnosis. Simple elective ligation and excision of the aneurysm are curative.^[3]

A-V malformations may be congenital or traumatic in origin. Clinical manifestation may include a loud continuous bruit, hemorrhage and throbbing headache, and in severe cases scalp necrosis. Angiography is the investigation of choice. Primary lesions can be excised completely without significant blood loss and scalp necrosis.^[4] Metastatic deposits from follicular carcinoma of thyroid have to be ruled out prior to the diagnosis of cirroid aneurysm.

References

1. Muthukumar N, Rajagopal V, Manoharan A, Durairaj N. Surgical management of cirroid aneurysms. *Acta Neurochir (Wien)* 2002;144: 349-56.
2. Gurkanlar D, Gonul M, Solmaz I, Gonul E. Cirroid aneurysms of the scalp. *Neurosurg Rev* 2006;29:208-12.
3. Lee ES, Gower DJ, McWhorter TM. Aneurysm of the superficial temporal artery. *Neurosurgery* 1988;23:499-500.
4. Shenoy SN, Raja A. Scalp arteriovenous malformations. *Neurol India* 2004;52:478-81.

How to cite this article: Tauro LF, Suhith G, Shetty P, Rao D. Cirroid aneurysm of scalp. *J Neurosci Rural Pract* 2012;3:95-6.

Source of Support: Nil. **Conflict of Interest:** None declared.