Role of endoscopy in evaluating upper gastrointestinal tract lesions in rural population

Sharanabasavaraj Javali, M. Madan¹, M. L. Harendrakumar², M. S. Mahesh¹

Department of General Surgery, Vydehi Institute of Medical Sciences, Bengaluru, ¹Departments of General Surgery and ²Pathology, Sri Devraj Urs Medical College, Kolar, Karnataka, India

Abstract

Background and Objectives: Before the advent of endoscopy direct access to the lesion for the confirmation of the diagnosis was difficult, this posed difficulty in contemplating adequate and appropriate surgery. Endoscopy as a diagnostic and therapeutic tool has grown in recent years. Upper gastrointestinal (GI) endoscopy is one of the most fascinating branch which serves not only as a means of resolving or amplifying the diagnosis made clinically or by X-ray, but also a primary diagnostic procedure for conditions not otherwise diagnosable on unoperated case. Fiber optic upper GI endoscopy has already become firmly established as a reliable, quick and inexpensive tool. This study was done to detect the upper gastrointestinal lesions in rural population of Kolar District, the distribution pattern of various upper GI lesions in patients presenting with upper GI symptoms and to follow the endoscopic diagnosis for medical and surgical management. **Materials and Methods:** The study group includes patients reporting to outpatient department and also the inpatients in wards of General Surgery and other departments, who have upper GI symptoms, were advised endoscopy at R. L. JALAPPA Hospital and Research Centre, Kolar, from a period of December 2011 to August 2013. **Results:** Of the 600 cases, 370 were males, and 230 were females. Disease incidence was highest in 51–70 years age group, that is, 21.6%. Pain abdomen was the most common symptom. Epigastric tenderness was the most common sign among the patients clinically. Reflux esophagitis and diffuse gastritis formed most common cases (307 cases). The incidence of duodenitis - 7.83%, peptic ulcer -3.3%, esophageal varices - 1.5%, the incidence of carcinoma esophagus and carcinoma stomach was approximately same that is, 4.5% and 4.6% respectively. The incidence of esophageal candidiasis was 4.16%. The majority of the patients had a normal study that is, 14.5%. **Conclusion:** Upper GI lesions were more common in males. The incidence of diseases was highest among the elderly age group. Most of the benign mucosal lesions were mainly due to spicy food and habit of tobacco consumption. The incidence of malignancy was mostly among older age group above 50 years. The incidence of the normal study was high owing to increased medical care, easy availability of the procedure and increased medical awareness among patients. In all these cases, upper GI endoscopy not only helped in diagnosing the disease but also helped to get information about pathology, extent of disease and complications that have occurred. This study highlights the importance of diagnostic and therapeutic uses, recording of the various gastroenterological diseases we come across in rural population.

Key words

Complications, endoscopy, extent of disease, pathology

Address for correspondence:

Dr. Sharanabasavaraj Javali, Department of General Surgery,
Vydehi Institute of Medical Sciences and Research Centre,
Bengaluru, Karnataka, India.
E-mail: dr.s.c.javali@gmail.com

Access this article online

Website:

www.jdeonline.in

DOI:

10.4103/0976-5042.159238

Quick Response Code

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Javali S, Madan M, Harendrakumar ML, Mahesh MS. Role of endoscopy in evaluating upper gastrointestinal tract lesions in rural population. J Dig Endosc 2015;6:59-65.

Introduction

What is endoscopy is it an instrument or technique or it is a “Revolution or Evolution”? Many have come to understand the meaning of endoscopy as merely that of technology or instrumentation. Diagnosis and localization of its afflictions relied for many decades on barium radiology, which provides indirect data in black and white. The situation has changed dramatically since the late 1960s with the introduction of fully flexible and maneuverable endoscopes. Upper gastrointestinal (GI) lesions include patients presenting with symptoms of acid-peptic disease recurrent vomiting, dysphagia, hematemesis, etc., The prevalence of gastro-esophageal reflux diseases is more in western countries compared to Asia. The complications such as duodenal ulcers and strictures, Barrett esophagus and esophageal adenocarcinoma, can be greatly reduced if early diagnosis and screening are done by endoscopy.^[1]

Peptic ulcer (duodenal and gastric ulcer) is common in men and prevalence is increased with age with a peak prevalence of 28.8% in 5th decade of life.^[2] The role of endoscopy in bleeding peptic ulcer disease is significant. It is an effective tool in diagnosis, prognostication, and therapy of bleeding peptic ulcer.

The studies have shown that endoscopic intervention has led to a reduction in a blood transfusion, shortened Intensive Care Unit and hospital stays. It has decreased the need for surgery and lowered the mortality rate.^[3,4]

In this rural population of Kolar, we routinely encounter patients with upper GI symptoms. Hence, the need for the study is early detection of upper GI lesions by endoscopy and later evaluating the disease for appropriate management.

Materials and Methods

This study entitled “Role of endoscopy in evaluating upper GI tract lesions in rural population” was conducted in 600 cases at R. L. Jalappa and Research Center Kolar during the year December 2011 to September 2013.

In all the cases, the endoscope used was the fiberoptic OLYMPUS endoscope connected to DELL LCD monitor 21” PC.

Patients were selected at random for the study and fell into two categories:

- Inpatients-referred and emergency
- Outpatients received through OPD.

In some of the cases, the radiological examination was done prior to endoscopy. In both the categories, the cases which presented with the history and clinical findings suggestive of upper GI tract disorders were accepted for endoscopic examinations.

Symptoms

- Upper abdominal pain vomiting
- Hematemesis/melaena
- Dysphagia
- Lump in abdomen
- Anorexia/weight loss
- Sensation of fullness after meals
- Weakness and tiredness
- Retrosternal burning with regurgitation.

Signs

- Epigastric tenderness
- Supraclavicular lymphnodes
- Lump in upper abdomen
- Visible upper abdominal peristalsis
- Ascites
- Jaundice.

Investigations

Following investigations were routinely done before endoscopic examination:

- Hemoglobin
- Stool for occult blood
- LFT in jaundice.

Barium examination (if necessary). The examination was carried out under local anesthetic solution using lignocaine viscus 2% spray, sprayed 5 min before the procedure.

Since fiberoptic endoscopy is simple and safe, we did not encounter any complications. Inadequate or improper explanation makes patient apprehensive, this is true especially with children and elderly. It is most important to have patient's co-operation to complete the procedure safely. Both safety of the patients and instrument are important for the endoscopist. Positioning of the patient is also equally important for better orientation during visualization of the interior, apart from passing the scope. Mild sedation definitely helps in apprehensive patients, but sometimes it may make the patient more uncooperative under sedation. Good anaesthetization (local) is very essential, as this procedure may irritate pharynx and cause severe bouts of cough in some patients.

Endoscopic methods of biopsy (wherever necessary) used in our series were multiple quadrant biopsy with a biopsy forceps (especially in cases of suspected carcinoma of the stomach, esophagus, etc.

Photographic studies and video recording were done in cases of any positive and diagnostic findings.

Out of these 600 cases, which were randomly selected, few could be followed up during their course of treatment in the hospital and after discharge for a period varying from 2 months to more than a year.

Results

Of the 600 patients studied, the maximum number of patients belonged to the age group of 61 years and above (25.1%). The minimum number of patients studied belonged to the age group of 11–20 years constituting 18 cases (3%).

Of the 600 patients studied, 370 patients were males (61.6%) and 230 were females (28.4%). Of the 600 cases studied, it has been observed that the maximum number of GI lesions was in the stomach followed by esophagus and duodenum. Benign lesion in esophagus, stomach and duodenum was seen in 108 cases, 241 cases and 81 cases respectively. Malignant lesions of stomach and esophagus were seen in 28 and 27 cases respectively. Only two cases of carcinoma duodenum were seen in our study.

Distribution of malignant lesions of the upper gastrointestinal tract

Of the 600 cases studied 57 cases were diagnosed on endoscopy as malignant lesions. Of the 57 cases studied 28 cases were carcinoma stomach, of which 20 were seen in males, eight were seen in females.

Of the 57 cases studied 27 cases were carcinoma esophagus, of which 15 were seen in males, 12 were seen in females.

Table 1: Distribution of patients according to age

Age group	Number of Patients	Percentage %
11-20	18	3
21-30	97	16.1
31-40	98	16.33
41-50	129	21.5
51-60	108	18.00
61- above	151	25.1
Total	600	100.00

Table 2: Distribution of patients according to sex

Sex	Number of patients	Percentage
Male	370	61.6
Female	230	28.4
Total	600	100

Table 3: Distribution of patients according to etiology

Aetiology	Number of Patients	Percentage
Reflux oesophagitis	41	6.8
Reflux oesophagitis with gastritis	30	5.9
Oesophageal varices	09	1.5
O. Candidiasis	25	4.16
Stricture oesophagus	02	0.33
Carcinoma oesophagus.	27	4.5
Acute diffuse gastritis	74	12.33
Chronic gastritis	75	13.8
Atrophic gastritis	04	0.6
Reflux bile gastritis	36	6
Erosive gastritis	43	7.16
Pyloric stenosis	04	0.6
Chronic gastric ulcer	05	0.8
Duodenal ulcer	15	2.5
Carcinoma stomach	28	4.6
Foreign body stomach	04	0.66
Duodenitis	37	6.16
Gastroduodenitis	20	0.6
Duodenal polyp	06	1
Worm infestation	10	0.6
Carcinoma duodenum	03	0.5
Duodenal diverticula	03	0.5
Oesophageal polyp	01	0.16
Normal study	88	14.5
Recurrence	02	0.33
Pt not prepared well	02	0.33
Un co-operative pt's.	02	0.33
Indentation of stomach.	02	0.33
Indentation of oesophagus	02	0.33
Total	600	100

Table 4: distribution of benign and malignant lesions of gastrointestinal tract

Type of Lesion	Oesophagus	Stomach	Duodenum	Total
Benign	108	241	81	430
Malignant	27	28	2	57
Total	135	269	83	487

Table 5: Distribution of malignant lesions of upper gastrointestinal tract

Type and site	Male	Female	Total	%
Carcinoma of Stomach	20	8	28	4.6
Carcinoma of Oesophagus.	15	12	27	4.5
Carcinoma Duodenum.	02	0	02	0.33
Total	37	20	57	9.5

Graph 1: Age distribution

Graph 2: Distribution of patients according to sex

Graph 3: Distribution of benign and malignant lesions in gastrointestinal tract

Graph 4: Distribution of malignant lesions of the upper gastrointestinal tract

Graph 5: Distribution of patients according to treatment offered

Upper gastrointestinal endoscopy [Figures 1-8]

Figure 1: Antral carcinoma

Figure 2: Ca oesophagus Lower 1/3rd

Of the 57 cases studied two cases were carcinoma duodenum seen in males.

Out of 600 cases studied, 592 (98.6%) patients were managed conservatively, 5 (0.83%) patients received radiation, and 3 (0.3%) had endoscopic treatment of foreign body removal [Tables 1-5 and Graphs 1-5].

Discussion

This is a prospective study conducted in Department of General Surgery from December 2011 to September 2013 at

Figure 3: Ca esophagus middle 1/3rd

Figure 4: Duodenal ulcer

Figure 5: Esophageal candidiasis

Figure 6: Esophageal varices

Figure 7: Foreign body esophagus

Figure 8: Foreign body removed with snare

Sri Devaraj Urs Academy of Higher Education And Centre, R.L. Jalappa Hospital, Tamaka, Kolar. 600 cases who fulfilled the inclusion criteria were included in the study and were subjected to endoscopy.

In the present study, maximum number of cases subjected to fiber optic endoscopy belonged to the age group of 61 years and above (25.1%) and 51–60 years (21.5%). In a similar study by Ray and Pal done at Referral Railway

Hospital, Kolkata, the mean age group of the study was 51–60 years.^[5] In Kuwait series maximum number of patients were in 21–30 years of age group (67/211) adult age group, that is, 21.6% was affected more than any other age groups.^[6]

In the present study, the maximum patients subjected to endoscopy were males (61.6%). In the study by naniwadekar and Kuwait series maximum number of patients subjected to endoscopy were males (72.5% and 62% respectively). The results were similar to our study.^[6,7]

In the present study, most of the patients presented with pain abdomen followed by vomiting. In the study by Ray and Pal 9398 cases were studied, and most of the patients presented with the similar complaints of pain abdomen.^[5] Similarly in a study of Kuwait series most of the patients presented with pain abdomen.

In this study, most of the patients on upper GI endoscopy were found to have chronic gastritis and acute diffuse gastritis.

The incidence of esophagitis and gastritis was 11.8% and 39.33% respectively, our results are comparable to study done by Ray and Pal and Shennak *et al.* In esophagitis group most of the patients were in the age group of 21–30 years, the incidence of gastritis was in the age group of 60 years and above. This can be explained because of heavy alcohol and smoking consumption in the laborer group of people in this area. Alcohol and smoking lowers the pyloric pressure, and that leads to reflux of bile leading to the damage of gastric mucosal barrier.

Gastric ulcer incidence was low about 0.8% compared to other studies by Al-Nakib and Al-Liddawi and Ray and Pal which was 5.9% and 4.9%. In this rural population incidence of gastric ulcer and duodenal ulcer (2.5%) was very low, and seen in elderly age group people. This may be attributed to the widespread use of proton pump inhibitors in the elder age group people as PPI are co-prescribed with NSAID'S and antiplatelet agents used for cardiovascular, cerebrovascular and rheumatological diseases. Also, the PPI are available easily over the counter without cost.

The incidence of gastric carcinoma in our study was found to be 4.6%. The M: F ratio was 2.5:1. The incidence of gastric carcinoma was 6.5% in a study by Ray and Pal.^[5] The ratio of affection of male: female was 3:1 in a study at Tata Memorial Hospital, Bombay. Database of Kidwai Memorial Institute of Oncology (KMIO) 2004–2005, Bangalore, Karnataka the incidence of gastric cancer was 9%. The maximum patients affected were in the age group of 51–70 years Mean age was 58.5 years comparable to 56 years quoted by Tata Memorial Hospital and with 53 years quoted by Nakib. Antral gastric

carcinoma was more predominant compared to carcinoma of the stomach in body and fundus.

The incidence of carcinoma esophagus was 4.5% in our series. Incidence pattern is comparable to KMIO, Bangalore with the incidence of 6.6% and Ray and Pal with the incidence of 4.6%.^[5] Out of the total 27 cases, 15 were Males and 12 were Females. In males, the maximum patients affected were in the age group of 60 years and above.

Esophageal Candidiasis in the present study, Incidence was 4.16%, that is, 25/600. It was seen more common among the male of the age group 50-60 years of the affected individuals only 6 of them were sero-positive for HIV. Candidiasis also affected other individuals (who were not sero-positive for HIV), this may be attributed to the advanced age, diabetes mellitus, alcoholism, inhaled gluco-corticoid use in asthmatics and acid suppressive therapy for patients with chronic gastritis.^[8,9] It was interesting that not all patients were symptomatic, and our study supports a prevalence of esophageal candidiasis in 20% of normal healthy individuals.^[8,9] All patients were followed and given a course of treatment with oral flucanazole 200 mg on 1st day, followed by 100 mg once daily for at least 2 weeks. Total number of normal study cases were 88 and incidence was 14.5%. Incidence was more in females compared with males. The normal study were more compared to all previous studies.^[6,7,10] The increase in normal studies of endoscopy on recent years is mainly because of increased medical care, easy availability of the procedure and increase awareness of health among the population.

The incidence of upper GI lesions namely gastroduodenitis, duodenal polyp were 3.33% and 0.5% respectively. We had two cases of stricture esophagus following corrosive acid consumption. And foreign body stomach constituted to around 0.66% (3 cases).

Financial support and sponsorship
Nil.

Conflict of interest
There are no conflict of interest.

References

1. Vakili N, van Zanten SV, Kahrilas P, Dent J, Jones R; Global Consensus Group. The Montreal definition and classification of gastroesophageal reflux disease: A global evidence-based consensus. *Am J Gastroenterol* 2006;101:1900-20.
2. Khuroo MS, Mahajan R, Zargar SA, Javid G, Munshi S. Prevalence of peptic ulcer in India: An endoscopic and epidemiological study in urban Kashmir. *Gut* 1989;30:930-4.
3. Barkun A, Bardou M, Marshall JK; Nonvariceal Upper GI Bleeding Consensus Conference Group. Consensus recommendations for managing patients with nonvariceal upper gastrointestinal bleeding. *Ann Intern Med* 2003;139:843-57.

4. Spiegel BM, Vakil NB, Ofman JJ. Endoscopy for acute nonvariceal upper gastrointestinal tract hemorrhage: Is sooner better? A systematic review. *Arch Intern Med* 2001;161:1393-404.
5. Ray G, Pal S. Trends in endodiagnosis of upper gastrointestinal diseases at a referral railway hospital. *J Dig Endosc* 2011;2:213-9.
6. Al-Nakib B, Al-Liddawi H. Upper gastrointestinal endoscopy experience in Kuwait: Analysis of 1019 cases. *Gastrointest Endosc* 1981;23:605-7.
7. Shroff CP, Nanivadekar SA. Endoscopic brushing cytology and biopsy in the diagnosis of upper gastrointestinal tract lesions. A study of 350 cases. *Acta Cytol* 1988;32:455-60.
8. Mimidis K, Papadopoulos V, Margaritis V, Thomopoulos K, Gatopoulou A, Nikolopoulou V, *et al.* Predisposing factors and clinical symptoms in HIV-negative patients with *Candida* oesophagitis: Are they always present? *Int J Clin Pract* 2005;59:210-3.
9. Underwood JA, Williams JW, Keate RF. Clinical findings and risk factors for *Candida* esophagitis in outpatients. *Dis Esophagus* 2003;16:66-9.
10. Shennak MM, Tarawneh MS, Al-Sheikh TM. Upper gastrointestinal diseases in symptomatic Jordanians: A prospective endoscopic study *Ann Saudi Med* 1997;17:4.