Prevalence of elongated styloid process in Saudi population of Aseer region

Mohammed Asif Shaik¹, Naheeda², Sultan Mohammed Kaleem¹,
Abdul Wahab³, Shahul Hameed¹

¹Department of Maxillofacial Diagnostic Sciences, College of Dentistry, King Khalid University, Abha, Kingdom of Saudi Arabia,
²Department of Periodontics, Mamatha Dental College, Khammam, Andhra Pradesh, India,
³Department of Preventive Dental Sciences, College of Dentistry, King Khalid University, Abha, Kingdom of Saudi Arabia

Correspondence: Dr. Naheeda
Email: dr.naheeda@gmail.com

ABSTRACT

Objective: The study was performed to investigate the prevalence, morphology and calcification pattern of elongated styloid process in Saudi population of Aseer (Southern) region and its relation to gender and sub-age groups. **Materials and Methods:** This study was analyzed digital panoramic radiographs of 1,162 adults. Any radiograph with questionable styloid process was excluded from the study. The apparent length of the styloid process was measured by a single experienced dental and maxillofacial Radiologist. The elongated styloid process was classified with the radiographic appearance based on the morphology and calcification pattern. The data were analyzed by using Student's *t*-test and Chi-square test with *P* value less than 0.05. **Results:** A total of 1,085 Digital panoramic radiographs showed elongated styloid process of which 686 (63.2%) were noticed in males and 399 (36.8%) were noticed in female patients. There was a statistical significant difference noticed in the mean difference of elongated styloid process between 20-29, 50-59 and 60 years and above sub-age groups. The elongated styloid process was more prevalent in elderly aged male patients ($P < 0.05$). Type I morphology with calcified out line (a) was the most frequent pattern of calcification noticed in the present study. **Conclusion:** The panoramic radiographs are economical, easily accessible and useful diagnostic tool for early detection of elongated styloid process with or without symptoms. However, studies with larger sample size would further help to assess the prevalence of this elongated styloid process in Saudi population of various other regions.

Key words: Eagle's syndrome, panoramic radiographs, Saudi population, styloid process

INTRODUCTION

Styloid process is derived from the Greek word "Stylos" means a pillar, which is normally a cylindrical structure arising from the temporal bone anterior to stylo mastoid foramen.^[1] The mean radiographic length of the styloid process is usually 2-3 cm.^[2] The apex of the styloid process is clinically important as it is located between internal and external carotid arteries. The tip of the styloid process is continuous with the stylohyoid ligament, which extends to the lesser cornu of the hyoid bone.^[3] The elongation

of this styloid process or ossification of stylohyoid ligament can result in Eagle's syndrome. This was first described in 1652 by Italian surgeon Pietro Marchetti. In 1937, Eagle coined the term Stylalgia to describe the pain associated with elongation of the styloid process caused due to compression on some neural and vascular structures.^[4] It has been estimated that between 2% and 4% of the general population presents radiographic evidence of an ossified portion of the stylohyoid chain. Majority of these patients are asymptomatic.^[5] When symptoms do persist there is little correlation between the extent of the symptoms

How to cite this article: Shaik MA, N, Kaleem SM, Wahab A, Hameed S. Prevalence of elongated styloid process in Saudi population of Aseer region. Eur J Dent 2013;7:449-54.

Copyright © 2013 Dental Investigations Society.

DOI: 10.4103/1305-7456.120687

and the severity of the ossification. More commonly, panoramic radiographs are used to determine the elongation of the styloid process. However, computed tomography is used as an adjunct for the information provided by panoramic radiographs.^[5,6] Langlais *et al.* proposed a classification on elongated styloid process based on the morphology of elongation [Figures 1-3]: Type I: Elongated, Type II: Pseudo articulated, Type III: Segmented and according to pattern of calcification [Figure 4]: (a) Calcified outline, (b) partially calcified, (c) nodular, (d) completely calcified.^[7] The aim of this study was to determine the prevalence of elongated styloid process in Saudi population of the southern region and to investigate the morphology and pattern of calcification of elongated styloid process in relation to gender and sub age groups with the help of digital panoramic radiograph.

MATERIALS AND METHODS

The study is based on 1,162 digital panoramic radiographs consecutively retrieved from the archival records. All radiographs were taken between 2009 and 2011 from Department Oral Medicine and Radiology College of Dentistry, King Khalid University, Abha, Saudi Arabia. The digital radiographs of 1,162 Saudi patients with dental problems had originally been taken for routine examination and not for the investigation. Hence, it was not necessary to seek ethical approval. All the panoramic radiographs were made by using Orthopantomograph® OP200D (Instrumentarium, Finland) operated at 85 kVp and 2.5 mA with an exposure cycle of 10 s as recommended by the manufacturer. The radiographs with questionable stylohyoid complex, positioning and magnification errors and with superimposed normal anatomical structures were excluded from the study. The selected radiographs were of patient's above 10 years with the maximum age limit of 70 years. To obtain the percentiles for different age groups, the subjects were divided in to 10-19, 20-29, 30-39, 40-49, 50-59, and ≥60 sub age groups. The apparent length of the styloid process was measured by an experienced oral and maxillofacial radiologist with the help of the measuring tools on the accompanying software (Clinic view 9.2, Instrumentarium) with a magnification factor of 1.4. The length of the styloid process was measured in a similar way described by Ilgüy *et al.*^[8] Any styloid process extending below an imaginary line connecting the anterior nasal spine and the mastoid process was considered elongated. The type of the elongation and calcification of the styloid process on right and left side were evaluated according to Langlais *et al.*^[7] To check

Figure 1: Type I: elongated styloid process

Figure 2: Type II: Pseudo articulated styloid process

Figure 3: Type III: Segmented styloid process

the intraobserver variations, measurements were repeated after 1 month on a subset of 550 panoramic radiographs by the same observer. The deviation of the mean length of the styloid process between 1st and

Figure 4: Calcification pattern of elongated styloid process

2nd measurements were <2%. The observed results were analyzed using SPSS 16.0 (statistical package for social science Inc. Chicago, USA) software. Student *t*-test and Chi-square test were used for statistical analysis. *P* values less than 0.05 were accepted as statistically significant.

RESULTS

A total of 1,162 digital panoramic radiographs with dental problems were enrolled in the present study. The panoramic radiographs of 77 patients who have questionable styloid process were excluded. Therefore, 1,085 panoramic radiographs showed elongated styloid process of which, 686 (63.2%) elongated styloid process were observed in males and 399 (36.8%) were observed in females. Table 1 shows the mean difference of elongated styloid process between the gender, age and the side affected. The mean difference of elongated styloid process between the age group of 20-29, 50-59, and 60 years and above was more in males when compared to female patients, which was statistically significant ($P < 0.05$). Though, the mean difference of elongated styloid process was more in females between the age group of 40 and 49, it was not statistically significant. A statistical significant difference was noticed in the morphology of elongated styloid process in males [Table 2]. Type I morphology on the left side above 60 years of age was noticed in males. There was no significant difference noticed in the morphology of elongated styloid process in females. Calcified outline (a) [Table 3] was the most common calcified pattern noticed on right and left side of the male and right side of the female patients, which was statistically significant [Tables 4 and 5] shows the percentage distribution of morphology and calcified pattern of elongated styloid process in different sub age groups. Type I and calcified outline (a) were the most common morphology and calcified pattern in the specified sub age groups.

Table 1: Mean difference of elongated styloid process in different age groups

Age	ESPs	Male		Female		<i>P</i> value
		Mean	SD	Mean	SD	
10-19	Right side	33	1.98	32.8	1.20	0.599
	Left side	33.2	1.89	33.1	1.06	1.35
20-29	Right side	34.2	2.23	33.5	1.89	0.029*
	Left side	34.5	2.01	33.8	1.44	0.003*
30-39	Right side	34.6	2.56	34.3	2.02	0.27
	Left side	34.7	2.07	34.2	2.09	0.148
40-49	Right side	35.4	2.42	35.8	2.51	0.325
	Left side	35.7	2.48	36.4	2.22	0.100
50-59	Right side	36.0	2.42	35.5	2.44	0.0031*
	Left side	36.6	2.42	35.6	2.67	0.0037*
≥60	Right side	36.1	2.34	35.6	2.51	0.0039*
	Left side	36.7	2.47	35.2	2.61	0.0035*

SD: Standard deviation, ESP: Elongated styloid process **t* test ($P < 0.05$)

DISCUSSION

The styloid process and ligaments are derived from the 1st and 2nd brachial arches, in addition to Reichert's cartilage. It has been demonstrated that during fetal development Reichert's cartilage links the styloid bone to the hyoid bone. In the adult, the stylohyoid ligament may retain some of its embryonic cartilage and may become partially or completely ossified. If these structures solidify they can cause pain and suffering present in symptomatic cases. The actual cause of elongation is poorly understood several theories have been proposed. It could be due to growth of osseous tissue at the insertion of stylohyoid ligament or it could be due to calcification of stylohyoid ligament due to unknown process or due to persistence of cartilaginous analog of stylohyal.^[9,10] There has always been a natural variation in the length of the styloid process as debated by various researchers. It has been previously reported that the normal radiographic length of the styloid process is 2-3 mm of which, the pathogenic importance is the spatial

Table 2: Morphology of elongated styloid process according to gender

Gender	Age	Gender age morphology on right side						P value	Morphology on left side						P value
		Type I		Type II		Type III			Type I		Type II		Type III		
		n	%	n	%	n	%		n	%	n	%	n	%	
Female	10-19	17	8.7	8	6.0	2	2.9	0.625	19	10.0	7	5.2	1	1.4	0.343
	20-29	37	19	28	21.1	12	17.1		36	18.9	25	18.2	16	22	
	30-39	31	15.8	14	10.5	14	20.0		31	16.3	21	15.3	7	9.7	
	40-49	26	13.3	16	12.0	10	14.3		23	12.2	17	12.4	12	16.7	
	50-59	37	19	28	21.1	11	15.7		36	18.9	25	18.2	15	20.8	
	≥60	48	24.2	39	29.3	21	30.0		45	23.7	42	30.7	21	29.4	
Male	10-19	32	10.1	17	5.9	1	1.4	<0.001*	36	9.8	14	6.2	0	0	<0.001*
	20-29	66	20.8	35	11.9	10	13.3		72	19.5	31	13.5	8	9.1	
	30-39	52	16.4	53	18.0	10	13.3		64	17.3	28	12.2	23	26.1	
	40-49	50	15.8	56	19.0	20	26.7		49	13.3	53	23.1	24	27.3	
	50-59	46	14.5	75	25.5	18	24.0		64	17.3	56	24.5	19	21.6	
	≥60	71	22.4	58	19.7	16	21.3		84	22.8	47	20.5	14	15.9	

*Chi-square test ($P<0.05$)**Table 3: Calcification pattern of elongated styloid process according to gender**

Gender	Age	Calcification on right side								P value	Calcification on left side								P value
		A		B		C		D			A		B		C		D		
		n	%	n	%	n	%	n	%		n	%	n	%	n	%	n	%	
Female	10-19	22	9.4	5	3.8	0	0	0	0	0.043*	20	12.2	3	2.9	0	0	4	3.9	0.106
	20-29	51	21.9	19	14.3	2	25	5	20		28	17.1	25	24	5	18	19	18.4	
	30-39	33	14.2	19	14.3	0	0	7	28		29	17.7	11	10.6	6	21.4	13	12.6	
	40-49	25	10.7	24	18.0	1	12.5	2	8.0		19	11.6	11	10.6	5	17.9	17	16.5	
	50-59	50	21.5	19	14.3	2	25.0	5	20		28	17.1	24	23.1	5	17.9	19	18.4	
	≥60	52	22.3	47	35.3	3	37.5	6	24		40	24.4	30	28.8	7	25.0	31	30.1	
Male	10-19	47	11.6	2	1.2	1	3.8	0	0	<0.001*	40	13.6	7	3.7	1	1.6	2	1.4	<0.001*
	20-29	91	22.5	18	10.5	1	3.8	1	1.2		63	21.4	34	17.9	2	3.3	12	8.5	
	30-39	77	19.0	34	19.9	0	0	4	4.8		42	14.3	20	10.5	24	39.3	29	20.6	
	40-49	36	8.9	63	36.8	17	65.4	10	11.9		45	15.3	22	11.6	0	0	59	41.8	
	50-59	60	14.8	22	12.9	4	15.4	53	63.1		34	11.6	67	35.3	27	44.3	11	7.8	
	≥60	94	23.2	32	18.7	3	11.5	16	19.0		70	23.8	40	21.1	7	11.5	28	19.9	

*Chi-square test ($P<0.05$)**Table 4: Percentage distribution of morphology of elongated styloid process in different sub age groups**

Age	Gender	Morphology on right side						Morphology on left side					
		Type I		Type II		Type III		Type I		Type II		Type III	
		n	%	n	%	n	%	n	%	n	%	n	%
10-19	Female	17	63	8	29.6	2	7.4	19	70.4	7	25.9	1	3.7
	Male	32	64	17	34.0	1	2.0	36	72.0	14	28.0	0	0
20-29	Female	37	48.1	28	36.4	12	15.6	36	46.8	25	32.5	16	20.8
	Male	66	59.5	35	31.5	10	9.0	72	64.9	31	27.9	8	7.2
30-39	Female	31	52.5	14	23.7	14	23.7	31	52.5	21	35.6	7	11.9
	Male	52	45.2	53	46.1	10	8.7	64	55.7	28	24.3	23	20.0
40-49	Female	26	50.0	16	30.8	10	19.2	23	44.2	17	32.7	12	23.1
	Male	50	39.7	56	44.4	20	15.9	49	38.9	53	42.1	24	19.0
50-59	Female	37	48.7	28	36.8	11	14.5	36	47.4	25	32.9	15	19.7
	Male	46	33.1	75	54.0	18	12.9	64	46.0	56	40.3	19	13.7
≥60	Female	45	41.7	42	38.9	21	19.4	45	41.7	42	38.9	21	19.4
	Male	71	49.0	58	40.0	16	11.0	84	57.9	47	32.4	14	9.7

position of the tip of this process.^[10] Kaufman *et al.*^[11] states that, if the length of the styloid process is more

than 30 mm on the radiograph it is considered as elongated. In the present study, the styloid process,

Table 5: Percentage distribution of calcification pattern of elongated styloid process in different sub age groups

Age	Gender	Calcification on right								Calcification on left side							
		A		B		C		D		A		B		C		D	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
10-19	Female	22	81.5	5	18.5	0	0	0	0	20	74.1	3	11.1	0	0	4	14.8
	Male	47	94	2	4.0	1	2	0	0	40	80.0	7	14.0	1	2	2	4.0
20-29	Female	51	66.2	19	24.7	2	2.6	5	6.5	28	36.4	25	32.5	5	6.5	19	24.7
	Male	91	82.0	18	16.2	1	0.9	1	0.9	63	56.8	34	30.6	2	1.8	12	10.8
30-39	Female	33	55.9	19	32.2	0	0	7	11.9	29	49.2	11	18.6	6	10.2	13	22.0
	Male	77	67.0	34	29.6	0	0	4	3.5	42	36.5	20	17.4	24	20.9	29	25.2
40-49	Female	25	48.1	24	46.2	1	1.9	2	3.8	19	36.5	11	21.2	5	9.6	17	32.7
	Male	36	28.6	63	50.0	17	13.5	10	7.9	45	35.7	22	17.5	0	0	59	46.8
50-59	Female	50	65.8	19	25.0	2	2.6	5	6.6	28	36.8	24	31.6	5	6.6	19	25.0
	Male	60	43.2	22	15.8	4	2.9	53	38.1	34	24.5	67	48.2	27	19.4	11	7.9
≥60	Female	52	48.1	47	43.5	3	2.8	6	5.6	40	37.0	30	27.8	7	6.5	31	28.7
	Male	94	64.8	32	22.1	3	2.1	16	11.0	70	48.3	40	27.6	7	4.8	28	19.3

which were longer than 30 mm were included for evaluation. There are many variations of styloid chain, including the thickness of segments, angle and direction of deviation, length of process and degree of calcification.^[12] Therefore, it is necessary to define the type of elongation and calcification of each styloid in order to describe its radiographic appearance. Hence, in order to simplify the description, Langlais *et al.*^[7] had classified styloid process based on the type of elongation and calcification. Similar classification was followed in the present study to describe the morphology and calcification of the styloid process. The panoramic radiograph rather than computed tomography are used to detect if the styloid process is elongated.^[13] Similarly, in the present study panoramic radiographs were used to identify the elongated styloid process. The average length of the styloid process was 3.67 ± 0.62 cm with a mean age of elongation was 42.95 ± 16.62 in males and 37.6 ± 14.2 in female patients. There was a significant difference in the mean elongation of the styloid process between the age group of 20-29, 50-59, and above 60 years of age. These findings were in consistent with the study conducted by Gokce *et al.* in Turkey population.^[14] In the present study, it was observed that there was an increase in the length of the styloid process with an increase in the age. These findings were in consistent with studies conducted by various authors.^[15,16] It was also observed that the styloid processes were elongated more in males when compared to females and more on the left side when compared to right side. However, these findings were not in consistent with studies conducted by various authors.^[15-17] Type I morphology was the most common type of morphology observed in male patients. However, it was not statistically

significant in females. The calcified outline (a) was the common calcified pattern was seen in males and on the right side of female patients. Similar findings were noticed by Ilgüy *et al.* in term of most common type of elongated styloid process. However, they found the partial calcification to be the most prevalent pattern. Similarly, Reddy *et al.*^[18] and Bagga *et al.*^[19] reported similar results in term of most common type of styloid process. However, they found completely calcified and partially calcified pattern most prevalent in south and north Indian population. The percentage distribution of elongated styloid process in different sub age group showed Type I (33.1-72%) as the most common type of elongation and calcified outline (28.6-94%) as the most common pattern of calcification in the specified subgroups of Saudi population.

CONCLUSION

It is important for the dentist to be aware of natural variations of the styloid process whose clinical importance is not well-understood. Styloid process elongation may often be a coincidental asymptomatic radiographic finding. Therefore, proper clinical and radiographic evaluation can detect an elongated styloid process. The panoramic radiographs are economical, easily access able and useful diagnostic aid for early detection of elongated styloid process with or without symptoms. To the best of our knowledge, this is the first of it kind of study in terms of the prevalence of elongated styloid process in Saudi population of Aseer region. However, studies with larger sample size would further help to assess the prevalence of this elongated styloid process in Saudi population of various other regions.

REFERENCES

1. Roopashri G, Vaishali MR, David MP, Baig M. Evaluation of elongated styloid process on digital panoramic radiographs. *J Contemp Dent Pract* 2012;13:618-22.
2. Bozkir MG, Boga H, Dere F. The evaluation of styloid process in panoramic radiographs in edentulous patients. *Turk J Med Sci* 1999;29:481-5.
3. Frommer J. Anatomic variations in the stylohyoid chain and their possible clinical significance. *Oral Surg Oral Med Oral Pathol* 1974;38:659-67.
4. Eagle WW. Symptomatic elongated styloid process; report of two cases of styloid process-carotid artery syndrome with operation. *Arch Otolaryngol* 1949;49:490-503.
5. Prasad KC, Kamath MP, Reddy KJ, Raju K, Agarwal S. Elongated styloid process (Eagle's syndrome): A clinical study. *J Oral Maxillofac Surg* 2002;60:171-5.
6. Jaju PP, Suvama P, Parikh N. Eagles syndrome: An enigma to dentists. *J Indian Acad Oral Med Radiol* 2007;19:424-9.
7. Langlais RP, Miles DA, Van Dis ML. Elongated and mineralized stylohyoid ligament complex: A proposed classification and report of a case of Eagle's syndrome. *Oral Surg Oral Med Oral Pathol* 1986;61:527-32.
8. Ilgüy M, Ilgüy D, Güler N, Bayirli G. Incidence of the type and calcification patterns in patients with elongated styloid process. *J Int Med Res* 2005;33:96-102.
9. Joshi V, Iyengar AR, Nagesh KS, Gupta J. Elongated styloid process: A radiographic study. *J Indian Acad Oral Med Radiol* 2007;19:498-502.
10. Okabe S, Morimoto Y, Ansai T, Yamada K, Tanaka T, Awano S, *et al.* Clinical significance and variation of the advanced calcified stylohyoid complex detected by panoramic radiographs among 80-year-old subjects. *Dentomaxillofac Radiol* 2006;35:191-9.
11. Kaufman SM, Elzay RP, Irish EF. Styloid process variation. Radiologic and clinical study. *Arch Otolaryngol* 1970;91:460-3.
12. More CB, Asrani MK. Evaluation of the styloid process on digital panoramic radiographs. *Indian J Radiol Imaging* 2010;20:261-5.
13. Keur JJ, Campbell JP, McCarthy JF, Ralph WJ. The clinical significance of the elongated styloid process. *Oral Surg Oral Med Oral Pathol* 1986;61:399-404.
14. Gokce C, Sisman Y, Ertas ET, Akgunlu F, Ozturk A. Prevalence of styloid process elongation on panoramic radiography in the Turkey population from Cappadocia region. *Eur J Dent* 2008;2:18-22.
15. Scaf G, Freitas DQ, Loffredo Lde C. Diagnostic reproducibility of the elongated styloid process. *J Appl Oral Sci* 2003;11:120-4.
16. Shah SP, Praveen NB, Syed V, Subhashini AR. Elongated styloid process: A retrospective panoramic radiograph study. *World J Dent* 2012;3:316-9.
17. Balcioglu HA, Kilic C, Akyol M, Ozan H, Kokten G. Length of the styloid process and anatomical implications for Eagle's syndrome. *Folia Morphol (Warsz)* 2009;68:265-70.
18. Reddy RS, Kiran CH, Madhavi NS, Raghavendra MN, Satish A. Prevalence of elongation and calcification patterns of elongated styloid process in South India. *J Clin Exp Dent* 2013;5:30-5.
19. Bagga MB, Kumar CA, Yeluri G. Clinicoradiologic evaluation of styloid process calcification. *Imaging Sci Dent* 2012;42:155-61.

Access this article online

Quick Response Code:

Website:
www.eurjdent.com

Source of Support: Nil.
Conflict of Interest: None declared

Announcement

Android App

Download
**Android
application**

FREE

A free application to browse and search the journal's content is now available for Android based mobiles and devices. The application provides "Table of Contents" of the latest issues, which are stored on the device for future offline browsing. Internet connection is required to access the back issues and search facility. The application is compatible with all the versions of Android. The application can be downloaded from <https://market.android.com/details?id=comm.app.medknow>. For suggestions and comments do write back to us.