Evaluation of non-surgical therapy on glutathione levels in chronic periodontitis

Pooja Palwankar¹, Minakshi Rana¹, Kapil Arora², C. Deepthy³

Correspondence: Dr. Pooja Palwankar
Email: poojapalwankar@gmail.com

¹Department of Periodontology, Manav Rachna Dental College, Faridabad, Haryana, India,

²Department of Public Health Dentistry, Manav Rachna Dental College, Faridabad, Haryana, India,

³Department of Biochemistry, Manav Rachna Dental College, Faridabad, Haryana, India

ABSTRACT

Objective: To compare the levels of glutathione (GSH), both oxidized and reduced forms in patients with and without chronic periodontitis in gingival crevicular fluid (GCF). **Materials and Methods:** Twenty GCF samples from maxillary quadrants were collected using capillary micropipettes from the chronic periodontitis patients (test group) at baseline before treatment, at 1-month, 3 months, and 6 months after scaling and root planing and samples from 20 patients without chronic periodontitis (control group) from maxillary quadrants were also collected. GSH, oxidized glutathione (GSSG) levels and GSH: GSSG ratios were determined using the spectrophotometric method. **Statistical Analysis:** Results were concluded for the test over control groups using paired Student's *t*-test. **Results:** Lower concentrations of GSH ($P < 0.001$) and GSSG ($P < 0.001$) were detected in GCF in patients with chronic periodontitis (test group) than patients without chronic periodontitis (control group) at baseline. Treatment had a significant effect in improving the GSH and reducing GSSG levels postscaling and root planing at 1-month and 3 months but not significant effect at 6 months. Scaling and root planing increased the GSH: GSSG ratio ($P < 0.001$) in the test group as compared to the control group ($P < 0.001$). **Conclusions:** The concentrations of GSH within GCF are reduced in chronic periodontitis patients. Scaling and root planing (nonsurgical therapy) restores GSH concentration in GCF post 1-month and 3 months along with redox balance (GSH: GSSG ratio), but at 6 months the balance is not maintained. Adjunctive use of micronutritional supplements to boost antioxidant concentration in tissues by preserving GSH or by elevating its level at the inflamed sites is recommended, as nonsurgical periodontal therapy alone is not able to maintain redox balance for longer duration.

Key words: Chronic periodontitis, oxidized glutathione, redox balance, reduced glutathione

INTRODUCTION

Periodontitis is a chronic inflammatory disease of supporting tissues of the teeth in response to noxious stimuli whether mechanical, chemical or infectious, resulting in progressive destruction of periodontal apparatus, hence leading to pocket formation, recession or both.^[1-5] The tissue destruction in periodontitis has been attributed to the production of enzymes and reactive oxygen species (ROS) by polymorphonuclear cells and other cells. It has been demonstrated that patients with periodontitis have higher oxidative DNA and lipid damage biomarkers and lower antioxidant (AO) enzymatic activities in saliva than healthy subjects.^[6] There are a variety of defense mechanisms in the body to combat excessive ROS production, and AOs are one among these. AOs are those substances when

present at a low concentration compared to those of an oxidizable substrate significantly delays or prevents oxidation of the substrate.^[7] They act at three different levels - prevention, interception, and repair.^[8] Superoxide and hydrogen peroxide are the main oxidative species produced, which are either

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Palwankar P, Rana M, Arora K, Deepthy C. Evaluation of non-surgical therapy on glutathione levels in chronic periodontitis. Eur J Dent 2015;9:415-22.

DOI: 10.4103/1305-7456.163226

enzymatically eliminated by preventive AOs, or metal ions are sequestered, hence preventing Fenton reactions and subsequent hydroxyl radical formation^[9] e.g. – catalase, glutathione (GSH) peroxidase, 7S-transferase. Interception involves scavenging/chain breaking AOs which inhibit chain initiation, chain propagation and also lipid peroxidation e.g. – lipophilic substances such as ubiquinol, Vitamin A, Vitamin E, carotenoids and hydrophilic substances like uric acid, ascorbic acid, albumin, and bilirubin. At repair level various AOs function by repairing the damaged and reconstituting membranes. These include DNA repair enzymes, protease, transferase, and lipase.

When the level of ROS increases intracellularly, the cellular AO defenses are insufficient to maintain these harmful molecules, this condition is generally referred to as “oxidative stress.” The concept of “oxidative stress” dates back to 1986 and was elaborated as the relation between free radicals and disease.^[10] GSH is a ubiquitous tripeptide made from the combination of three amino acids, that is, cysteine, glutamate, and glycine. It is a low molecular weight thiol (up to 5–10 mM) present in the cell and existing in two forms which are oxidized glutathione (GSSG) and reduced GSH forms.^[11] Out of the total GSH present in the body, the reduced GSH constitutes about 90%, and GSSG is 10%. For survival of cell, it is imperative to maintain optimal GSH: GSSG ratios. Oxidative damage results when there is a deficiency of reduced GSH in the cell, which puts it at risk for oxidative damage.^[11] Reduced GSH plays three major functions in the body.^[12]

- Anti-oxidant function [Figure 1]
- Detoxification function
- Immune function.

The purpose of this study is to compare the levels of GSH, both oxidized and reduced forms in patients with and without chronic periodontitis in gingival crevicular fluid (GCF).

MATERIALS AND METHODS

Study groups and design

This study was a 6 months randomized case-control study, which was conducted at the Department of Periodontology. An approval for the study was obtained from the ethical committee, and written informed consent was obtained before enrolling the subjects for the study. The study enrolled 40 patients, which included patients with chronic periodontitis $n = 20$ (test group) and patients

Figure 1: Schematic representation of the biochemical interactions between neutrophil superoxide and glutathione

without chronic periodontitis $n = 20$ (control group). Subjects with no relevant medical history of tobacco usage were enrolled for the study. Subjects in the test group were selected on the basis of clinical criteria of having at least two nonadjacent sites per quadrant with probing pocket depths ≥ 5 mm, along with bleeding on probing and demonstrable radiographic bone loss. Control patients did not show evidence of attachment loss or probing pocket depths ≥ 3 mm and bleeding scores were below 10%. The subjects with a history of taking vitamin supplements, anti-inflammatory or antibiotic medication in the preceding 3 months and pregnant ladies and patient with special dietary needs were not included in the study. The study was started in January 2013 and ended in July 2014. After enrollment, baseline GCF samples were collected, before recording clinical measures which included Silness and Loe plaque index, Loe and Silness gingival index, Muhlemann and Son sulcus bleeding index and probing pocket depth. After data collection, chronic periodontitis patients (test group) underwent scaling and root planing and oral hygiene instruction which included brushing technique, dental flossing were reinforced at baseline. Patient without chronic periodontitis (control group) received only oral hygiene instructions. Subjects were then recalled at 1-month, 3 months and 6 months posttherapy for recording clinical and biochemical parameters.

Gingival crevicular fluid collection and glutathione estimation

GCF samples were collected from mesiobuccal/distolingual sites on any teeth in the maxillary quadrant (test and control group) using microcapillary pipettes. GCF samples were then

immediately transferred to top sealed cuvettes to prevent oxidation of AOs present in it. Levels of reduced and GSSG were analyzed using “beutler” spectrophotometric test.^[13]

Data analysis

All the results of this study were obtained by comparing intragroup and intergroup parameters, at various designated phases at baseline, 1st month, 3rd month and 6th month for test over control groups using paired Student’s *t*-test using software Computer software SPSS version 20 (IBM SPSS Inc, Chicago, IL, USA).

RESULTS

Clinical data

The nonsurgical periodontal therapy provided to the test group resulted in observed reductions of whole mouth mean plaque scores ($P < 0.001$), mean gingival scores, and mean sulcus bleeding index scores in the group ($P < 0.001$) Which in turn showed reduced inflammation as compared to control group [Tables 1-6].

Glutathione levels in gingival crevicular fluid

In all groups, mean reduced GSH and GSSG levels were detected in the millimolar range (range: 0.30–5.14 mM [Table 7]). Lower levels of GSH and GSSG concentrations were reported in the GCF from chronic periodontitis patients before and after treatment, compared with those detected in nonchronic periodontitis subjects (control group) [Table 8].

Comparative analysis of difference in mean GSH and GSSG value for test group was done at different time intervals, namely, baseline - 1-month, baseline - 3 months, baseline - 6 months, 1–3 months and 1–6 months. The differences were observed to be statistically significant ($P < 0.001$) at a confidence interval of 95%, but statistically nonsignificant at the 3–6 months interval [Tables 9 and 10]. Comparison of mean GSH and GSSG score with a standard deviation between test and control group is depicted [Figures 2 and 3] GSH and GSSG value for control group were depicted at different time intervals, namely, baseline - 1 month, baseline - 3

Table 1: Intra-group comparison of mean plaque scores and SD in test group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	22	1.98	0.23	0.92	0.50	9.178	<0.001	Statistically significant
Baseline-3 rd month	21	2.00	0.21	0.84	0.53	9.483	<0.001	Statistically significant
Baseline-6 th month	21	2.00	0.21	0.86	0.49	9.635	<0.001	Statistically significant
1 st month-3 rd month	21	0.90	0.51	0.84	0.53	0.835	0.414	Statistically nonsignificant
1 st month-6 th month	21	0.90	0.51	0.86	0.49	0.393	0.699	Statistically nonsignificant
3 rd month-6 th month	21	0.84	0.53	0.86	0.49	0.419	0.68	Statistically nonsignificant

SD: Standard deviation

Table 2: Intra-group comparison of mean gingival index scores and SD in test group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	22	2.23	1.31	0.86	0.49	4.25	<0.001	Statistically significant
Baseline-3 rd month	21	2.25	1.34	0.87	0.56	4.042	0.001	Statistically nonsignificant
Baseline-6 th month	21	2.25	1.34	1.00	0.67	3.471	0.002	Statistically nonsignificant
1 st month-3 rd month	21	0.84	0.49	0.87	0.56	0.528	0.603	Statistically nonsignificant
1 st month-6 th month	21	0.84	0.49	1.00	0.67	1.778	0.091	Statistically nonsignificant
3 rd month-6 th month	21	0.87	0.56	1.00	0.67	1.961	0.064	Statistically nonsignificant

SD: Standard deviation

Table 3: Intragroup comparison of mean SBI scores and SD in test group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	22	2.20	0.79	0.93	0.48	7.341	<0.001	Statistically significant
Baseline-3 rd month	21	2.21	0.81	0.94	0.63	4.968	<0.001	Statistically significant
Baseline-6 th month	21	2.21	0.81	0.85	0.63	5.214	<0.001	Statistically significant
1 st month-3 rd month	21	0.90	0.47	0.94	0.63	0.375	0.712	Statistically nonsignificant
1 st month-6 th month	21	0.90	0.47	0.85	0.63	0.41	0.686	Statistically nonsignificant
3 rd month-6 th month	21	0.94	0.63	0.85	0.63	2.29	0.033	Statistically nonsignificant

SD: Standard deviation, SBI: Sulcus bleeding index

Table 4: Intragroup comparison of mean plaque scores and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	20	0.23	0.15	0.19	0.26	0.654	0.521	Statistically nonsignificant
Baseline-3 rd month	20	0.23	0.15	0.24	0.43	0.146	0.886	Statistically nonsignificant
Baseline-6 th month	20	0.23	0.15	0.13	0.10	2.747	0.013	Statistically nonsignificant
1 st month-3 rd month	20	0.18	0.25	0.24	0.42	0.566	0.578	Statistically nonsignificant
1 st month-6 th month	20	0.18	0.25	0.13	0.10	0.993	0.333	Statistically nonsignificant
3 rd month-6 th month	20	0.24	0.42	0.13	0.10	1.173	0.255	Statistically nonsignificant

SD: Standard deviation

Table 5: Intragroup comparison of mean gingival index scores and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	20	0.20	0.09	0.21	0.27	0.235	0.817	Statistically nonsignificant
Baseline-3 rd month	20	0.20	0.09	0.21	0.23	0.223	0.826	Statistically nonsignificant
Baseline-6 th month	20	0.20	0.09	0.15	0.11	1.466	0.159	Statistically nonsignificant
1 st month-3 rd month	20	0.21	0.27	0.21	0.23	0.101	0.92	Statistically nonsignificant
1 st month-6 th month	20	0.21	0.27	0.15	0.11	1.241	0.23	Statistically nonsignificant
3 rd month-6 th month	20	0.21	0.23	0.15	0.11	1.375	0.185	Statistically nonsignificant

SD: Standard deviation

Table 6: Intragroup comparison of mean sulcus bleeding index score and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	20	0.16	0.08	0.19	0.23	0.36	0.722	Statistically nonsignificant
Baseline-3 rd month	20	0.16	0.08	0.17	0.21	0.09	0.929	Statistically nonsignificant
Baseline-6 th month	20	0.16	0.08	0.16	0.20	0.908	0.375	Statistically nonsignificant
1 st month-3 rd month	20	0.19	0.23	0.17	0.21	0.845	0.409	Statistically nonsignificant
1 st month-6 th month	20	0.19	0.23	0.16	0.20	0.841	0.411	Statistically nonsignificant
3 rd month-6 th month	20	0.17	0.21	0.16	0.20	0.882	0.389	Statistically nonsignificant

SD: Standard deviation

Table 7: Mean ratio of reduced glutathione and oxidized glutathione and SD in test group

Time interval	Number of subjects	Mean (Umol/dl)	SD	Mean (Umol/dl)	SD	<i>t</i>	<i>P</i>	Inference
Baseline-1 st month	22	0.30	0.19	1.76	0.95	7.129	<0.001	Statistically significant
Baseline-3 rd month	21	0.31	0.19	4.33	1.67	10.855	<0.001	Statistically significant
Baseline-6 th month	21	0.31	0.19	5.14	1.55	14.274	<0.001	Statistically significant
1 st month-3 rd month	21	1.79	0.96	4.33	1.67	11.337	<0.001	Statistically significant
1 st month-6 th month	21	1.79	0.96	5.14	1.55	9.513	<0.001	Statistically significant

SD: Standard deviation

months, baseline - 6 months, 1-3 months and 1-6 months in Tables 11 and 12.

Ratio of reduced GSH and GSSG for test group at different time interval during the study was reported to be statistically significant ($P < 0.001$) at confidence interval of 95% at baseline - 1-month, baseline - 3 months, baseline - 6 months, 1-3 months, 1-6 months [Table 7]. Comparison of the mean ratio of reduced and oxidized GSH (GSH: GSSG) score with a standard deviation between test and control group is depicted in Figure 4. The GSH and GSSG ratio values for control group were shown in Table 13. This improvement in the GSH: GSSG ratio after

treatment was as a result of an increase in levels of reduced GSH and a reduction in levels of oxidized GSH.

DISCUSSION

The present data confirm the study conducted by Chapple *et al.*, 2002 suggesting that a millimolar concentration of GSH was present in GCF, which decreases during periodontitis. The reduction in GSH levels is suggestive of its protective role for vital cells and tissue structures from host-derived free radicals. Due to the conversion of GSH to GSSG, there is a reduction in GSH levels at inflamed tissue sites as

Table 8: Intergroup comparison between various parameters used in study between test and control group

	Groups	n	Mean	SD	t	P
GSH (B)	Test group	22	108.67	42.39	15.486	<0.001
	Control group	20	519.34	116.29		
GSSG (B)	Test group	22	318.41	139.69	8.538	<0.001
	Control group	20	50.80	9.97		
GSH_GSSG (B)	Test group	22	0.30	0.19	15.7	<0.001
	Control group	20	10.73	3.11		
GSH (1-month)	Test group	22	233.91	107.74	8.683	<0.001
	Control group	20	525.15	109.48		
GSSG (1-month)	Test group	22	145.86	76.22	5.34	<0.001
	Control group	20	53.70	12.27		
GSH_GSSG (1-month)	Test group	22	1.76	0.95	13.011	<0.001
	Control group	20	10.31	2.92		
GSH (3 month)	Test group	21	362.00	145.30	4.567	<0.001
	Control group	20	533.65	86.48		
GSSG (3 month)	Test group	21	89.43	38.09	4.248	<0.001
	Control group	20	51.85	10.88		
GSH_GSSG (3 month)	Test group	21	4.33	1.67	9.679	<0.001
	Control group	20	10.70	2.48		
GSH (6 month)	Test group	21	386.38	109.20	5.576	<0.001
	Control group	20	555.50	82.40		
GSSG (6 month)	Test group	21	80.48	27.33	4.141	<0.001
	Control group	20	52.60	12.91		
GSH_GSSG (6 month)	Test group	21	5.14	1.55	8.895	<0.001
	Control group	20	11.08	2.61		
Plaque index (B)	Test group	22	1.98	0.23	28.667	<0.001
	Control group	19	0.23	0.15		
SBI (B)	Test group	22	2.20	0.79	11.482	<0.001
	Control group	20	0.16	0.08		
Gingival index (B)	Test group	22	2.23	1.31	6.938	<0.001
	Control group	20	0.20	0.09		
Plaque index (1-month)	Test group	22	0.92	0.50	5.9	<0.001
	Control group	20	0.18	0.25		
SBI (1-month)	Test group	22	0.93	0.48	6.336	<0.001
	Control group	20	0.19	0.23		
Gingival index (1-month)	Test group	22	0.86	0.49	5.233	<0.001
	Control group	20	0.21	0.27		
Plaque index (3 month)	Test group	21	0.84	0.53	3.996	<0.001
	Control group	20	0.24	0.42		
SBI (3 month)	Test group	21	0.94	0.63	5.243	<0.001
	Control group	20	0.17	0.21		
Gingival index (3 month)	Test group	21	0.87	0.56	4.919	<0.001
	Control group	20	0.21	0.23		
Plaque index (6 month)	Test group	21	0.86	0.49	6.558	<0.001
	Control group	20	0.13	0.10		
SBI (6 month)	Test group	21	0.85	0.63	0.803	0.427
	Control group	20	0.52	1.76		
Gingival index (6 month)	Test group	21	1.00	0.67	5.539	<0.001
	Control group	20	0.15	0.11		

SD: Standard deviation, GSH: Reduced glutathione, GSSG: Oxidized glutathione, SBI: Sulcus bleeding index, B: Baseline

also seen in this study. Similarly, the reduced GSH: GSSG ratio at baseline was due to reduced levels of GSH as well as increased accumulation of GSSG. These results were consistent with findings reported in earlier studies demonstrating a reduction in total

antioxidant capacity of saliva in patients with chronic periodontitis along with reduced levels of GSH.^[15-17]

In this study, a significant increase in GSH levels along with improvement in GSH: GSSG ratio at 1-month was

Table 9: Intragroup comparison of mean reduced glutathione level and SD in test group

Time interval	Number of subjects	Mean (Umol/dl)	SD	Mean (Umol/dl)	SD	t	P	Inference
Baseline-1 st month	22	108.67	42.39	233.91	107.74	7.64	<0.001	Statistically significant
Baseline-3 rd month	21	112.15	40.10	362.00	145.30	9.415	<0.001	Statistically significant
Baseline-6 th month	21	112.15	40.10	386.38	109.20	13.262	<0.001	Statistically significant
1 st month-3 rd month	21	240.48	105.79	362.00	145.30	8.047	<0.001	Statistically significant
1 st month-6 th month	21	240.48	105.79	386.38	109.20	8.746	<0.001	Statistically significant

SD: Standard deviation

Table 10: Intragroup comparison of mean oxidized glutathione level and SD in test group

Time interval	Number of subjects	Mean (Umol/dl)	SD	Mean (Umol/dl)	SD	t	P	Inference
Baseline-1 st month	22	318.41	39.69	233.91	76.22	6.769	<0.001	Statistically significant
Baseline-3 rd month	21	326.76	137.39	362.00	38.09	8.755	<0.001	Statistically significant
Baseline-6 th month	21	326.76	137.39	386.38	27.33	8.926	<0.001	Statistically significant
1 st month-3 rd month	21	148.52	77.04	145.86	38.09	5.119	<0.001	Statistically significant
1 st month-6 th month	21	148.52	77.04	89.43	27.33	4.721	<0.001	Statistically significant
3 rd month-6 th month	21	89.43	38.09	80.48	27.33	1.565	0.133	Statistically nonsignificant

SD: Standard deviation

Table 11: Intragroup comparison of mean reduced glutathione level and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	t	P	Inference
Baseline-1 st month	20	519.34	116.29	525.15	109.48	0.424	0.676	Statistically nonsignificant
Baseline-3 rd month	20	519.34	116.29	533.65	86.48	0.905	0.377	Statistically nonsignificant
Baseline-6 th month	20	519.34	116.29	555.50	82.40	1.722	0.101	Statistically nonsignificant
1 st month-3 rd month	20	525.15	109.48	533.65	86.48	0.602	0.554	Statistically nonsignificant
1 st month-6 th month	20	525.15	109.48	555.50	82.40	1.834	0.082	Statistically nonsignificant

SD: Standard deviation

Table 12: Intragroup comparison of mean oxidized glutathione level and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	t	P	Inference
Baseline-1 st month	20	50.80	9.97	53.70	12.27	0.998	0.331	Statistically nonsignificant
Baseline-3 rd month	20	50.80	9.97	51.85	10.88	0.434	0.67	Statistically nonsignificant
Baseline-6 th month	20	50.80	9.97	52.60	12.91	0.755	0.459	Statistically nonsignificant
1 st month-3 rd month	20	53.70	12.27	51.85	10.88	0.96	0.349	Statistically nonsignificant
1 st month-6 th month	20	53.70	12.27	52.60	12.91	0.641	0.529	Statistically nonsignificant
3 rd month-6 th month	20	51.85	10.88	52.60	12.91	0.408	0.688	Statistically nonsignificant

SD: Standard deviation, GSSG: Oxidized glutathione

Table 13: Intragroup comparison of mean ratio of reduced glutathione and oxidized glutathione and SD in control group

Time interval	Number of subjects	Mean	SD	Mean	SD	t	P	Inference
Baseline-1 st month	20	10.73	3.11	10.31	2.92	0.852	0.405	Statistically nonsignificant
Baseline-3 rd month	20	10.73	3.11	10.70	2.48	0.056	0.956	Statistically nonsignificant
Baseline-6 th month	20	10.73	3.11	11.08	2.61	0.893	0.383	Statistically nonsignificant
1 st month-3 rd month	20	10.31	3.11	10.70	2.48	1.181	0.252	Statistically nonsignificant
1 st month-6 th month	20	10.31	2.92	10.70	2.61	2.192	0.041	Statistically nonsignificant

SD: Standard deviation

observed following nonsurgical periodontal therapy viz., scaling and root planing, which is consistent with the observations made by several others studies.^[18-20] These results were suggestive of an improvement in AO status and reduction in oxidative stress following

nonsurgical periodontal therapy. Other possibilities for the improvement seen following nonsurgical therapy is the reduction in proteolytic activity of microorganisms and also reductions in inflammation posttherapy, thereby lowering levels of oxidants

Figure 2: Reduced glutathione levels at different time intervals for test and control groups

Figure 3: Oxidized glutathione levels at different time intervals for test and control groups

Figure 4: Ratio of reduced to oxidized glutathione levels at different time intervals for test and control groups

in GCF, both exogenously and endogenously. The improvement was seen in GSH: GSSG ratio at

3 months is due to relatively long duration taken by subgingival microflora to reestablish, hence their reduced activity.^[2]

The preliminary findings of studies stated that millimolar concentrations of GSH is present in GCF gets significantly reduced in chronic periodontitis patients.^[14,19] In addition, this study demonstrated that successful, nonsurgical periodontal therapy does restore the redox balance (GSH: GSSG) till 3 months, but at 6 months no significant improvement in either GSH or GSH: GSSG ratio was noted. Thus after 3 months thorough oral hygiene practice is not self-sufficient for restoring the GSH levels and GSH: GSSG ratio to health, this implies that there is a reduced potential of periodontal tissues against ROS activity in chronic periodontitis patients, even after successful nonsurgical therapy. Such findings have opened up the potential of using pharmacological agents to elevate buffering capacity within tissues by elevating GSH levels, e.g. use of the GSH promoting drug N-acetyl cysteine. This pharmacological approach is currently under investigation in the management of rheumatoid arthritis.^[20,21] Anti-inflammatory tissue redox state can be created by supplementing the individuals with micronutrients for elevating GSH concentrations.^[22] For various chronic inflammatory diseases that are associated with redox imbalance, AO supplementation is actively pursued as a preventive and therapeutic measure.

Limitations

Duration of the study, the reliability of the spectroscopic method used for GSH estimation and method of GCF collection. This study opens up scope for various future studies dealing with the role of GSH as biological diagnostic disease marker, AO, detoxifier, and immune modulator. Long-term studies can be undertaken to evaluate the efficacy of novel therapeutic approaches for improving the buffering capacity within periodontal tissues by elevating the GSH levels.

CONCLUSIONS

The study concluded that at baseline due to prevailing oxidative stress the levels of reduced GSH were reduced in patients with chronic periodontitis. Following nonsurgical therapy the levels of GSH were improved at 1-month and 3 months. At 6 months recall following nonsurgical therapy, an adjunctive use of micronutritional supplements to boost AO concentration in tissues by preserving GSH or by elevating its level at the inflamed sites is recommended.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL Jr. Microbial complexes in subgingival plaque. *J Clin Periodontol* 1998;25:134-44.
2. Newman MG, Takei HH, Klokkevoid PR, Carranza FA. Classification of disease and conditions affecting the periodontium. Carranza's Clinical Periodontology. 10th ed. St. Louis, Missouri: Saunders; 2009. p. 104.
3. Fredriksson M, Gustafsson A, Asman B, Bergström K. Hyper-reactive peripheral neutrophils in adult periodontitis: Generation of chemiluminescence and intracellular hydrogen peroxide after *in vitro* priming and FcγR-stimulation. *J Clin Periodontol* 1998;25:394-8.
4. Gustafsson A, Asman B, Bergström K. Priming response to inflammatory mediators in hyperreactive peripheral neutrophils from adult periodontitis. *Oral Dis* 1997;3:167-71.
5. Loe H, Theilade E, Jensen SB. Experimental gingivitis in man. *J Periodontol* 1965;36:177-87.
6. Canakci CF, Cicek Y, Yildirim A, Sezer U, Canakci V. Increased levels of 8-hydroxydeoxyguanosine and malondialdehyde and its relationship with antioxidant enzymes in saliva of periodontitis patients. *Eur J Dent* 2009;3:100-6.
7. Halliwell B. Reactive oxygen species in living systems: Source, biochemistry, and role in human disease. *Am J Med* 1991;91:14S-22S.
8. Battino M, Bullon P, Wilson M, Newman H. Oxidative injury and inflammatory periodontal diseases: The challenge of anti-oxidants to free radicals and reactive oxygen species. *Crit Rev Oral Biol Med* 1999;10:458-76.
9. Sies H. Oxidative stress, angewandte chemie int . London academic press 1986; 25:1058-71.
10. Cnubben NH, Rietjens IM, Wortelboer H, van Zanden J, van Bladeren PJ. The interplay of glutathione-related processes in antioxidant defense. *Environ Toxicol Pharmacol* 2001;10:141-52.
11. Townsend DM, Tew KD, Tapiero H. The importance of glutathione in human disease. *Biomed Pharmacother* 2003;57:145-55.
12. Meister A, Anderson ME. Glutathione. *Annu Rev Biochem* 1983;52:711-60.
13. Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J Lab Clin Med* 1963;61:882-8.
14. Chapple IL, Brock G, Eftimiadi C, Matthews JB. Glutathione in gingival crevicular fluid and its relationship to local antioxidant capacity in periodontal health and disease. *J Clin Periodontol* 2002;42:1-73.
15. Borges I Jr, Moreria EA, Filho DW, de Oliveira TB, Da Silva MB, Frode TS. Proinflammatory and oxidative stress markers in patients with periodontal disease. *Mediators Inflamm* 2007;12:457-64.
16. Tsai CC, Chen HS, Ho YP, Ho KY, Wu YM, Hou GL. Periodontopathogens and Oxidative Stress in Periodontal Diseases. Vol. 6. Paper 0595, 81st General Session of International Association for Dental Research; 2003. p. 25-8.
17. Moore S, Calder KA, Miller NJ, Rice-Evans CA. Antioxidant activity of saliva and periodontal disease. *Free Radic Res* 1994;21:417-25.
18. Guentsch A, Preshaw PM, Bremer-Streck S, Klinger G, Glockmann E, Sigusch BW. Lipid peroxidation and antioxidant activity in saliva of periodontitis patients: Effect of smoking and periodontal treatment. *Clin Oral Investig* 2008;12:345-52.
19. Grant MM, Brock GR, Matthews JB, Chapple IL. Crevicular fluid glutathione levels in periodontitis and the effect of non-surgical therapy. *J Clin Periodontol* 2010;37:17-23.
20. Reiners JJ Jr, Mathieu P, Okafor C, Putt DA, Lash LH. Depletion of cellular glutathione by conditions used for the passaging of adherent cultured cells. *Toxicol Lett* 2000;115:153-63.
21. Kelly GS. Clinical applications of N-acetylcysteine. *Altern Med Rev* 1998;3:114-27.
22. Johnston CS, Meyer CG, Srilakshmi JC. Vitamin C elevates red blood cell glutathione in healthy adults. *Am J Clin Nutr* 1993;58:103-5.

Access this article online

Quick Response Code:

Website:

www.eurjdent.com