Biomarkers in neurocritical care

Benedict Tan, Jose I. Suarez

INTRODUCTION

One of the great challenges faced by practitioners is reaching the correct diagnosis in patients with acute neurological problems. Usually, excellent history taking complemented with skilful neurological examinations will shed light on the possible diagnosis. However, in practice, the inconspicuousness of disease often eclipses the real problem, and the lack of readily reliable data may delay or misdirect the care.

Given these circumstances, biomarkers play a role and act as a surrogate measure. In a highly dynamic Neurointensive Care Unit (NICU) environment, it can help guide the treatment decisions or provide risk stratification. It is not meant to replace the traditional methods, but should rather serve as a useful adjunctive tool. We aim to review the different types, strengths and limitation of biomarkers in this setting.

DEFINITION OF BIOMARKERS

According to the Biomarkers Definitions Working Group paper, a biomarker is a characteristic, that is, objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes or pharmacologic responses to a therapeutic intervention.^[1] There are several features that are desirable for biomarkers to be useful in the Neurocritical Care Unit (NICU). They have to be brain specific; they have to increase or decrease significantly during the relevant neurological insult; they must be available within a few hours. All these features will help a treating physician to make better diagnosis and direct care.

Department of Neurology, Division of Vascular Neurology and Neurocritical Care, Baylor College of Medicine, CHI Baylor St. Luke's Medical Center, Houston, TX, USA

HISTORY OF BIOMARKERS

Since the 1950s, interests in biomarkers of neuronal injuries have increased significantly. Most of the earliest work involves studies in traumatic brain injury (TBI) and cerebral ischaemia. In 1976, Rudman *et al.* published a classic study on cerebrospinal fluid (CSF) cyclic adenosine monophosphate levels in TBI as a putative marker of the depth of coma after injury.^[2,3] In 1988, Vaagenes *et al.* studied the levels of brain creatine kinase (CK) in CSF in dogs after cardiac arrest and shows that peak activity at 48–72 h post-arrest correlated with poor outcome.^[4] This was translated to human 6 years later and found that the increase in CK activity in CSF reflects permanent brain damage.^[5] Despite the ubiquitous use of CK to investigate cardiac injury, it had never been part of the routine blood work for neurological injury.

For the next 25 years, the list of putative neurological biomarkers continued to grow as did the conflicting results of their utility in different types of brain injury. The ones that have received the most attention are S-100B and neuron-specific enolase (NSE). Both were first described in the 1960s. These biomarkers have shown effectiveness as a diagnostic and prognostic marker in the setting of wide range acute neurological diseases.

More recently, a systematic review on biomarkers recommended against the routine use of biomarkers for outcome prognostication for all disorders except hypoxic ischemic encephalopathic injuries, where NSE may be used in conjunction with clinical data for neurologic prognostication.^[6]

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Tan B, Suarez JI. Biomarkers in neurocritical care. J Neuroanaesthesiol Crit Care 2016;3:4-11.

Access this article online	
Quick Response Code:	Website: www.jnaccjournal.org
DOI: 10.4103/2348-0548.174726

APPLICATIONS IN THE NEUROINTENSIVE CARE UNIT

The majority of the patients in the NICU suffer from acute ischemic stroke (AIS), TBI, intracerebral haemorrhage (ICH) or sub-arachnoid haemorrhage (SAH). All these involve an active process going on with the nervous tissues. The outcomes of these patients often depend on how well we can mitigate the disastrous pathophysiology. Given the nature of most of the neurocritical diseases, a good neurological examination may be difficult to obtain, and the procurement of imaging studies may also be delayed for various reasons. These factors lead to the search for neuro-biomarkers. Public interest involving TBI in sports and accidents, along with the military personnel from the Iraq and Afghanistan Wars have also helped to fuel the development of biomarkers with both public and private funding.

Of the biomarkers that have been studied, NSE, S-100B, glial fibrillary acidic protein (GFAP) and ubiquitin C-terminal hydrolase L1 (UCH-L1) are the most well researched [Tables 1 and 2]. Newer biomarkers such as neurofilaments, amyloid beta, spectrin, apolipoprotein E (APOE) and many others are currently under investigation.

APPROACHES

Traditionally, researchers have focused on the investigation of a single or few molecules processes related specifically to the pathological process in the brain, such as the use of S-100B and NSE. In this review, we will discuss some of the newer approaches and what lies in the future [Figure 1].

Proteomic approach

Acute neurological diseases set into motion a complex secondary injury cascade. Utilising the proteomic approaches allow assessment of hundreds or thousands of proteins simultaneously. Several traditional techniques have been used including gel electrophoresis, mass spectrometry, antibody arrays and high-throughput immunoblotting, which are then combined with bioinformatics.^[7] For example, Jenkins *et al.* used two-dimensional gel electrophoresis in TBI and identified over 1500 proteins with 10% demonstrating a 10-fold change between injury and control conditions.^[8] Recently, serum amyloid A using the same method was identified in children with mild TBI.^[9] Overall, proteomic screenings of blood and CSF of acute neurological injuries hold promise in the identification of newer biomarkers.

Lipidomics approach

Lipid peroxidation has been long associated with brain injury.^[10] Lipidomics, first introduced by Han and Gross in 2003, is the qualitative and quantitative analyses of the lipid components from serum, plasma, tissue or

cell.^[11] Cardiolipin, a phospholipid, found exclusively in the mitochondria. Cytochrome C in the mitochondria interacts with cardiolipin and acts as a cardiolipin oxygenase. This oxygenase is activated in a variety of cellular stresses which subsequently trigger apoptotic cell death.^[12,13] Limiting factors to this approach include the origin of the lipids,^[14] and the lack of proper oxidised phospholipid standards.^[13]

Blood-based genetic marker

Several studies have shown that APOE genotype may be related to clinical outcome in brain injuries. Athletes with APOE $\epsilon 4$ reported greater symptomatology post-concussion than those without.^[15] It is also associated with an increased risk of unfavourable outcome in TBI^[16] and SAH.^[17] There is also strong associations between APOE variants and lobar ICH.^[18] Suggesting that low-density lipoprotein levels may modulate this effect.^[19] Possible limitations of this genetic marker include the fact that it may not reflect the current brain injury pathology; it may be as a result of systemic lipid metabolism, innate immunity or endocytic trafficking that act as contributors to brain injuries, as suggested in the pathology of Alzheimer's disease.^[20] Despite not being conclusive, it holds a promising future.

MicroRNAs

MicroRNAs (miRNAs) was first identified in 1993^[21] are an abundant class of short, non-coding RNA molecules, approximately 22 nucleotides (nt) in length, which regulate gene expression through RNA interference.^[22] It has been linked with neuroinflammation in acute brain injury, infection, neuroimmune and neurodegenerative diseases.^[23] Exposure to any pathogen leads to significant changes in the expression of specific miRNAs in immune cells, such as in the case of miR-155 and miR-146a, which have an important role in the Japanese encephalitis.^[23,24] Several circulating miRNAs have been shown to change in the plasma of TBI patients.^[25] Bioinformatic analysis indicates that miR-let-7i may regulate TBI-related proteins and inflammatory cytokines, including S-100B, GFAP and UCH-L1, suggesting the potential use of it as diagnostic biomarkers.^[26] Ischaemic stroke patients also display changes in the level of it.^[27] Although still in its early stages research, it holds great potentials to monitor the courses of disease.

Other approaches

The search for pathological fingerprint continues with excitement in the era of advancement in translational medicine and unfolds the possibilities of newer approaches. Multiplex bead technology provides quantitative measurement of large numbers of analytes and this will broaden attempts at discovering newer biomarkers.^[28] Exosomes are nanovesicles secreted into the blood upon internal vesicle fusion with the

Table 1: A summary of potentially useful biomarkers in the Neurocritical Care Units

Marker	Structure affected	Findings in relation to brain injury	
		CSF	Blood
S-100B	Astroglial cells	Elevated	Elevated but may be confounded by extracerebral tissues
GFAP	Astroglial cells	Elevated	Elevated
MBP	Axon	N/A	Not sensitive
NFLP	Axon	Peak at 4-10 days post-injury	N/A
NSE	Neuron	Elevated, but may be confounded by RBC lysis	
UCH-L1	Neuron	Elevated	Elevated
Amyloid beta 40, 42	Plaque pathology	No change	N/A
MMP-9	Blood-brain barrier	Elevated	Elevated
SBDP 145, 150	Membranes, cytoskeleton	Elevated	Elevated
Total tau protein	Axon	Peak at 4-10 days post-injury	Elevated

Other potential biomarkers: Leptin; glutamate; interleukins; APOE; copeptin; BDNF

NSE = Neuron-specific enolase, MBP = Myelin basic protein, GFAP = Glial fibrillary acidic protein, UCH-L1 = Ubiquitin C-terminal hydrolase, NFLP = Neurofilament light polypeptide, MMP = Matrix metalloproteinases, BDNF = Brain-derived neurotrophic factor, SBDP = Spectrin breakdown product, N/A = Not available, CSF = Cerebrospinal fluid, RBC = Red blood cell, APOE = Apolipoprotein E

Table 2: A summary of the types of tests in the Neurocritical Care Units

Diseases	Type of tests	Potential biomarkers
Intracranial hypertension	Head CT, MRI brain, ultrasound, ICP monitor	IL-6
Ischaemic stroke	Head CT, MRI DWI-ADC and angiography	S-100B
Intracerebral haemorrhage	Head CT, MRI SWI FFE, angiography	Leptin in basal ganglia haemorrhage, fibrinogen in post-tPA
Traumatic brain injury	Head CT, MRI brain, MRS and DTI	S-100B, NSE, UCH-L1, NFLP, MBP, GFAP
SAH vasospasm	TCD, CT angiography, invasive cerebral angiography and EEG	Nitrate, nitrite, ADMA, S-100B
Status epilepticus	EEG, MRI	UCH-L1, MiRNA
Cardiac injury	EKG, 2D/3D echogram and MRI heart	Troponin, CKMB

CT = Computed tomography, MRI = Magnetic resonance imaging, ICP = Intracranial pressure, DWI = Diffusion-weighted imaging, ADC = Apparent diffusion coefficient, MRS = Magnetic resonance spectroscopy, DTI = Diffusion tensor imaging, TCD = Transcranial Doppler ultrasound, EEG = Electroencephalogram, EKG = Electrocardiogram, CK = Creatine kinase, NSE = Neuron-specific enolase, MBP = Myelin basic protein, GFAP = Glial fibrillary acidic protein, UCH-L1 = Ubiquitin C-terminal hydrolase, NFLP = Neurofilament light polypeptide, MMP = Matrix metalloproteinases, ADMA = Asymmetric dimethylarginine, IL-6 = Interleukin-6, 2D = Two-dimensional, 3D = Three-dimensional, MiRNA = MicroRNA, SWI = Susceptibility-weighted imaging, FFE = Fast field echo, SAH = Sub-arachnoid haemorrhage

plasma membrane and has been implicated in the pathophysiology of neuronal injuries. It has already

Figure 1: Biomarkers approaches in acute neurological diseases. TBI = Traumatic brain injury, SAH = Sub-arachnoid haemorrhage, AIS = Acute ischaemic stroke, ICH = Intracerebral haemorrhage, SE = Status epilepticus

been used to look for biomarkers in Alzheimer's disease and also being proposed as a possible treatment of neurological injuries.^[29] Plasma post-translational modifications are non-DNA-coded modifications to the structure of proteins that generate novel and unique parts of a protein is another interesting aspect, for example, phosphorylated tau in Alzheimer's disease.^[30] Last, neurosteroids are endogenous brain molecules, and many demonstrate pleiotropic actions that are potentially relevant to TBI and its therapeutics.^[31] Although much work is on-going, these promising discoveries could yield transformative approaches in the diagnosis of acute neurological diseases.

WHAT DO WE KNOW SO FAR REGARDING SPECIFIC BIOMARKERS?

S-100B

First described in 1965,^[32] it is found in the cytosol of central nervous system (CNS) glial cells predominantly the astrocytes, but also extracranially, such as chondrocytes, melanocytes and adipocytes.^[33] It is the first identified member of the S-100 protein multigenic family and participates in an extra- and intra-cellular regulation of a cellular calcium metabolism.^[34,35] It can be detected in blood and CSF.

A few studies showed that S-100B is elevated in SAH compared to healthy subjects and is associated with vasospasm and poor outcome.^[36-38] External ventricular drain in SAH is associated with decreased blood S-100B, which may confound the usability of it.^[38]

A major limitation of S-100B is the acceptability range. At present, none of the reviewed assays established an acceptable range.^[39] The other drawback is its short half-life of 2 h,^[40] therefore, making it only relatively useful in the most severe form of brain injuries.^[41]

Neuron-specific enolase

NSE is a glycolytic enzyme found predominantly in the neuronal cytoplasm. First described in the 1960s,^[32] and has shown sensitivity or specificity in TBI, stroke and cardiac arrest and does not appear to exhibit age-dependent liabilities as seen in S-100B.^[3] Studies found that patients with NSE >28–97 µg/L post-cardiac arrest had poor outcome.^[6,42-45] Another prospective cohort study on 61 patients treated with therapeutic hypothermia after cardiac arrest showed that NSE and electroencephalogram findings were strongly correlated and while five survivors (3 with good outcome) had NSE levels 33 µg/L,^[46] which raised caution on the validity of NSE in the era of therapeutic hypothermia. However, a recently published randomised study of 686 patients to targeted temperature management at either 33°C or 36°C concluded that serial high NSE values have a high predictive value of poor outcome in cardiac arrest patients.^[47] An interesting study also found that in pre-eclampsia, the levels of NSE remained high throughout pregnancy,^[48] therefore raising the possibility of using it to monitor such a condition. As a result of some of these studies, a weak recommendation on the use of serum NSE in conjunction with clinical data for neurologic prognostication was given.^[6] Last, NSE is not a biomarker to be used alone as it has also been implicated as a marker for neuroendocrine bladder of tumour, small cell lung cancer and neuroblastomas.^[49]

Glial fibrillary acidic protein

GFAP is an intermediate filament protein that is only found in the glial cells of the CNS. First described in

1971,^[50] the usefulness of it as biomarkers has been reported in several neurological injuries. It has been shown to have excellent specificity and moderate sensitivity for TBI, while also having good specificity for computed tomography (CT)-confirmed brain injury^[51] and shown to have higher levels in patients with mass lesions compared with diffuse injury.^[52] These raise the redundancy of this biomarkers, as CT scans are readily available in most hospitals. In a recent prospective cohort study of 67 patients, serum GFAP levels on admission and during the first 5 days of injury were increased in patients with severe TBI and were predictive of neurological outcome at 6 months.^[53] However, they do not add predictive power to commonly used prognostic variables in a TBI population of varying severities.^[54] Overall, GFAP has the potential to be a useful biomarker, but more studies need to be done.

Matrix metalloproteinases

Matrix metalloproteinases (MMP) belong to a large family of proteolytic enzymes that degrades basement membrane components such as collagen IV, laminin and fibronectin, which are the major constituents of the blood-brain barrier.^[55] The presence of some of it such as MMP-2 and MMP-9 has been implicated as a negative prognostic factor in stroke.^[56,57] It is also implicated in other pathogenic mechanisms such as post-tissue plasminogen activator (tPA) haemorrhage in stroke, ICH, SAH and TBI. Higher levels are associated with thrombolysis failure.^[58-61] MMP-9 has also been correlated with haemorrhage transformation^[62] and malignant cerebral oedema^[63,64] in AIS. Plasma levels of cellular fibronectin 3.6 µg/mL and of MMP-9 140 ng/mL have been associated with parenchymal hematoma after treatment with rt-PA in patients with AIS in a multicentre confirmatory study.^[65,66] Therefore, in the last two decades, it is one of the most vigorously studied biomarkers for risk stratification and even as a possible neuroprotection agent through the inhibition of it.^[67] Despite all these, MMPs have not shown sufficient sensitivity and specificity for use in the clinical setting. They also have shown weak discriminative ability between AIS and stroke mimics in comparison to baseline clinical parameters, with non-significant improvement in receiver operating characteristic area under the curve.^[68]

Ubiquitin C-terminal hydrolase L1

UCH-L1 is a protein that is involved in the addition or removal of ubiquitin from proteins that are destined for metabolism. First detected in 1980 as protein gene product 9.5.^[69] It presents in human brain at concentrations at least 50-fold greater than in other organs.^[70] However, its exclusivity was questioned when it was also found in non-neuroendocrine carcinomas, such as those of the breast, kidney, prostate, pancreas, lung and colon.^[71] Increased CSF and blood concentrations have

been associated with neuron destruction and increased blood–brain barrier permeability.^[72] It has also been reported in neurodegenerative diseases.^[73] Remarkably, this neuronal protein can be readily detected in CSF and blood very early after brain injury,^[74] status epilepticus^[75] and carbon monoxide poisoning^[76] which allow it to be used as valuable time-window biomarkers for potential neuroprotective strategies. However, in a recently published prospective study of 324 patients with TBI, despite significantly associated with outcome, it does not add predictive power to commonly used prognostic variables.^[54]

Others

Other biomarkers may also have the potential to either being used alone or being used along with others [Table 1]. For example, fibrinogen level reduction by more than 200 mg/dL after tPA shows a higher risk of symptomatic ICH and, therefore being suggested to monitor post-rt-PA haemorrhage.^[77] However, a multicentre retrospective study published recently concluded that it did not significantly reduce the likelihood of in-hospital mortality or hematoma expansion.^[78] Thereby, raising the futility of fibrinogen testing post-tPA. Inflammatory proteins such as interleukin-6 have also been shown to be elevated in CSF in TBI.^[79] However, given its low specificity, it has not been applied clinically. Serum and CSF increased tumour necrosis factor- α levels have been described in patients following a severe TBI,^[80] however, these increased levels did not equate to an increased mortality rate,^[81] along with its low specificity and sensitivity, limiting its use clinically. Future potential markers include spectrin breakdown proteins, microtubule-associated protein, soluble urokinase plasminogen activator receptor and many other more.

LIMITATIONS

Recent reviews summarised the potentials of some of the most common biomarkers discussed^[6,49] and concluded that no single biomarker alone has been proven to have substantial clinical use and more research needs to be undertaken.

Despite progress over the last 25 years in translational science, much of the work has followed a typical pattern. After an initial period of optimism, there is, then, a realisation that although an individual biomarker may be of some use, there are often multiple confounding factors.^[41]

Unlike most organs in the body, the brain itself consists of multiple sub-structures that may share the same biomarkers that serve very different functions. Many of these sub-structures are small and when damaged, it may have a huge morbidity impact. For instance, a small

injury to the pons and frontal lobe may lead to equal elevation of biomarkers, but the consequences are widely different. Hence, how do we interpret it meaningfully?

The rising costs of medical care also raise the concern of the practicality of these tests. Even when proven to be clinically useful, the cost of it has to be compared with more traditional approaches. Last, there is also a lack of consistency across research methods on these topics, and these discrepancies hamper the progress. Qualification of a predictive biomarker signature must be based on randomised prospective clinical trials to demonstrate predictability in the biomarker-defined responders.^[82]

THE FUTURE

New developments in the field of proteomics, lipidomic, genetic markers, exosomes and miRNAs hold great promise in the discovery of newer makers, with trend in the development of personalised medicine, these approaches may even be used as treatment. Therefore, despite many inconclusive and even conflicting studies, the future of biomarkers look very bright.

CONCLUSIONS

Despite multiple studies and the enthusiasm towards the development of it, no single biomarker has proven to be applicable clinically. In the foreseeable future, it may be used as an adjunct, supplementing a good neurological examination and neuroimaging to help in the diagnosis and prognostication; this incorporation will be an important tool for neurocritical care specialist. Looking forward, the challenge will be to address the validity of it in different spectrum of brain injuries and to demonstrate that effective treatment can be as a result of it. To advance this field, multinational and multi-institution collaborations will be needed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Biomarkers Definitions Working Group. Biomarkers and surrogate endpoints: Preferred definitions and conceptual framework. *Clin Pharmacol Ther* 2001;69:89-95.
2. Rudman D, Fleischer A, Kutner MH. Concentration of 3', 5' cyclic adenosine monophosphate in ventricular cerebrospinal fluid of patients with prolonged coma after head trauma or intracranial hemorrhage. *N Engl J Med* 1976;295:635-8.
3. Kochanek PM, Berger RP, Bayir H, Wagner AK, Jenkins LW, Clark RS. Biomarkers of primary and evolving damage in traumatic and ischemic brain injury: Diagnosis, prognosis, probing mechanisms, and therapeutic decision making. *Curr Opin Crit Care* 2008;14:135-41.

4. Vaagenes P, Safar P, Diven W, Moossy J, Rao G, Cantadore R, *et al*. Brain enzyme levels in CSF after cardiac arrest and resuscitation in dogs: Markers of damage and predictors of outcome. *J Cereb Blood Flow Metab* 1988;8:262-75.
5. Vaagenes P, Mullie A, Fodstad DT, Abramson N, Safar P. The use of cytosolic enzyme increase in cerebrospinal fluid of patients resuscitated after cardiac arrest. *Brain Resuscitation Clinical Trial I Study Group. Am J Emerg Med* 1994;12:621-4.
6. Chou SH, Robertson CS; Participants in the International Multi-disciplinary Consensus Conference on the Multimodality Monitoring. Monitoring biomarkers of cellular injury and death in acute brain injury. *Neurocrit Care* 2014;21 Suppl 2:S187-214.
7. Kulbe JR, Geddes JW. Current status of fluid biomarkers in mild traumatic brain injury. *Exp Neurol* 2015. pii: S0014-488600151-X.
8. Jenkins LW, Peters GW, Dixon CE, Zhang X, Clark RS, Skinner JC, *et al*. Conventional and functional proteomics using large format two-dimensional gel electrophoresis 24 hours after controlled cortical impact in postnatal day 17 rats. *J Neurotrauma* 2002;19:715-40.
9. Gao W, Lu C, Kochanek PM, Berger RP. Serum amyloid A is increased in children with abusive head trauma: A gel-based proteomic analysis. *Pediatr Res* 2014;76:280-6.
10. Bazan NG, Marcheselli VL, Cole-Edwards K. Brain response to injury and neurodegeneration: Endogenous neuroprotective signaling. *Ann N Y Acad Sci* 2005;1053:137-47.
11. Han X, Gross RW. Global analyses of cellular lipidomes directly from crude extracts of biological samples by ESI mass spectrometry: A bridge to lipidomics. *J Lipid Res* 2003;44:1071-9.
12. Bayir H, Tyurin VA, Tyurina YY, Viner R, Ritov V, Amoscato AA, *et al*. Selective early cardiolipin peroxidation after traumatic brain injury: An oxidative lipidomics analysis. *Ann Neurol* 2007;62:154-69.
13. Samhan-Arias AK, Ji J, Demidova OM, Sparvero LJ, Feng W, Tyurin V, *et al*. Oxidized phospholipids as biomarkers of tissue and cell damage with a focus on cardiolipin. *Biochim Biophys Acta* 2012;1818:2413-23.
14. Wood PL. Lipidomics of Alzheimer's disease: Current status. *Alzheimers Res Ther* 2012;4:5.
15. Merritt VC, Arnett PA. Apolipoprotein E (APOE) $\epsilon 4$ allele is associated with increased symptom reporting following sports concussion. *J Int Neuropsychol Soc* 2016;22:89-94.
16. Li L, Bao Y, He S, Wang G, Guan Y, Ma D, *et al*. The association between apolipoprotein E and functional outcome after traumatic brain injury: A Meta-Analysis. *Medicine (Baltimore)* 2015;94:e2028.
17. Lanterna LA, Ruigrok Y, Alexander S, Tang J, Biroli F, Dunn LT, *et al*. Meta-analysis of APOE genotype and subarachnoid hemorrhage: Clinical outcome and delayed ischemia. *Neurology* 2007;69:766-75.
18. Biffi A, Sonni A, Anderson CD, Kissela B, Jagiella JM, Schmidt H, *et al*. Variants at APOE influence risk of deep and lobar intracerebral hemorrhage. *Ann Neurol* 2010;68:934-43.
19. Raffeld MR, Biffi A, Battey TW, Ayres AM, Viswanathan A, Greenberg SM, *et al*. APOE $\epsilon 4$ and lipid levels affect risk of recurrent nonlobar intracerebral hemorrhage. *Neurology* 2015;85:349-56.
20. Holtzman DM, Goate A, Kelly J, Sperling R. Mapping the road forward in Alzheimer's disease. *Sci Transl Med* 2011;3:114ps48.
21. Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993;75:843-54.
22. Ambros V. The functions of animal microRNAs. *Nature* 2004;431:350-5.
23. Cardoso AL, Guedes JR, de Lima MC. Role of microRNAs in the regulation of innate immune cells under neuroinflammatory conditions. *Curr Opin Pharmacol* 2015;26:1-9.
24. Pareek S, Roy S, Kumari B, Jain P, Banerjee A, Vrati S. MiR-155 induction in microglial cells suppresses Japanese encephalitis virus replication and negatively modulates innate immune responses. *J Neuroinflammation* 2014;11:97.
25. Redell JB, Moore AN, Ward NH 3rd, Hergenroeder GW, Dash PK. Human traumatic brain injury alters plasma microRNA levels. *J Neurotrauma* 2010;27:2147-56.
26. Yu B, Zhou S, Yi S, Gu X. The regulatory roles of non-coding RNAs in nerve injury and regeneration. *Prog Neurobiol* 2015;134:122-39.
27. Jickling GC, Ander BP, Zhan X, Noblett D, Stamova B, Liu D. microRNA expression in peripheral blood cells following acute ischemic stroke and their predicted gene targets. *PLoS One* 2014;9:e99283.
28. Elshal MF, McCoy JP. Multiplex bead array assays: Performance evaluation and comparison of sensitivity to ELISA. *Methods* 2006;38:317-23.
29. Properzi F, Logozzi M, Fais S. Exosomes: The future of biomarkers in medicine. *Biomark Med* 2013;7:769-78.
30. Henriksen K, O'Bryant SE, Hampel H, Trojanowski JQ, Montine TJ, Jeromin A, *et al*. The future of blood-based biomarkers for Alzheimer's disease. *Alzheimers Dement* 2014;10:115-31.
31. Marx CE, Naylor JC, Kilts JD, *et al*. Neurosteroids and traumatic brain injury: Translating biomarkers to therapeutics; overview and pilot investigations in Iraq and Afghanistan Era Veterans. In: Laskowitz D, Grant G, editors. *Translational Research in Traumatic Brain Injury*. Boca Raton (FL): CRC Press/Taylor and Francis Group; 2015.
32. Moore BW, McGregor D. Chromatographic and electrophoretic fractionation of soluble proteins of brain and liver. *J Biol Chem* 1965;240:1647-53.
33. Zhu L, Okano S, Takahara M, Chiba T, Tu Y, Oda Y, *et al*. Expression of S100 protein family members in normal skin and sweat gland tumors. *J Dermatol Sci* 2013;70:211-9.
34. Schäfer BW, Heizmann CW. The S100 family of EF-hand calcium-binding proteins: Functions and pathology. *Trends Biochem Sci* 1996;21:134-40.
35. Hajduková L, Sobek O, Prchalová D, Bílková Z, Koudelková M, Lukášková J, *et al*. Biomarkers of brain damage: S100B and NSE concentrations in cerebrospinal fluid-a normative study. *Biomed Res Int* 2015;2015:379071.
36. Wiesmann M, Missler U, Hagenström H, Gottmann D. S-100 protein plasma levels after aneurysmal subarachnoid haemorrhage. *Acta Neurochir (Wien)* 1997;139:1155-60.
37. Oertel M, Schumacher U, McArthur DL, Kästner S, Böker DK. S-100B and NSE: Markers of initial impact of subarachnoid haemorrhage and their relation to vasospasm and outcome. *J Clin Neurosci* 2006;13:834-40.
38. Stranjalis G, Korfiatis S, Psachoulia C, Kouyialis A, Sakas DE, Mendelow AD. The prognostic value of serum S-100B protein in spontaneous subarachnoid haemorrhage. *Acta Neurochir (Wien)* 2007;149:231-7.
39. Strathmann FG, Schulte S, Goerl K, Petron DJ. Blood-based biomarkers for traumatic brain injury: Evaluation of research approaches, available methods and potential utility from the clinician and clinical laboratory perspectives. *Clin Biochem* 2014;47:876-88.
40. Žurek J, Fedora M. The usefulness of S100B, NSE, GFAP, NF-H, secretogogin and Hsp70 as a predictive biomarker of outcome in children with traumatic brain injury. *Acta Neurochir (Wien)* 2012;154:93-103.
41. Toman E, Harrison S, Belli T. Biomarkers in traumatic brain injury: A review. *J R Army Med Corps* 2015. pii: Jramc-2015-000517.

42. Cronberg T, Rundgren M, Westhall E, Englund E, Siemund R, Rosén I, *et al.* Neuron-specific enolase correlates with other prognostic markers after cardiac arrest. *Neurology* 2011;77:623-30.
43. Rundgren M, Karlsson T, Nielsen N, Cronberg T, Johnsson P, Friberg H. Neuron specific enolase and S-100B as predictors of outcome after cardiac arrest and induced hypothermia. *Resuscitation* 2009;80:784-9.
44. Daubin C, Quentin C, Allouche S, Etard O, Gaillard C, Seguin A, *et al.* Serum neuron-specific enolase as predictor of outcome in comatose cardiac-arrest survivors: A prospective cohort study. *BMC Cardiovasc Disord* 2011;11:48.
45. Zandbergen EG, Hijdra A, Koelman JH, Hart AA, Vos PE, Verbeek MM, *et al.* Prediction of poor outcome within the first 3 days of postanoxic coma. *Neurology* 2006;66:62-8.
46. Rossetti AO, Carrera E, Oddo M. Early EEG correlates of neuronal injury after brain anoxia. *Neurology* 2012;78:796-802.
47. Stammet P, Collignon O, Hassager C, Wise MP, Hovdenes J, Åneman A, *et al.* Neuron-Specific Enolase as a predictor of death or poor neurological outcome after out-of-hospital cardiac arrest and targeted temperature management at 33°C and 36°C. *J Am Coll Cardiol* 2015;65:2104-14.
48. Bergman L, Åkerud H. Plasma levels of the cerebral biomarker, neuron-specific enolase, are elevated during pregnancy in women developing preeclampsia. *Reprod Sci* 2015. pii: 1933719115604732.
49. Forde CT, Karri SK, Young AM, Ogilvy CS. Predictive markers in traumatic brain injury: Opportunities for a serum biosignature. *Br J Neurosurg* 2014;28:8-15.
50. Eng LF, Vanderhaeghen JJ, Bignami A, Gerstl B. An acidic protein isolated from fibrous astrocytes. *Brain Res* 1971;28:351-4.
51. Sandler SJ, Figaji AA, Adelson PD. Clinical applications of biomarkers in pediatric traumatic brain injury. *Childs Nerv Syst* 2010;26:205-13.
52. Mondello S, Papa L, Buki A, Bullock MR, Czeiter E, Tortella FC, *et al.* Neuronal and glial markers are differently associated with computed tomography findings and outcome in patients with severe traumatic brain injury: A case control study. *Crit Care* 2011;15:R156.
53. Lei J, Gao G, Feng J, Jin Y, Wang C, Mao Q, *et al.* Glial fibrillary acidic protein as a biomarker in severe traumatic brain injury patients: A prospective cohort study. *Crit Care* 2015;19:362.
54. Takala RS, Posti JP, Runtti H, Newcombe VF, Outtrim J, Katila AJ, *et al.* GFAP and UCH-L1 as outcome predictors in traumatic brain injury. *World Neurosurg* 2015. pii: S1878-875001484-9.
55. Horstmann S, Kalb P, Koziol J, Gardner H, Wagner S. Profiles of matrix metalloproteinases, their inhibitors, and laminin in stroke patients: Influence of different therapies. *Stroke* 2003;34:2165-70.
56. Montaner J, Alvarez-Sabín J, Molina C, Anglés A, Abilleira S, Arenillas J, *et al.* Matrix metalloproteinase expression after human cardioembolic stroke: Temporal profile and relation to neurological impairment. *Stroke* 2001;32:1759-66.
57. Clark AW, Krekoski CA, Bou SS, Chapman KR, Edwards DR. Increased gelatinase A (MMP-2) and gelatinase B (MMP-9) activities in human brain after focal ischemia. *Neurosci Lett* 1997;238:53-6.
58. Heo JH, Kim SH, Lee KY, Kim EH, Chu CK, Nam JM. Increase in plasma matrix metalloproteinase-9 in acute stroke patients with thrombolysis failure. *Stroke* 2003;34:e48-50.
59. Ning M, Furie KL, Koroshetz WJ, Lee H, Barron M, Lederer M, *et al.* Association between tPA therapy and raised early matrix metalloproteinase-9 in acute stroke. *Neurology* 2006;66:1550-5.
60. Castellanos M, Sobrino T, Millán M, *et al.* Plasma MMP-9 and fibronectin levels confirm their value as powerful predictors of hemorrhagic transformation after thrombolytic therapy in acute ischemic stroke: A multicenter prospective study. In: *Stroke*. Philadelphia, PA, USA: Lippincott Williams & Wilkins; 2007. p. 463.
61. Montaner J, Molina CA, Monasterio J, Abilleira S, Arenillas JF, Ribó M, *et al.* Matrix metalloproteinase-9 pretreatment level predicts intracranial hemorrhagic complications after thrombolysis in human stroke. *Circulation* 2003;107:598-603.
62. Jha R, Battey TW, Pham L, Lorenzano S, Furie KL, Sheth KN, *et al.* Fluid-attenuated inversion recovery hyperintensity correlates with matrix metalloproteinase-9 level and hemorrhagic transformation in acute ischemic stroke. *Stroke* 2014;45:1040-5.
63. Moldes O, Sobrino T, Millán M, Castellanos M, Pérez de la Ossa N, Leira R, *et al.* High serum levels of endothelin-1 predict severe cerebral edema in patients with acute ischemic stroke treated with t-PA. *Stroke* 2008;39:2006-10.
64. Serena J, Blanco M, Castellanos M, Silva Y, Vivancos J, Moro MA, *et al.* The prediction of malignant cerebral infarction by molecular brain barrier disruption markers. *Stroke* 2005;36:1921-6.
65. Castellanos M, Leira R, Serena J, Blanco M, Pedraza S, Castillo J, *et al.* Plasma cellular-fibronectin concentration predicts hemorrhagic transformation after thrombolytic therapy in acute ischemic stroke. *Stroke* 2004;35:1671-6.
66. Castellanos M, Sobrino T, Millán M, García M, Arenillas J, Nombela F, *et al.* Serum cellular fibronectin and matrix metalloproteinase-9 as screening biomarkers for the prediction of parenchymal hematoma after thrombolytic therapy in acute ischemic stroke: A multicenter confirmatory study. *Stroke* 2007;38:1855-9.
67. Chang JJ, Emanuel BA, Mack WJ, Tsvigoulis G, Alexandrov AV. Matrix metalloproteinase-9: Dual role and temporal profile in intracerebral hemorrhage. *J Stroke Cerebrovasc Dis* 2014;23:2498-505.
68. Raman K, Paré G. Of stroke and biomarkers: The elusive quest for a clinical biomarker panel. *Clin Biochem* 2013;46:705-6.
69. Jackson P, Thompson RJ. The demonstration of new human brain-specific proteins by high-resolution two-dimensional polyacrylamide gel electrophoresis. *J Neurol Sci* 1981;49:429-38.
70. Thompson RJ, Doran JF, Jackson P, Dhillon AP, Rode J. PGP 9.5 – A new marker for vertebrate neurons and neuroendocrine cells. *Brain Res* 1983;278:224-8.
71. Campbell LK, Thomas JR, Lamps LW, Smoller BR, Folpe AL. Protein gene product 9.5 (PGP 9.5) is not a specific marker of neural and nerve sheath tumors: An immunohistochemical study of 95 mesenchymal neoplasms. *Mod Pathol* 2003;16:963-9.
72. Blyth BJ, Farahvar A, He H, Nayak A, Yang C, Shaw G, *et al.* Elevated serum ubiquitin carboxy-terminal hydrolase L1 is associated with abnormal blood-brain barrier function after traumatic brain injury. *J Neurotrauma* 2011;28:2453-62.
73. Zhang M, Cai F, Zhang S, Zhang S, Song W. Overexpression of ubiquitin carboxyl-terminal hydrolase L1 (UCHL1) delays Alzheimer's progression *in vivo*. *Sci Rep* 2014;4:7298.
74. Brophy GM, Mondello S, Papa L, Robicsek SA, Gabrielli A, Tepas J 3rd, *et al.* Biokinetic analysis of ubiquitin C-terminal hydrolase-L1 (UCH-L1) in severe traumatic brain injury patient biofluids. *J Neurotrauma* 2011;28:861-70.
75. Mondello S, Palmio J, Streeter J, Hayes RL, Peltola J, Jeromin A. Ubiquitin carboxy-terminal hydrolase L1 (UCH-L1) is increased in cerebrospinal fluid and plasma of patients after epileptic seizure. *BMC Neurol* 2012;12:85.
76. Pang L, Wu Y, Dong N, Xu DH, Wang DW, Wang ZH, *et al.* Elevated serum ubiquitin C-terminal hydrolase-L1 levels in patients with carbon monoxide poisoning. *Clin Biochem* 2014;47:72-6.

77. Yaghi S, Eisenberger A, Willey JZ. Symptomatic intracerebral hemorrhage in acute ischemic stroke after thrombolysis with intravenous recombinant tissue plasminogen activator: A review of natural history and treatment. *JAMA Neurol* 2014;71:1181-5.
78. Yaghi S, Boehme AK, Dibu J, Leon Guerrero CR, Ali S, Martin-Schild S, *et al.* Treatment and outcome of thrombolysis-related hemorrhage: A Multicenter Retrospective Study. *JAMA Neurol* 2015;72:1451-7.
79. Kumar RG, Diamond ML, Boles JA, Berger RP, Tisherman SA, Kochanek PM, *et al.* Acute CSF interleukin-6 trajectories after TBI: Associations with neuroinflammation, polytrauma, and outcome. *Brain Behav Immun* 2015;45:253-62.
80. Ross SA, Halliday MI, Campbell GC, Byrnes DP, Rowlands BJ. The presence of tumour necrosis factor in CSF and plasma after severe head injury. *Br J Neurosurg* 1994;8:419-25.
81. Crespo AR, Da Rocha AB, Jotz GP, Schneider RF, Grivicich I, Pinheiro K, *et al.* Increased serum sFas and TNFalpha following isolated severe head injury in males. *Brain Inj* 2007;21:441-7.
82. Chen JJ, Lu TP, Chen YC, Lin WJ. Predictive biomarkers for treatment selection: Statistical considerations. *Biomark Med* 2015;9:1121-35.