

Management of Mild Brain Trauma in the Elderly: Literature Review

Abstract

Purpose: The world population is aging. As direct consequence, geriatric trauma is increasing both in absolute number and in the proportion of annual admissions causing a challenge for the health-care system worldwide. The aim of this review is to delineate the specific and practice rules for the management of mild brain trauma in the elderly. **Methods:** Systematic review of the last 15 years literature on mild traumatic brain injury (mTBI) in elderly patients. **Results:** A total of 68 articles met all eligibility criteria and were selected for the systematic review. We collected 29% high-quality studies and 71% low-quality studies. **Conclusion:** Clinical advices for a comprehensive management are provided. Current outcome data from mTBIs in the elderly show a condition that cannot be sustained in the future by families, society, and health-care systems. There is a strong need for more research on geriatric mild brain trauma addressed to prevent falls, to reduce the impact of polypharmacy, and to define specific management strategies.

Keywords: Brain trauma, elderly, head injury, head trauma, mild brain trauma

Introduction

The world population is aging. It is estimated that almost two billion people will be older than 60 years by 2050.^[1] In the USA, in 2030, one in five residents is expected to be 65 years or older with a projection to doubling the current number in 2050.^[2] A similar trend has been observed in Canada.^[3]

As direct consequence, geriatric trauma is increasing both in absolute number and in the proportion of annual admissions,^[4] causing a challenge for the health-care system worldwide.

Traumatic brain injury (TBI) represents a major cause of mortality and morbidity which involves globally more than 10 million individuals annually,^[5] with no substantial differences in the incidence between low- and high-income countries.^[6]

Since TBI used to be prevalent in young people, nowadays, we are witnessing in a shift in the epidemiology of TBI toward the elderly.^[5] This is the result of mainly two factors: first, the aging of population as seen above; second, the decrease of traffic accidents due to the improvement of traffic safety which allowed falls to become the

most important cause of TBI. Moreover, falls account for over 65% of TBI-related hospital admissions in the elderly.^[7]

Epidemiologic studies show that the majority of TBI in older patients is mild with an incidence ranging from 60% to 80%.^[3,8,9]

Mild TBI (mTBI) is defined by the Neurotraumatology Committee of the World Federation of Neurosurgical Societies as a head injury presenting with Initial Glasgow Coma Score (GCS) of 14 or 15.^[10] History of loss of consciousness (LoC) and posttraumatic amnesia may be or not be present. Consider 65 years as a cutoff for the elderly is widely accepted in the international literature, mostly after the report of Mosenthal *et al.* which demonstrates the age 65 and more to be a predictor of increased mortality after TBI.^[11]

This cohort of patients represents a challenge for clinicians because of the unique physiologic and clinical factors, including frailty, chronic health diseases and polypharmacy, which make them more prone to primary and secondary injury following TBI, regardless of severity.

Frailty is defined as a syndrome of decreased reserves (energy, physical ability,

**Federica Marrone^{1,2},
Luca Zavatto^{1,2},
Mario Allevi^{1,2},
Hambra Di
Vitantonio¹,
Daniele Francesco
Millimaggi¹,
Soheila Raysi
Dehcordi¹,
Alessandro Ricci¹,
Graziano Taddei¹**

¹Department of Neurosurgery,
San Salvatore Hospital,

²Department of Life,
Health and Environmental
Sciences (MESVA), University of
L'Aquila, L'Aquila, Italy

Address for correspondence:

Dr. Graziano Taddei,
Department of Neurosurgery,
San Salvatore Hospital,
L'Aquila, SS80 67100, Italy.
E-mail: gr.taddei@gmail.com

Access this article online

Website: www.asianjns.org

DOI: 10.4103/ajns.AJNS_205_20

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Marrone F, Zavatto L, Allevi M, Di Vitantonio H, Millimaggi DF, Dehcordi SR, *et al.* Management of mild brain trauma in the elderly: Literature review. Asian J Neurosurg 2020;15:809-20.

Submitted: 05-May-2020

Revised: 14-Jun-2020

Accepted: 03-Jul-2020

Published: 21-Dec-2020

cognition, and health) that gives rise to vulnerability in times of acute stress.^[12] It has been calculated that about 7% of the people older than 65 years and 20% older than 80 years have frailty syndrome.^[12]

Despite what was discussed above, the researcher in the last 15 years does not seem to have found interest in exploring a common address for the management of mTBI in the elderly.

In 2004, the World Health Organization instituted a task force on mTBI with the scope of finding methodological issues and research recommendations.^[13] After 10 years, the main research priorities included focus on the elderly and the development of validated clinical prediction rules for the prognosis.^[14]

In 2015, the American Association for the Surgery of Trauma performed a member survey disclosing that 65% of the participants thought that geriatric trauma patients are undertriaged and not transferred appropriately to the higher levels of care. Reported concerns were the lack of uniformly accepted protocols, primarily for mTBI.^[15]

More recently, a group from the University of Cambridge completed an extensive epidemiologic literature review with the conclusion that though increased admissions of elderly patients following TBI there was little or no evidence of corresponding increase of age of patients recruited for TBI in studies around the world.^[16]

Finally, in a large cohort study including about 6000 mTBI older patients, Cheng *et al.* concluded that mTBI is an independent significant risk factor of death in the elderly and emphasized the need for head injury prevention and research.^[17]

The aim of this review is to point out the state of the art of the literature focused on mTBI in the elderly with the scope of delineating specific and practice rules for the management and to define the lacking areas of research.

Methods

This literature review study was conducted in accordance to the recommendations outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement during the reporting of the current study's findings.^[18] A literature search using PubMed was performed for English human studies published in the last 15 years using the following keywords: "traumatic brain injury AND elderly" and "head injury AND elderly." The literature search was conducted on March 6, 2019.

Study selection

Studies were included if they met focus on mTBI in the elderly defined as people aged >65 years. The exclusion criteria were as follows: (1) animal studies and case reports, (2) studies focusing on severe and/or moderate TBI, (3) studies focusing on pediatric patients, (4) studies

that had reported data without distinction of age and trauma severity, (5) clinical trial still in progress, and (6) full-text reports unavailable on-line.

Each retrieved article was screened for eligibility by two independent reviewers (FM and GT) first at the title and abstract level, and then full-text level if potentially relevant. The references of the included articles had been also evaluated and articles that met our criteria had been comprised in the review [Figure 1].

Results

A total of 388 studies were identified, and after duplicates exclusion, 359 unique abstracts were screened. Abstracts were evaluated by the first author (FM), and 148 potentially relevant articles were selected for full-text review and subsequently obtained and assessed for eligibility by two authors (FM and GT). Additional 15 full-text articles were identified through references. Finally, a total of 68 articles met all eligibility criteria and were selected for the review.

Quality assessment was achieved categorizing each study by level of evidence (LoE), according to the framework proposed by Sauaia *et al.* for the surgical research.^[19] We found 10 prospective prognostic and diagnostic studies (LoE I); 4 prospective therapeutic/care management studies (LoE II); 3 prospective prognostic and epidemiological studies (LoE II); 31 retrospective prognostic studies and 2 prognostic case-control studies (LoE III); one therapeutic case-control study and one retrospective epidemiological study (LoE III); 2 epidemiological studies (LoE IV–V) and 3 prognostic studies (LoE V). We collected three systematic




Figure 1: Flow diagram for studies identification and selection process

reviews of predominantly level II studies with no negative criteria (LoE II); 4 systematic reviews with up to two negative criteria (LoE III); 4 reviews (LoE IV). Overall, we collected 29% high-quality studies (LoE I or II) and 71% low-quality studies (LoE III or V). Studies were published between 1997 and 2018.

Extensive reading of the selected articles permitted to identify the major concerns of the recent literature about mTBI in the elderly to be discussed on. A total of 8 focused on epidemiology. Risk factors were reported in six articles. Thirty-five addressed the management of mTBI whom a total of 25 focusing patients on anticoagulation/antiplatelet (AP) therapy, and one described seizure therapy. Outcome and mortality were analyzed by 19 papers. All studies and their main characteristics are presented in Table 1 [Appendix A].

Only 8 studies (11,76%) were designed to assess exclusively elder mTBI patients, with the rest presenting mixed cohorts for age and severity. Due to the high heterogeneity of the topics found, statistical analysis was not performed.

Discussion

The aging of the world population and the increase in life expectancy subject both emergency and neurosurgery departments to everyday challenges related to the management of TBI in the elderly. This is especially true for those with a mild trauma because it is estimated that 82% of older patients hospitalized for mTBI return to independent living after completing the medical, surgical, and physical treatments.^[11] Triage and appropriate treatment should be uniformed to obtain the better results in terms of outcome for patients and cost-effectiveness for health-care system. As most of the international literature is focused on younger patients or severe TBI, and nonspecific guidelines exist, older mTBI patients are often undertriaged and the entity of trauma may be underestimated.

The results of our literature review were grouped and evaluated according to the major concerns emerged.

Risk factors for traumatic brain injury and prevention

Mechanisms accounting for TBI in the elderly may be the same of younger people. Nonetheless, most of TBI in the elderly are attributable to falls either from standing or from heights,^[9,20-23] accounting for over 80% of TBI hospitalizations and in-hospital mortality.^[21]

Mild cognitive impairment or dementias, comorbidities, ambulation with assistance, and polypharmacy have been demonstrated as the risk factors for falls in older adults and subsequent admission for fall-related TBI (fr-TBI).^[9,20,21]

However, even in the absence of comorbidities, older age is associated with an increased risk of falls due to physiologic age-related deterioration of physical functions

in balance, vision and hearing, physical strength, gait, and dexterity.^[21,22]

Finally, older adults are particularly susceptible to suffering iatrogenic effects of medications because of the increased risk of polypharmacy, defined as the simultaneous use of multiple medications such as anticoagulants (ACs), psychotropics, and sedatives, defined as “fall risk-increasing drugs.”^[22,24] Polypharmacy represents a relevant problem as it has been shown that more than 20% of older Americans receive at least one inappropriate drug.^[25] Physicians must carefully weigh the risk-benefit ratio when making decisions to prescribe medication such as AC therapy to patients who are at a high risk of falling.^[26] The Prevention of Falls in the Elderly Trial demonstrated the fundamental role of falls prevention with appropriate interventions, including the following: exercise programs aimed at improving strength and balance; vision assessments to correct poor eyesight; educational programs to encourage safer behavior among patients and families; periodic medication review to eliminate unnecessary medications or mitigate iatrogenic effects.^[27]

However, there is some controversy in the literature regarding the optimal combination of these interventions, and the best prevention strategy for fr-TBI in the elderly has still to be elucidated.^[21,26]

Admission versus observation

The overall assessment of a geriatric mTBI patient requests an interdisciplinary collaboration to improve diagnosis and treatment of underlying comorbidities, management of medications, and to address the most appropriate care setting.^[4]

It is estimated that 65% of annually visited TBI elderly patients in the emergency departments require admission,^[28] thus establishing the guidelines to define the criteria for observation and discharge is mandatory.

Mild head trauma in older population is not as mild as in younger patients. Several studies showed a discrepancy of observed GCS between the elderly and young patients with similar anatomic severity of TBI, with geriatric TBI having better GCS than younger patients.^[29,30] This creates a false idea that the elderly could tolerate equivalent injuries better, leading to underestimate the entity of TBI. The mechanisms behind this phenomenon are unknown, but they are certainly related to comorbidities and differences in physiological response to injury.

Although LoC has a strong predictive value for intracranial hemorrhage (ICH),^[31] most older patients with mTBI who require surgical intervention do not have a history of LoC.^[32] These data suggest that GCS and neurology examination alone cannot reliably exclude significant intracranial pathology in the elderly with mTBI. Given the lack of validated clinical evidence for identifying mTBI older patients who can be managed without cranial

Table 1: Characteristics of selected studies

	Number of studies	Type of studies	Authors and year	LoE
Epidemiology	8	2 prospective prognostic	Jiang J <i>et al.</i> 2013	I
			de Guise <i>et al.</i> 2015	
		1 prospective epidemiological	Stocchetti N <i>et al.</i> 2014	II
		1 systematic review	Gaastra B <i>et al.</i> 2016	II
		1 epidemiological retrospective	Van den Brand CL <i>et al.</i> 2018	III
		1 epidemiological retrospective	Cetty L <i>et al.</i> 2017	IV
		1 review	Adams SD <i>et al.</i> 2015	IV
		1 epidemiological	Kristman VL <i>et al.</i> 2014	V
Risk factors	6	1 prospective prognostic	Fu WW <i>et al.</i> 2017	I
		2 retrospective prognostic	Evans D <i>et al.</i> 2015	III
			Teo DB <i>et al.</i> 2018	
		3 prognostic	Filler W <i>et al.</i> 2015	V
			Karibe H <i>et al.</i> 2017	
			Krishnamoorthy V <i>et al.</i> 2015	
Management, imaging and complication	35	3 prospective prognostic	Batey M <i>et al.</i> 2018	I
			Chenoweth JA <i>et al.</i> 2018	
			Nishijima DK <i>et al.</i> 2012	
		2 diagnostic	Clement CM <i>et al.</i> 2006	I
			Haydel M <i>et al.</i> 2000	
		2 prospective therapeutic/management	Ivascu FA <i>et al.</i> 2005	II
			Menditto VG <i>et al.</i> 2012	
		2 prospective prognostic	Kaen A <i>et al.</i> 2010*	II
			Riccardi A <i>et al.</i> 2017*	
		1 systematic review	Jagoda AS <i>et al.</i> 2008	II
		20 retro prognostic	Beynon C <i>et al.</i> 2015	III
			Campiglio L <i>et al.</i> 2017	
			Cohen D.B. <i>et al.</i> 2006	
			Collins CE <i>et al.</i> 2014	
			Franko J <i>et al.</i> 2006	
			Gangavati A.S. <i>et al.</i> 2009	
			Grandhi R. <i>et al.</i> 2015	
			Helmes E <i>et al.</i> 2011	
			Howard J.L. <i>et al.</i> 2009	
			Julien J <i>et al.</i> 2017	
			Kim DY <i>et al.</i> 2012	
			Lavoie A <i>et al.</i> 2004	
			Li J <i>et al.</i> 2001	
			Mann N <i>et al.</i> 2018*	
			Moore MM <i>et al.</i> 2012	
			Parra MW <i>et al.</i> 2013	
			Pieracci M <i>et al.</i> 2007	
			Riccardi A <i>et al.</i> 2013*	
			Scantling D <i>et al.</i> 2017	
			Swap C <i>et al.</i> 2016*	
		2 case-control prognostic	Mahler B <i>et al.</i> 2015	III
			Tseng JH <i>et al.</i> 2014*	
		2 systematic reviews	Callaway DW <i>et al.</i> 2007	III
			Miller J <i>et al.</i> 2015	
		1 review	Haydel M <i>et al.</i> 2012	IV
Outcome and rehabilitation	19	2 prospective prognostic	Brousseau AA <i>et al.</i> 2016*	I
			Mosenthal AC <i>et al.</i> 2004	
		1 prospective therapeutic	Pedersen A <i>et al.</i> 2015	II

Contd...

Table 1: Contd...

Number of studies	Type of studies	Authors and year	LoE
	1 retrospective therapeutic	Susman M <i>et al.</i> 2002	II
	1 systematic review	Werman HA <i>et al.</i> 2011	II
	9 retrospective prognostic	Bouras T <i>et al.</i> 2007	III
		Cheng PL <i>et al.</i> 2014*	
		Depreitere B <i>et al.</i> 2012	
		Flaada JT <i>et al.</i> 2007	
		Haring RS <i>et al.</i> 2015	
		Herou E <i>et al.</i> 2015	
		LeBlanc J <i>et al.</i> 2006	
		Salotto K <i>et al.</i> 2014	
		Shimoda K <i>et al.</i> 2014	
	1 case-control therapeutic	Stapert S <i>et al.</i> 2006	III
	2 systematic reviews	Levin HS <i>et al.</i> 1997	III
		Menzel JC <i>et al.</i> 2008	
	2 reviews	Cruise CM <i>et al.</i> 2006	IV
		Flanagan SR <i>et al.</i> 2005	

*Exclusively elder mTBI patients. Complete references are available as supplemental material (Appendix A). LoE – Level of evidence; mTBI – Mild traumatic brain injury

computed tomography (CT), radiological evaluation is always needed.^[33-35]

Clinical observation is also often involved because it has been showed that 80% of patients in need of neurosurgical intervention can be identified by worsening of symptoms during the first 6 h.^[36]

Admission criteria to a trauma center are strongly correlated with the presence of co-injuries, clotting disorders, and preexistent diseases. The role of anticoagulation/AP therapy is discussed further in this section. Certainly, elderly patients on anticoagulation and documented intracranial pathology should be admitted to the neurosurgical department.^[28] In case of normal CT and elevated international normalized ratio (INR), mTBI patients should be observed for 12–24 h,^[37] and those with normal INR and CT, and no other associated injury, may be discharged and followed-up at home by a care provider.^[28]

Several authors suggested that most of observed patients can be safely discharged without a second CT scan, in the absence of clinical impairment, due to the low rates of delayed ICH occurring after 24 h from trauma.^[38-43]

Finally, coinjuries and chronic diseases are considered the positive predictors of worse outcome and mortality when associated to mTBI in the elderly, claiming admission to a trauma center.^[1]

A comprehensive flow chart for the management is provided in Figure 2.

The role of anticoagulation/antiplatelet therapy

Anticoagulants (ACs), APs, and the more recent direct oral anticoagulants (DOACs) are frequently used among the elderly. This trend, together with the high risk of falling,

determines a difficult dilemma for clinicians who have to balance the risk of ICH with thromboembolic complications. Therefore, benefits may not always outweigh the risks. Even though an ICH may also occur in patients not taking any ACs or APs medications,^[44] it is well known that long-term oral anticoagulation is an independent risk factor for both immediate and delayed ICH, and it is significantly associated with worse outcome and higher mortality.^[45,46] Risk's classification for ICH requiring surgery according with GCS score and clinical manifestations is resumed in Table 2.^[44] The management of these older mTBI patients is still largely debated in the literature and a common strategy has not yet been achieved. ACs drugs like warfarin are associated with the higher increased risk for ICH in the setting of TBI^[47] and with elevated chances of ICH progression in the follow-up period.^[48] Conversely, DOACs have a considerably lower rate of development of ICH compared to warfarin^[49,50] and lesser chances of ICH progression.^[48] These data support the safer profile of DOACs than ACs, suggesting that they should be used as first choice in older patients with high risk of falling. However, warfarin and DOACs users have similar mortality rates,^[48] reflecting the fact that every effort to prevent falls should be accomplished in patients who need ACs or DOACs therapy. APs use also confer a slight increase rate of ICH,^[47,51] particularly clopidogrel more than aspirin,^[41] but they are not associated with higher mortality if not used in combination.^[48] Part of the management of mTBI elderly patients on ACs or APs therapy involves the need of discontinuation and/or reversal of these drugs. In case of ICH at the initial CT scan, prompt reversal of warfarin with fresh frozen plasma demonstrated to be effective in decreasing ICH progression and mortality.^[52] Furthermore, APs and DOACs should be discontinued in this setting, but potential drawbacks are the inability to rapidly reverse DOACs, as an antidote is not always available,




Figure 2: Flowchart for management of mild traumatic brain injury in elderly patients. Co-injuries and anticoagulant therapy must be combined (GCS: Glasgow Coma Score; CT: Computed tomography; LoE: Level of evidence)

Table 2: Stratification of risk of hematoma requiring neurosurgical intervention

Categories	Characteristic	Risk of neurosurgical intervention (%)
Low risk	GCS 15	<0.1
	No LOC	
	No amnesia	
	No vomiting	
	No HA	
Medium risk	GCS 15 with one or more of the following	1-3
	LOC	
	Amnesia	
	Vomiting	
High risk	HA	10
	GCS 14-15 or neurologic deficits or	
	Drug induced coagulopathies	
	Alcoholism	
	Drug abuse	
	Epilepsy	
	Previous neurologic treatment	
	Disabled elderly patients	

LOC – Loss of consciousness; GCS – Glasgow Coma Score; HA – Headache

and that platelet transfusion is not associated with benefit in mortality and outcome in case of aspirin or clopidogrel usage.^[53] Patients on ACs or APs therapy with initial normal CT scan have a cumulative incidence of delayed ICH very low (<1%), and there is no evidence that their therapy must be reversed or discontinued.^[41] Nevertheless, supratherapeutic INR levels should be taken back to therapeutic range in all mTBI patients. When neurosurgical intervention for ICH is needed, ACs or DOACs reversal may facilitate clot evacuation and limit recurrences. A flowchart for a proposed management based on our review is provided

in Figure 3. Despite this extensive literature search, no clear indications are emerged on when to restart ACs or APs medications if discontinued after a mTBI.

Seizure risk and prophylaxis

TBI is known to contribute to 4%–9% of all epilepsy cases, particularly mTBI is associated to a double risk for seizures in the general population.^[54] Related TBI epilepsy is defined as at least one seizure occurring 7 days or later after the trauma; isolated seizure occurring in the first 7 days is considered as acute symptom.^[54]




Figure 3: Flowchart for management of anticoagulant/antiplatelet therapy in mild traumatic brain injury elderly patients (ACs: Anticoagulants; APs: Antiplatelets; DOACs: Direct oral anticoagulants; INR: International normalized ratio; ICH: Intracranial hemorrhage; mTBI: Mild traumatic brain injury; CT: Computed tomography; LoE: Level of evidence)

New onset seizures in older patients are often focal and can differ for semiology from younger patients. Subtle and noncharacteristic symptoms lead to delayed diagnosis and include syncope, memory problems, confusional episodes, or mental status changes.^[55] Only one report matches our inclusion criteria. In this large prospective case-control study, Mahler *et al.* assessed a cumulative relative risk of 2.5 in mTBI patients older than 65 years, which increased six times with a brain contusion. Conversely, an isolated skull fracture was not associated with an increased risk.^[54] Based on these few data, seizure prophylaxis cannot be recommended in all cases, but physicians should be aware of the subtle and noncharacteristic symptoms to promptly confirm the diagnosis with an electroencephalogram and to start therapy if needed.

Outcome

Long-term impact of mTBI on the patient, their family and society cannot be underestimated. It is mainly associated with loss of independence and cognitive decline secondary to the trauma. The high impact for the health-care systems is also reflected by the prolonged hospitalization and the needing for rehabilitative therapy. In the acute settings at the time of hospitalization, literature data showed that 45% of mTBI elderly patients could be discharged at home, 38% need physical or cognitive rehabilitation, and 8% need a long-term placement.^[56] Mortality is still high, reported as about 8%,^[46,56-58] and it has a strong and linear correlation with age. Moreover, 23% of discharged mTBI elderly patients have severe disability.^[11] It is possible that these discouraging reports may have led to an undertreatment for these patients in the last years. However, there are suggestions that outcome may be improved with more invasive monitoring and aggressive care.^[58,59] Furthermore,

a recent prospective study demonstrates that elderly TBI patients benefit from highly specialized rehabilitation, and it fails to prove inferiority regarding the younger age group.^[60] Furthermore, for patients who need neurosurgical intervention, evidence that craniotomy will negatively affect outcome is lacking. When operated, mTBI elderly patients show a more favorable outcome than who do not undergo surgery.^[56,61] Our review supports the idea that age should not be the sole factor for limiting care in mTBI patients. Physicians should be aware that older patients with mTBI deserve prompt and aggressive care and rehabilitation, as they can respond similarly to younger patients. However, the retrospective nature of this supporting data must be reinforced by more prospective studies.

Limitations

This study is mainly subject to the classical limitation of all reviews: the overall quality of the studies included. The majority of studies presented level II or less, and only 6 studies (8%) were LoE I. Moreover, there is a high heterogeneity in the literature focused on elderly TBI patients, and very few articles consider primarily mTBI. This made it necessary to carry out a research work oriented in extracting data that could involve some bias in patients' selection and outcome results. Finally, the obtained data were not suitable for meta-analysis to reach the evidence of a systematic review and the given recommendations are only based directly on the LoE of the studies selected.

Conclusion

Assuming that life expectancy is constantly increasing as well as the world population over the age of 65 years, reduction in mortality and morbidity should be the aim of managing patients with TBI in this cohort. Since severe

TBI still represents a challenge in both young and older people, current outcome data from mTBI in elderly show a condition that cannot be sustained in the future by families, society, and health-care systems.

There is a strong need for more research on geriatric mTBI addressed to prevent falls, to reduce the impact of polypharmacy, and to define specific management strategies based on the identification of patients who have chance of unfavorable outcome despite mild trauma.

Age alone should not limit care in these patients because they can have a good outcome if properly treated.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Werman HA, Erskine T, Caterino J, Riebe JF, Valasek T, Members of the Trauma Committee of the State of Ohio EMS Board. Development of statewide geriatric patients trauma triage criteria. *Prehosp Disaster Med* 2011;26:170-9.
- Bruns J Jr., Hauser WA. The epidemiology of traumatic brain injury: A review. *Epilepsia* 2003;44:2-10.
- de Guise E, LeBlanc J, Dagher J, Tinawi S, Lamoureux J, Marcoux J, *et al.* Traumatic brain injury in the elderly: A level 1 trauma centre study. *Brain INJ* 2015;29:558-64.
- Adams SD, Holcomb JB. Geriatric trauma. *Curr Opin Crit Care* 2015;21:520-526 doi:10.1097/MCC.0000000000000246.
- van den Brand CL, Karger LB, Nijman ST, Hunink MG, Patka P, Jellema K. Traumatic brain injury in the Netherlands, trends in emergency department visits, hospitalization and mortality between 1998 and 2012. *Eur J Emerg Med* 2018;25:355-61.
- Cetty L, Abdin E, Vaingankar JA, Jeyagurunathan A, Chua BY, Picco L, *et al.* Prevalence and correlates of traumatic brain injury (TBI) in older adults: Results from the Well being of the Singapore Elderly (WiSE) study. *Int Psychogeriatr* 2017;29:1899-907.
- Haring RS, Narang K, Canner JK, Asemota AO, George BP, Selvarajah S, *et al.* Traumatic brain injury in the elderly: Morbidity and mortality trends and risk factors. *J Surg Res* 2015;195:1-9.
- Jiang JY. Head trauma in China. *Injury* 2013;44:1453-57.
- Tagliaferri F, Compagnone C, Korsic M, Servadei F, Kraus J. A systematic review of brain injury epidemiology in Europe. *Acta Neurochir (Wien)* 2006;148:255-68; discussion 268.
- Servadei F, Teasdale G, Merry G, Societies NCotWfoN. Defining acute mild head injury in adults: A proposal based on prognostic factors, diagnosis, and management. *J Neurotrauma* 2001;18:657-64.
- Mosenthal AC, Livingston DH, Lavery RF, Knudson MM, Lee S, Morabito D, *et al.* The effect of age on functional outcome in mild traumatic brain injury: 6 month report of a prospective multicenter trial. *J Trauma* 2004;56:1042-8.
- Polidoro A, Dornbusch T, Vestri A, Di Bona S, Alessandri C. Frailty and disability in the elderly: A diagnostic dilemma. *Arch Gerontol Geriatr* 2011;52:e75-78.
- Carroll L, Cassidy JD, Holm L, Kraus J, Coronado V. Methodological issues and research recommendations for mild traumatic brain injury: The who collaborating centre task force on mild traumatic brain injury. *J Rehabil Med* 2004;36:113-25.
- Kristman VL, Borg J, Godbolt AK, Salmi LR, Cancelliere C, Carroll LJ, *et al.* Methodological issues and research recommendations for prognosis after mild traumatic brain injury: Results of the international collaboration on mild traumatic brain injury prognosis. *Arch Phys Med Rehabil* 2014;95:S265-77.
- Kozar RA, Arbabi S, Stein DM, Shackford SR, Barraco RD, Biffl WL, *et al.* Injury in the aged: Geriatric trauma care at the crossroads. *J Trauma Acute Care Surg* 2015;78:1197-209.
- Gaasstra B, Longworth A, Matta B, Snelson C, Whitehouse T, Murphy N, *et al.* The ageing population is neglected in research studies of traumatic brain injury. *Br J Neurosurg* 2016;30:221-6.
- Cheng PL, Lin HY, Lee YK, Hsu CY, Lee CC, Su YC. Higher mortality rates among the elderly with mild traumatic brain injury: A nationwide cohort study. *Scand J Trauma Resusc Emerg Med* 2014;22:7.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, *et al.* The PRISMA statement for reporting systematic reviews and meta analyses of studies that evaluate health care interventions: Explanation and elaboration. *J Clin Epidemiol* 2009;62:e1-34.
- Sauaia A, Moore EE, Crebs JL, Maier RV, Hoyt DB, Shackford SR. Evidence level of individual studies: A proposed framework for surgical research. *J Trauma Acute Care Surg* 2012;72:1484-90.
- Evans D, Pester J, Vera L, Jeanmonod D, Jeanmonod R. Elderly fall patients triaged to the trauma bay: Age, injury patterns, and mortality risk. *Am J Emerg Med* 2015;33:1635-8.
- Fu WW, Fu TS, Jing R, McFaul SR, Cusimano MD. Predictors of falls and mortality among elderly adults with traumatic brain injury: A nationwide, population based study. *PLoS One* 2017;12:e0175868.
- Karibe H, Hayashi T, Narisawa A, Kameyama M, Nakagawa A, Tominaga T. Clinical characteristics and outcome in elderly patients with traumatic brain injury: For establishment of management strategy. *Neurol Med Chir (Tokyo)* 2017;57:418-25.
- Ramanathan DM, McWilliams N, Schatz P, Hillary FG. Epidemiological shifts in elderly traumatic brain injury: 18 year trends in Pennsylvania. *J Neurotrauma* 2012;29:1371-8.
- Hajjar ER, Cafiero AC, Hanlon JT. Polypharmacy in elderly patients. *Am J Geriatr Pharmacother* 2007;5:345-51.
- Zhan C, Sangl J, Bierman AS, Miller MR, Friedman B, Wickizer SW, *et al.* Potentially inappropriate medication use in the community dwelling elderly: Findings from the 19.
- Krishnamoorthy V, Distelhorst JT, Vavilala MS, Thompson H. Traumatic brain injury in the elderly: Burden, risk factors, and prevention. *J Trauma Nurs* 2015;22:204-08.
- Close J, Ellis M, Hooper R, Glucksman E, Jackson S, Swift C. Prevention of falls in the elderly trial (PROFET): A randomised controlled trial. *Lancet (London, England)* 1999;353:93-7.
- Callaway DW, Wolfe R. Geriatric trauma. *Emerg Med Clin North Am* 2007;25:837-60.
- Rau CS, Wu SC, Chen YC, Chien PC, Hsieh HY, Kuo PJ, *et al.* Effect of age on glasgow coma scale in patients with moderate and severe traumatic brain injury: An approach with propensity score matched population. *Int J Environ Res Public Health* 2017;14:1378. doi:10.3390/ijerph1411137.
- Salottolo K, Levy AS, Slone DS, Mains CW, Bar Or D. The effect of age on Glasgow coma scale score in patients with traumatic brain injury. *JAMA Surg* 2014;149:727.
- Mack LR, Chan SB, Silva JC, Hogan TM. The use of head

- computed tomography in elderly patients sustaining minor head trauma. *J Emerg Med* 2003;24:157-62.
32. Gangavati AS, Kiely DK, Kulchyski LK, Wolfe RE, Mottley JL, Kelly SP, *et al.* Prevalence and characteristics of traumatic intracranial hemorrhage in elderly fallers presenting to the emergency department without focal findings: Intracranial hemorrhage in elderly fallers. *J Am Geriatr Soc* 2009;57:1470-4.
 33. Jagoda AS, Bazarian JJ, Bruns JJ, Cantrill SV, Gean AD, Howard PK, *et al.* Clinical policy: Neuroimaging and decisionmaking in adult mild traumatic brain injury in the acute setting. *Ann Emerg Med* 2008;52:714-48.
 34. Smits M, Dippel DW, de Haan GG, Dekker HM, Vos PE, Kool DR, *et al.* External validation of the Canadian CT head rule and the new Orleans criteria for CT scanning in patients with minor head injury. *JAMA* 2005;294:1519-25.
 35. Vos PE, Alekseenko Y, Battistin L, Ehler E, Gerstenbrand F, Muresanu DF, *et al.* Societies EFoN. Mild traumatic brain injury. *Eur J Neurol* 2012;19:191-8.
 36. Clement CM, Stiell IG, Schull MJ, Rowe BH, Brison R, Lee JS, *et al.* Clinical features of head injury patients presenting with a glasgow coma scale score of 15 and who require neurosurgical intervention. *Ann Emerg Med* 2006;48:245-51.
 37. Itshayek E, Rosenthal G, Fraifeld S, Perez Sanchez X, Cohen JE, Spektor S. Delayed posttraumatic acute subdural hematoma in elderly patients on anticoagulation. *Neurosurgery* 2006;58:A851-6.
 38. Campiglio L, Bianchi F, Cattalini C, Belvedere D, Rosci CE, Casellato CL, *et al.* Mild brain injury and anticoagulants: Less is enough. *Neurol Clin Pract* 2017;7:296-305.
 39. Kaen A, Jimenez Roldan L, Arrese I, Delgado MA, Lopez PG, Alday R, *et al.* The value of sequential computed tomography scanning in anticoagulated patients suffering from minor head injury. *J Trauma* 2010;68:895-8.
 40. Miller J, Lieberman L, Nahab B, Hurst G, Gardner Gray J, Lewandowski A. Delayed intracranial hemorrhage in the anticoagulated patient: A systematic review. *J Trauma Acute Care Surg* 2015;79:310-3.
 41. Nishijima DK, Offerman SR, Ballard DW, Vinson DR, Chettipally UK, Rauchwerger AS, *et al.* Immediate and delayed traumatic intracranial hemorrhage in patients with head trauma and preinjury warfarin or clopidogrel use. *Ann Emerg Med* 2012;59:460-8.e467.
 42. Scantling D, Fischer C, Gruner R, Teichman A, McCracken B, Eakins J. The role of delayed head CT in evaluation of elderly blunt head trauma victims taking antithrombotic therapy. *Eur J Trauma Emerg Surg* 2017;43:741-6.
 43. Swap C. Risk of delayed intracerebral hemorrhage in anticoagulated patients after minor head trauma: The role of repeat cranial computed tomography. *Perm J* 2016; 20:14-6. doi: 10.7812/TPP/15-095.
 44. Chenoweth JA, Gaona SD, Faul M, Holmes JF, Nishijima DK, Consortium ft SC. Incidence of delayed intracranial hemorrhage in older patients after blunt head trauma. *JAMA Surg* 2018;153:570.
 45. Pieracci FM, Eachempati SR, Shou J, Hydo LJ, Barie PS. Degree of anticoagulation, but not warfarin use itself, predicts adverse outcomes after traumatic brain injury in elderly trauma patients. *J Trauma* 2007;63:525-30.
 46. Ritchie PD, Cameron PA, Ugoni AM, Kaye AH. A study of the functional outcome and mortality in elderly patients with head injuries. *J Clin Neurosci* 2000;7:301-4.
 47. Collins CE, Witkowski ER, Flahive JM, Anderson FA, Santry HP. Effect of preinjury warfarin use on outcomes after head trauma in Medicare beneficiaries. *Am J Surg* 2014;208:544-9. e541.
 48. Scotti P, Séguin C, Lo BW, de Guise E, Troquet JM, Marcoux J. Antithrombotic agents and traumatic brain injury in the elderly population: Hemorrhage patterns and outcomes. *J Neurosurg* 2019; 5:1-10.
 49. Kobayashi L, Barmparas G, Bosarge P, Brown CV, Bukur M, Carrick MM, *et al.* Group AMPOSoTPoNOAS. Novel oral anticoagulants and trauma: The results of a prospective American Association for the Surgery of Trauma Multi Institutional Trial. *J Trauma Acute Care Surg* 2017;82:827-35.
 50. Riccardi A, Spinola B, Minuto P, Ghinatti M, Guidido G, Malerba M, *et al.* Intracranial complications after minor head injury (MHI) in patients taking vitamin K antagonists (VKA) or direct oral anticoagulants (DOACs). *Am J Emerg Med* 2017;35:1317-9.
 51. Julien J, Alsieiri G, Marcoux J, Hasen M, Correa JA, Feyz M, *et al.* Antithrombotic agents intake prior to injury does not affect outcome after a traumatic brain injury in hospitalized elderly patients. *J Clin Neurosci* 2017;38:122-5.
 52. Ivascu FA, Howells GA, Junn FS, Bair HA, Bendick PJ, Janczyk RJ. Rapid warfarin reversal in anticoagulated patients with traumatic intracranial hemorrhage reduces hemorrhage progression and mortality. *J Trauma* 2005;59:1131-7.
 53. Grandhi R, Harrison G, Voronovich Z, Bauer J, Chen SH, Nicholas D, *et al.* Preinjury warfarin, but not antiplatelet medications, increases mortality in elderly traumatic brain injury patients. *J Trauma Acute Care Surg* 2015;78:614-21.
 54. Mahler B, Carlsson S, Andersson T, Adelöw C, Ahlbom A, Tomson T. Unprovoked seizures after traumatic brain injury: A population based case control study. *Epilepsia* 2015;56:1438-44.
 55. Motika PV, Spencer DC. Treatment of epilepsy in the elderly. *Curr Neurol Neurosci Rep* 2016;16:96. doi:10.1007/s11910-016-0696-8.
 56. Lai Fung L, Wai Man L, Wong HT, Yuen WK, Leung GK. Outcome after operative intervention for traumatic brain injuries in the elderly. *Asian J Neurosurg* 2017;12:37.
 57. Flaada JT, Leibson CL, Mandrekar JN, Diehl N, Perkins PK, Brown AW, *et al.* Relative risk of mortality after traumatic brain injury: A population based study of the role of age and injury severity. *J Neurotrauma* 2007;24:435-45.
 58. Susman M, DiRusso SM, Sullivan T, Risucci D, Nealon P, Cuff S, *et al.* Traumatic brain injury in the elderly: Increased mortality and worse functional outcome at discharge despite lower injury severity. *J Trauma* 2002;53:219-23; discussion 223-4.
 59. Scalea TM, Simon HM, Duncan AO, Atweh NA, Sclafani SJ, Phillips TF, *et al.* Geriatric blunt multiple trauma: Improved survival with early invasive monitoring. *J Trauma* 1990;30:129 34; discussion 134-6.
 60. Pedersen AR, Severinsen K, Nielsen JF. The effect of age on rehabilitation outcome after traumatic brain injury assessed by the functional independence measure (FIM). *Neurorehabil Neural Repair* 2015;29:299-307.
 61. Shimoda K, Maeda T, Tado M, Yoshino A, Katayama Y, Bullock MR. Outcome and surgical management for geriatric traumatic brain injury: Analysis of 888 cases registered in the Japan Neurotrauma Data Bank. *World Neurosurg* 2014;82:1300-6.

Appendix A

Complete references for Table 1 (alphabetic order)

- Adams SD, Holcomb JB. Geriatric trauma. *Curr Opin Crit Care* 2015;21:520–526 doi:10.1097/MCC.0000000000000246.
- Batey M, Hecht J, Callahan C, Wahl W. Direct oral anticoagulants do not worsen traumatic brain injury after low-level falls in the elderly. *Surgery* 2018;164:814-9.
- Beynon C, Potzy A, Sakowitz OW, Unterberg AW. Rivaroxaban and intracranial haemorrhage after mild traumatic brain injury: A dangerous combination? *Clin Neurol Neurosurg* 2015;136:73-8.
- Bouras T, Stranjalis G, Korfiatis S, Andrianakis I, Pitaridis M, Sakas DE. Head injury mortality in a geriatric population: differentiating an “edge” age group with better potential for benefit than older poor-prognosis patients. *J Neurotrauma* 2007;24:1355-61.
- Brousseau AA, Emond M, Sirois MJ, Daoust R, Griffith LE, Lang E, *et al.* Comparison of functional outcomes in elderly who have sustained a minor trauma with or without head injury: A prospective multicenter cohort study. *CJEM* 2017;19:329-37.
- Callaway DW, Wolfe R. Geriatric trauma. *Emerg Med Clin North Am* 2007;25:837-60.
- Campiglio L, Bianchi F, Cattalini C, Belvedere D, Rosci CE, Casellato CL, *et al.* Mild brain injury and anticoagulants: Less is enough. *Neurol Clin Pract* 2017;7:296-305.
- Cetty L, Abdin E, Vaingankar JA, Jeyagurunathan A, Chua BY, Picco L, *et al.* Prevalence and correlates of traumatic brain injury (TBI) in older adults: Results from the Well-being of the Singapore Elderly (WiSE) study. *Int Psychogeriatr* 2017;29:1899-907.
- Cheng PL, Lin HY, Lee YK, Hsu CY, Lee CC, Su YC. Higher mortality rates among the elderly with mild traumatic brain injury: A nationwide cohort study. *Scand J Trauma Resusc Emerg Med* 2014;22:7.
- Chenoweth JA, Gaona SD, Faul M, Holmes JF, Nishijima DK; Consortium ftSCPR. Incidence of delayed intracranial hemorrhage in older patients after blunt head trauma. *JAMA Surg* 2018;153:570.
- Clement CM, Stiell IG, Schull MJ, Rowe BH, Brison R, Lee JS, *et al.* Clinical features of head injury patients presenting with a glasgow coma scale score of 15 and who require neurosurgical intervention. *Ann Emerg Med* 2006;48:245-51.
- Cohen DB, Rinker C, Wilberger JE. Traumatic brain injury in anticoagulated patients. *J Trauma* 2006;60:553-7.
- Collins CE, Witkowski ER, Flahive JM, Anderson FA, Santry HP. Effect of preinjury warfarin use on outcomes after head trauma in medicare beneficiaries. *Am J Surg* 2014;208:544-9.e541.
- Cruise CM, Sasson N, Lee MH. Rehabilitation outcomes in the older adult. *Clin Geriatr Med* 2006;22:257-67.
- de Guise E, LeBlanc J, Dagher J, Tinawi S, Lamoureux J, Marcoux J, *et al.* Traumatic brain injury in the elderly: A level 1 trauma centre study. *Brain Inj* 2015;29:558-64.
- Depreitere B, Meyfroidt G, Roosen G, Ceuppens J, Grandas FG. Traumatic brain injury in the elderly: A significant phenomenon. *Acta Neurochir Suppl* 2012;114:289-94.
- Evans D, Pester J, Vera L, Jeanmonod D, Jeanmonod R. Elderly fall patients triaged to the trauma bay: Age, injury patterns, and mortality risk. *Am J Emerg Med* 2015;33:1635-8.
- Filer W, Harris M. Falls and traumatic brain injury among older adults. *N C Med J* 2015;76:111-4.
- Flaada JT, Leibson CL, Mandrekar JN, Diehl N, Perkins PK, Brown AW, *et al.* Relative risk of mortality after traumatic brain injury: A population-based study of the role of age and injury severity. *J Neurotrauma* 2007;24:435-45.
- Flanagan SR, Hibbard MR, Gordon WA. The impact of age on traumatic brain injury. *Phys Med Rehabil Clin N Am* 2005;16:163-77.
- Franko J, Kish KJ, O'Connell BG, Subramanian S, Yuschak JV. Advanced age and preinjury warfarin anticoagulation increase the risk of mortality after head trauma. *J Trauma* 2006;61:107-10.
- Fu WW, Fu TS, Jing R, McFaull SR, Cusimano MD. Predictors of falls and mortality among elderly adults with traumatic brain injury: A nationwide, population-based study. *PLoS One* 2017;12:e0175868.
- Gaastra B, Longworth A, Matta B, Snelson C, Whitehouse T, Murphy N, *et al.* The ageing population is neglected in research studies of traumatic brain injury. *Br J Neurosurg* 2016;30:221-6.
- Gangavati AS, Kiely DK, Kulchyski LK, Wolfe RE, Mottley JL, *et al.* Prevalence and characteristics of traumatic intracranial hemorrhage in elderly fallers presenting to the emergency department without focal findings: Intracranial hemorrhage in elderly fallers. *J Am Geriatr Soc* 2009;57:1470-4.
- Grandhi R, Harrison G, Voronovich Z, Bauer J, Chen SH, Nicholas D, *et al.* Preinjury warfarin, but not antiplatelet medications, increases mortality in elderly traumatic brain injury patients: . *J Trauma Acute Care Surg* 2015;78:614-21.
- Haring RS, Narang K, Canner JK, Asemota AO, George BP, Selvarajah S, *et al.* Traumatic brain injury in the elderly: Morbidity and mortality trends and risk factors. *J Surg Res* 2015;195:1-9.
- Haydel M. Management of mild traumatic brain injury in the emergency department. *Emerg Med Pract* 2012;14:1-24.
- Haydel MJ, Preston CA, Mills TJ, Luber S, Blaudeau E, DeBlieux PM. Indications for computed tomography in patients with minor head injury. *N Engl J Med* 2000;343:100-5.
- Helmes E, Ostbye T, Steenhuis RE. Incremental contribution of reported previous head injury to the prediction of diagnosis and cognitive functioning in older adults. *Brain Inj* 2011;25:338-47.
- Herou E, Romner B, Tomasevic G. Acute traumatic brain injury: Mortality in the elderly. *World Neurosurg* 2015;83:996-1001.
- Howard JL 2nd, Cipolle MD, Horvat SA, Sabella VM, Reed JF 3rd, Fulda G, *et al.* Preinjury warfarin worsens outcome in elderly patients who fall from standing. *J Trauma* 2009;66:1518-22.
- Ivascu FA, Howells GA, Junn FS, Bair HA, Bendick PJ, Janczyk RJ. Rapid warfarin reversal in anticoagulated patients with traumatic intracranial hemorrhage reduces hemorrhage progression and mortality. *J Trauma* 2005;59:1131-7.

33. Jagoda AS, Bazarian JJ, Bruns JJ, Cantrill SV, Gean AD, Howard PK, *et al.* Clinical policy: Neuroimaging and decisionmaking in adult mild traumatic brain injury in the acute setting. *Ann Emerg Med* 2008;52:714-48.
34. Jiang JY. Head trauma in China. *Injury* 2013;44:1453-7.
35. Julien J, Alsideiri G, Marcoux J, Hasen M, Correa JA, Feyz M, *et al.* Antithrombotic agents intake prior to injury does not affect outcome after a traumatic brain injury in hospitalized elderly patients. *J Clin Neurosci* 2017;38:122-5.
36. Kaen A, Jimenez-Roldan L, Arrese I, Delgado MA, Lopez PG, Alday R, *et al.* The value of sequential computed tomography scanning in anticoagulated patients suffering from minor head injury. *J Trauma* 2010;68:895-8.
37. Karibe H, Hayashi T, Narisawa A, Kameyama M, Nakagawa A, Tominaga T. Clinical characteristics and outcome in elderly patients with traumatic brain injury: For establishment of management strategy. *Neurol Med Chir (Tokyo)* 2017;57:418-25.
38. Kim DY, Kobayashi L, Barmparas G, Fortlage D, Curry T, Coimbra R. Venous thromboembolism in the elderly: The result of comorbid conditions or a consequence of injury? *J Trauma Acute Care Surg* 2012;72:1286-91.
39. Krishnamoorthy V, Distelhorst JT, Vavilala MS, Thompson H. Traumatic brain injury in the elderly: Burden, risk factors, and prevention. *J Trauma Nurs* 2015;22:204-8.
40. Kristman VL, Borg J, Godbolt AK, Salmi LR, Cancelliere C, Carroll LJ, *et al.* Methodological issues and research recommendations for prognosis after mild traumatic brain injury: Results of the international collaboration on mild traumatic brain injury prognosis. *Arch Phys Med Rehabil* 2014;95:S265-77.
41. Lavoie A, Ratte S, Clas D, Demers J, Moore L, Martin M, *et al.* Preinjury warfarin use among elderly patients with closed head injuries in a trauma center. *J Trauma* 2004;56:802-7.
42. LeBlanc J, de Guise E, Gosselin N, Feyz M. Comparison of functional outcome following acute care in young, middle-aged and elderly patients with traumatic brain injury. *Brain Inj* 2006;20:779-90.
43. Levin HS, Goldstein FC, MacKenzie EJ. Depression as a secondary condition following mild and moderate traumatic brain injury. *Semin Clin Neuropsychiatry* 1997;2:207-15.
44. Li J, Brown J, Levine M. Mild head injury, anticoagulants, and risk of intracranial injury. *Lancet* 2001;357:771-2.
45. Mahler B, Carlsson S, Andersson T, Adelöw C, Ahlbom A, Tomson T. Unprovoked seizures after traumatic brain injury: A population-based case-control study. *Epilepsia* 2015;56:1438-44.
46. Mann N, Welch K, Martin A, Subichin M, Wietecha K, Birmingham LE, *et al.* Delayed intracranial hemorrhage in elderly anticoagulated patients sustaining a minor fall. *BMC Emerg Med* 2018;18:27.
47. Menditto VG, Lucci M, Polonara S, Pomponio G, Gabrielli A. Management of minor head injury in patients receiving oral anticoagulant therapy: A prospective study of a 24-hour observation protocol. *Ann Emerg Med* 2012;59:451-5.
48. Menzel JC. Depression in the elderly after traumatic brain injury: A systematic review. *Brain Inj* 2008;22:375-80.
49. Miller J, Lieberman L, Nahab B, Hurst G, Gardner-Gray J, Lewandowski A, *et al.* Delayed intracranial hemorrhage in the anticoagulated patient: A systematic review. *J Trauma Acute Care Surg* 2015;79:310-3.
50. Moore MM, Pasquale MD, Badellino M. Impact of age and anticoagulation: Need for neurosurgical intervention in trauma patients with mild traumatic brain injury. *J Trauma Acute Care Surg* 2012;73:126-30.
51. Mosenthal AC, Livingston DH, Lavery RF, Knudson MM, Lee S, Morabito D, *et al.* The effect of age on functional outcome in mild traumatic brain injury: 6-month report of a prospective multicenter trial. *J Trauma* 2004;56:1042-8.
52. Nishijima DK, Offerman SR, Ballard DW, Vinson DR, Chettipally UK, Rauchwerger AS, *et al.* Immediate and delayed traumatic intracranial hemorrhage in patients with head trauma and preinjury warfarin or clopidogrel use. *Ann Emerg Med* 2012;59:460-8.e467.
53. Parra MW, Zucker L, Johnson ES, Gullett D, Avila C, Wichner ZA, *et al.* Dabigatran bleed risk with closed head injuries: Are we prepared? *J Neurosurg* 2013;119:760-5.
54. Pedersen AR, Severinsen K, Nielsen JF. The effect of age on rehabilitation outcome after traumatic brain injury assessed by the functional independence measure (FIM). *Neurorehabil Neural Repair* 2015;29:299-307.
55. Pieracci FM, Eachempati SR, Shou J, Hydo LJ, Barie PS. Degree of anticoagulation, but not warfarin use itself, predicts adverse outcomes after traumatic brain injury in elderly trauma patients. *J Trauma* 2007;63:525-30.
56. Riccardi A, Frumento F, Guidido G, Spinola MB, Corti L, Minuto P, *et al.* Minor head injury in the elderly at very low risk: A retrospective study of 6 years in an Emergency Department (ED). *Am J Emerg Med* 2013;31:37-41.
57. Riccardi A, Spinola B, Minuto P, Ghinatti M, Guidido G, Malerba M, *et al.* Intracranial complications after minor head injury (MHI) in patients taking Vitamin K antagonists (VKA) or direct oral anticoagulants (DOACs). *Am J Emerg Med* 2017;35:1317-9.
58. Salottolo K, Levy AS, Slone DS, Mains CW, Bar-Or D. The effect of age on Glasgow coma scale score in patients with traumatic brain injury. *JAMA Surg* 2014;149:727.
59. Scantling D, Fischer C, Gruner R, Teichman A, McCracken B, Eakins J. The role of delayed head CT in evaluation of elderly blunt head trauma victims taking antithrombotic therapy. *Eur J Trauma Emerg Surg* 2017;43:741-6.
60. Shimoda K, Maeda T, Tado M, Yoshino A, Katayama Y, Bullock MR. Outcome and surgical management for geriatric traumatic brain injury: Analysis of 888 cases registered in the Japan Neurotrauma Data Bank. *World Neurosurg* 2014;82:1300-6.
61. Stapert S, Houx P, de Kruijk J, Ponds R, Jolles J. Neurocognitive fitness in the sub-acute stage after mild TBI: the effect of age. *Brain Inj* 2006;20:161-5.
62. Stocchetti N, Paterno R, Citerio G, Beretta L, Colombo A. Traumatic brain injury in an aging population. *J Neurotrauma* 2012;29:1119-25.
63. Susman M, DiRusso SM, Sullivan T, Risucci D, Nealon P, Cuff S, *et al.* Traumatic brain injury in the elderly: Increased mortality and worse functional outcome at discharge despite lower injury severity. *J Trauma* 2002;53:219-23.
64. Swap C. Risk of delayed intracerebral hemorrhage in anticoagulated patients after minor head trauma: The role of repeat cranial computed tomography. *Perm J* 2016;20:214-6. [doi:10.7812/TPP/15-095].

65. Teo DB, Wong HC, Yeo AW, Lai YW, Choo EL, Merchant RA. Characteristics of fall-related traumatic brain injury in older adults. *Int Med J* 2018;48:1048-55.
66. Tseng JH, Tseng MY, Liu AJ, Lin WH, Hu HY, Hsiao SH. Risk factors for chronic subdural hematoma after a minor head injury in the elderly: A population-based study. *Biomed Res Int* 2014. pii: 218646.
67. Van den Brand CL, Karger LB, Nijman ST, Hunink MG, Patka P, Jellema K. Traumatic brain injury in the Netherlands, trends in emergency department visits, hospitalization and mortality between 1998 and 2012. *Eur J Emerg Med* 2018;25:355-61.
68. Werman HA, Erskine T, Caterino J, Riebe JF, Valasek T, Board MotTCotSoOE. Development of statewide geriatric patients trauma triage criteria. *Prehosp Disaster Med* 2011;26:170-9.