

## Original Article

# Comparison of subdermal and perforator delay techniques on a rat flap model

Dincer Altinel, Burcu Biltekin<sup>1</sup>, Merdan Serin, Ugur Sahin<sup>2</sup>, Muzaffer Altindas<sup>3</sup>

Department of Plastic and Reconstructive Surgery, Istanbul Training and Research Hospital, <sup>1</sup>Department of Histology and Embryology, Istanbul University Cerrahpasa Medical School, <sup>2</sup>Department of Plastic and Reconstructive Surgery, Bahcelievler Medica Hospital, <sup>3</sup>Department of Plastic and Reconstructive Surgery, Istanbul Diabetes Hospital, Istanbul, Turkey

**Address for correspondence:** Dr. Merdan Serin, Department of Plastic Surgery, Istanbul Research and Teaching Hospital, Istanbul, Turkey.  
E-mail: merdanserin@gmail.com

## ABSTRACT

**Background:** In this study, we investigated the subdermal and perforator delay phenomena as a method to improve flap survival. **Materials and Methods:** In this experimental study, we used 24 rats in three groups. In the control group, the dorsal flaps were elevated and reinserted back to their place. In the experimental groups, we practiced the delay phenomena with two different techniques. In the first experimental group, cranial and lateral side incisions were performed; however, the flaps were not cut-off from the underlying fascia. In the second experimental group, we placed a silicon sheet under the planned flap to cut-off the circulation from the perforator vessels. Four weeks after the delay procedure, the flaps were raised completely and reinserted back to their place. **Results:** The average of necrotic area in the control group was 21.9% ( $\pm 7.70$ ). There was no necrosis in both experimental groups ( $P < 0.0001$ ). Histological examination revealed that collagen density in both of the experimental groups was increased in comparison to the control group, it has only been found a significant first experimental group ( $P = 0.0315$ ). We have not found any significant difference in lymphocyte density between the groups. Angiographic imaging has showed an increase in the vascular density in the flaps of the first experimental group. **Conclusion:** We believe that both of these delay techniques can be adapted to clinical applications and used safely to increase flap survival.

## KEY WORDS

Flap delay; rat; skin flap

## INTRODUCTION

Flap survival depends on many internal and external factors.<sup>[1-5]</sup> Understanding of flap physiology is crucial for the success in reconstructive surgery.

Flap microcirculation is composed of small vessels such as arterioles, capillaries and venules. These can be affected by many biochemical and neurological impulses. Flap circulation can be severed by the impairment of arterial

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**How to cite this article:** Altinel D, Biltekin B, Serin M, Sahin U, Altindas M. Comparison of subdermal and perforator delay techniques on a rat flap model. Indian J Plast Surg 2018;51:196-201.

Access this article online	
Quick Response Code:	Website: www.ijps.org
	DOI: 10.4103/ijps.IJPS_28_17

or venous circulation.<sup>[6]</sup> In this regard, venous flaps have certain advantages to traditional arterial flaps among which are easier design and less donor-site morbidity.<sup>[7]</sup> Many factors have been shown to have effects on flap circulation such as topical applications,<sup>[8]</sup> pharmacological agents,<sup>[9]</sup> lasers,<sup>[10]</sup> photodynamic therapy,<sup>[11]</sup> oxygen pressure,<sup>[12]</sup> and scar penetration of neovascularisation.<sup>[13,14]</sup>

Delay procedure has been shown to have significant impact on flap survival.<sup>[15]</sup> Although the developments in the techniques of axial and free flaps have decreased the need for this procedure, many surgical and non-surgical delay procedures have been defined.

Delay procedure is frequently performed to increase the viability of flaps. Classically, this procedure is performed 2–3 weeks before the surgery. Axial pattern skin flaps can also benefit from the delay procedure. In previous studies, rat skin island flap models have been used as a model for the delay procedure.<sup>[16]</sup> As a result, the delay procedure has been shown to have positive effects on flap circulation and has been adopted by plastic surgeons.

The delay procedure performed on the TRAM flap is one of the best examples of this application. In the practice of plastic surgery, the delay procedure can be performed with surgical techniques (incisions, saturations and ligations), pharmacological agents (vasoconstrictor and vasodilator agents), lasers<sup>[17]</sup> and mechanical impulses (extracorporeal shock wave).<sup>[18]</sup>

Subdermal plexus from the adjacent skin and musculocutaneous/septocutaneous perforators are the two major blood supplies of skin flaps. In this study, our aim was to compare the effectiveness of subdermal and perforator delay procedures with clinical and histological findings in a rat skin flap model.

## MATERIALS AND METHODS

This study was conducted in the Istanbul University Cerrahpasa Medical School Animal Research Laboratory with the approval of Istanbul University local Ethical Committee. Twenty-four female Sprague-Dawley rats averaging at 225 g weight were used. 9 cm × 3 cm caudally based dorsal flaps were designed on the back skin of each animal [Figure 1].<sup>[19]</sup> The flaps were raised at the subcutaneous level, and subdermal plexus was preserved during this procedure. The deep fascia was not included in the flap to avoid any vascular supply from the subfascial and suprafascial plexuses.

Animals were divided into three groups with eight in each group.

### First experimental group (subdermal delay)

In this group, the first delay procedure was performed in which the flaps were only incised at the cranial edge. One week later, a second delay procedure involving the excision of the lateral edges was performed [Figure 2]. This procedure was performed in two stages to reduce the surgical stress and to enhance the new vessel formation. The connection of the flap with the underlying fascia was kept intact during these procedures. Four weeks later, the first delay procedure, the flaps were raised completely and sutured back to their location with 4.0 polypropylene sutures together with the control group.

### Second experimental group (perforator delay)

In this group, a delay procedure was performed in which the flaps were only incised at the cranial edge, and the flaps were dissected from the underlying fascia. A silicon sheet was placed under the flap. Lateral edges were kept intact [Figure 3]. Four weeks after the delay procedure, the flaps were raised completely and sutured back to their location with 4.0 polypropylene sutures together with the control group.

### Control group

Caudally based flaps were raised without any delay and sutured back to their location with 4.0 polypropylene sutures along with the other animals from the experimental groups [Figure 4].

Post-operative follow-up was performed by photographing of the flaps at 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and 7<sup>th</sup> days following the final surgery. At the end of the follow-up period, flap areas




Figure 1: Mc Farlane dorsal skin flap model


were marked and photographed. The marked flap areas were copied to an X-ray film, were photographed. Flap area calculations were performed with the VistaMetrix software (© SkillCrest, LLC).

All the animals were sacrificed with high-dose barbiturate, and the tissue samples were taken for the histological examination. Samples from the proximal and distal portion of the flaps were prepared in 10% formaldehyde. Lymphocyte and collagen densities were evaluated under light microscopy. H and E staining and Masson staining were used, and scoring was performed with a cell density scale.

Kruskal–Wallis and Dunn's multiple comparison tests were used for statistical analysis.

Angiography was performed on two rats on the 7<sup>th</sup> post-operative day before sacrifice.

Inferior vena cava and aorta were cannulated [Figure 5]. A mixture of 40 ml 40% barium sulphate and 10% gelatine was prepared inside saline solution.<sup>[20]</sup> This solution was injected into the vessels, and the animals were kept in – 20°C overnight. The flaps were removed from the animals on the next day. The flaps were stabilised on plastic plates from corners. The flaps were photographed on high-resolution mammography machines.



**Figure 2:** (a) Superior and (b) lateral incisions on the first experimental group



**Figure 4:** Skin flaps raised on caudal pedicle, 4 weeks after the delay

## RESULTS


During the follow-ups, the flaps of the control group started to show discoloration on the first post-operative day. Demarcation of the necrotic areas became visible between on 5<sup>th</sup> and the 7<sup>th</sup> day postoperatively. The average of the necrotic area on the flaps was 21.9% ( $\pm 7.70$ ).

In the experimental groups, two animals on Group 1 and three animals on Group 3 showed some discoloration on the first post-operative day. These findings were all confined in areas which were 1–2 cm in diameter. On the follow-ups, these areas recovered completely; hence, there was no necrosis at the end of the study on both of these groups. These results were significantly different than the control group ( $P < 0.0001$ ) [Figure 6].


### Evaluation of histological findings

Lymphocyte and collagen densities were evaluated under light microscopy [Figure 7]. Seven pairs of samples were taken from each group. Density scoring was performed in accordance with the following scale:

- 0: None
- 1+: Low density
- 2+: Medium density
- 3+: High density.




**Figure 3:** (a) Silicone sheet used in the second experimental group. (b) Silicone sheet is placed under the flap from a superior incision



**Figure 5:** (a) Dissection (b) Cannulation for the angiographic studies





**Figure 6:** Photographs of the flaps at the end of the study from selected samples. (a) Control group. (b) First experimental group. (c) Second experimental group

There was only a statistically significant difference in the collagen density results of the control group and the proximal of the first experimental group ( $P = 0.0315$ ) [Figure 8].

In the samples taken from the distal flap portions of the experimental groups, there was a significant increase in the vascular density; however, this parameter has not been scored.


### Evaluation of angiography results

The angiography images were magnified and evaluated. In the subjective evaluation of the images, the experimental groups showed a slight increase in the amount of vascularisation when compared to the control group. This difference was especially visible in the first experimental group [Figure 9].

## DISCUSSION

Delay procedure on the skin flaps makes larger-sized flaps possible. The exact mechanisms behind this procedure are not clear. Many studies have been conducted to answer this question.<sup>[21-24]</sup> The amount of vascularity increases required for an adequate delay still remains as an important question. This will depend on many variables as well as the delay technique. Although we believe that it will not be possible to quantitatively measure this amount, the future studies involving more variables might give us valuable insight into this point.

At the end of the study, none of the experimental groups showed a permanent circulation problem or necrosis.



**Figure 7:** Collagen density shown with the Masson's trichrome staining. (a) Control group, (c) first experimental group, (e) second experimental group. Lymphocyte density shown with haematoxylin and eosin staining. (b) Control group, (d) first experimental group, (f) second experimental group

This finding can be further supported by the fact that no distortion on the epidermis was found microscopically. Under light microscopy, there was an increase in the vascular density and neovascularisation. The 4 weeks of delay time used in our study was probably effective in reducing the amount of necrosis for the perforator delay group. There was a significant amount of discoloration after the flap elevation in three animals in this group. This discoloration did not eventually result in permanent necrosis. We think that this might have been different if a shorter duration of delay time was chosen. Although 2 weeks of delay time was shown to be adequate for dermal delay techniques,<sup>[25]</sup> to the best of our knowledge, there was not any previous study on the adequate amount of delay time for the perforator technique. We thought that the dermal vascularity increase will be much slower in perforator delay group, which in principle will be similar to the expander techniques in which the perforator circulation is similarly interrupted.

In our study, there was not any significant increase in the lymphocyte density in the experimental groups. These results suggest that the flaps were not in an inflammation state. Collagen density has increased in both experimental groups. The collagen formation suggests an increase in




Figure 8: (a) Average of lymphocyte density scorings. (b) Average of collagen density scorings




Figure 9: Angiography results from selected samples. (a) Control group. (b) First experimental group

neovascularisation and fibroblast activity. Zhou *et al.* have shown in a study that the injection of fibroblast growth factor and collagen matrix increases neovascularisation on rabbits.<sup>[26]</sup> Rao *et al.* have also shown an increase in angiogenesis with collagen matrix and endothelial and mesenchymal stem cells.<sup>[27]</sup> All of these studies indicate that the importance of collagen matrix for neovascularisation.

The angiographic imaging performed in our study was technically demanding. We believe that angiographic studies on small animals such as rats will usually result in a low success rate and are difficult to interpret. A slight increase in vascularisation in experimental groups was detected with these studies. This increase in vascularisation has to be further investigated for venous and arterial components. Lymphatic vessels also seem to play an important role on flap viability. We believe that these can be further investigated with lymphoscintigraphic imaging on larger mammals such as pigs.

It is crucial to point out the fact that regional variations have to be taken into consideration. These results may turn out differently in other flap locations. Differences between human and rat physiologies are another

factor to be considered.<sup>[28]</sup> These differences might be particularly important in neovascularisation and delay durations. There are many unknowns in the equation of flap physiology. Although many of these factors have been clarified by previous studies, it should not be discouraging to see that similar studies reveal different results.

Histological results show that collagen and vascular densities were lower on the second experimental group. Although there was not any permanent necrosis in any of the experimental groups, these results might suggest that necrosis would be more likely in this second experimental group. The incisions that were performed in stages on the first experimental group might have positive effects on flap circulation by decreasing the amount of stress and ischaemia on the tissues. It seems that by gradually increasing the stress we can enhance the circulation of the flap. In a previous study, Callegari *et al.* have showed that choked vessels can enlarge with the ligation of blood vessels from the adjacent angiosomes.<sup>[29]</sup> This result is consistent with our findings. Lateral incisions performed in the first experimental group do have a similar effect described in the earlier study of George *et al.* In this study, an occlusion clamp was used instead of the lateral incisions performed in our study. We believe that both of these techniques would open adjacent choked vessels as described earlier which would yield to similar results regarding the flap survival.

## CONCLUSION

Our results support the idea that the delay procedure has a positive effect on flap viability. In conclusion, flap viability and histological data at the end of the study suggest that

either subdermal or perforator circulation can be interrupted for a successful delay procedure. The papers published in angiosomes,<sup>[30]</sup> and the perforasomes<sup>[31]</sup> have changed our understanding of the random flaps in a profound way. These theories actually made it clear that there are actually no random flaps. However, in practice when the surgeon is unable to identify the angiosomes and the perforasomes associated with the skin area where the flap is to be raised the term random flap is used. Circulation problems are common with these types of flaps since they are raised without the knowledge of their vascular supply. These problems usually limit the usage of these of flaps on smaller defects.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Kaufman T, Eichenlaub EH, Levin M, Hurwitz DJ, Klain M. Tobacco smoking: Impairment of experimental flap survival. *Ann Plast Surg* 1984;13:468-72.
- Cutting C. Critical closing and perfusion pressures in flap survival. *Ann Plast Surg* 1982;9:524.
- German C, Koss N. Infection and flap survival. *Surg Forum* 1977;28:533-5.
- Reinisch J, Myers B. The effect of local anesthesia with epinephrine on skin flap survival. *Plast Reconstr Surg* 1974;54:324-7.
- Earle AS, Fratianne RB, Nunez FD. The relationship of hematocrit levels to skin flap survival in the dog. *Plast Reconstr Surg* 1974;54:341-44.
- Seyhan T, Deniz M, Borman H, Ulusal Güzel B, Çağlar B. Comparison of two different vascular delay methods in a rat cranial epigastric perforator flap model. *Ann Plast Surg* 2010;64:89-92.
- Yan H, Brooks D, Jackson WD, Angel MF, Akdemir O, Zhang F, *et al.* Improvement of prearterialized venous flap survival with delay procedure in rats. *J Reconstr Microsurg* 2010;26:193-200.
- Huemer GM, Froschauer SM, Pachinger T, Kwasny O, Schoffl H. A comparison of pretreatment with a topical combination of nonivamide and nicoboxil and surgical delay in a random pattern skin flap model. *J Plast Reconstr Aesthet Surg* 2009;62:914-9.
- Karacaoglu E, Yuksel F, Turan SO, Zienowicz RJ. Chemical delay: An alternative to surgical delay experimental study. *Ann Plast Surg* 2002;49:73-80.
- Acikel C, Kenkel JM, Nojima K, Hoopman JE, Saboorian MH, Brown SA, *et al.* Evaluation of flashlamp-pumped pulsed-dye laser (585 nm) in nonsurgical delay of dorsal rat cutaneous flaps. *Plast Reconstr Surg* 2005;115:2032-41.
- Belmont MJ, Marabelle N, Mang TS, Hall R, Wax MK. Effect of photodynamic therapy on revascularization of fasciocutaneous flaps. *Laryngoscope* 2000;110:942-5.
- Stewart RJ, Moore T, Bennett B, Easton M, Newton GW, Yamaguchi KT, *et al.* Effect of free-radical scavengers and hyperbaric oxygen on random-pattern skin flaps. *Arch Surg* 1994;129:982-7.
- Serin M, Bayramicli M. Evaluation of scar penetrating neovascularisation in a rat epigastric flap model. *J Plast Surg Hand Surg* 2015;19:1-5.
- Edstrom LE, Robson MC, Hopp D. Survival of skin flaps based on scar. *Ann Plast Surg* 1981;7:44-7.
- Holzbach T, Neshkova I, Vlaskou D, Konerding MA, Gansbacher B, Biemer E, *et al.* Searching for the right timing of surgical delay: Angiogenesis, vascular endothelial growth factor and perfusion changes in a skin-flap model. *J Plast Reconstr Aesthet Surg* 2009;62:1534-42.
- Komorowska-Timek E, Timek TA, Brady S, Buncke HJ. Perfusion, viability, and pedicle dependence in acute and delayed rat island skin flaps. *Microsurgery* 2007;27:146-51.
- Cohen M, Kattan A, Gat A, Almogi N, Katzir A, Villan A, *et al.* Comparison of flash lamp pulsed-dye laser (585 nm) and conventional surgery in the delay of random dorsal rat flaps. *Lasers Surg Med* 1999;25:178-86.
- Reichenberger MA, Keil H, Mueller W, Herold-Mende C, Gebhard MM, Germann G, *et al.* Comparison of extracorporeal shock wave pretreatment to classic surgical delay in a random pattern skin flap model. *Plast Reconstr Surg* 2011;127:1830-7.
- Mcfarlane RM, Deyoung G, Henry RA. The design of a pedicle flap in the rat to study necrosis and its prevention. *Plast Reconstr Surg* 1965;35:177-82.
- Quinodoz P, Quinodoz M, Nussbaum JL, Montandon D, Pittet B. Barium sulphate and soft-tissue radiology: Allying the old and the new for the investigation of animal cutaneous microcirculation. *Br J Plast Surg* 2002;55:664-7.
- Reinisch JF. The pathophysiology of skin flap circulation. The delay phenomenon. *Plast Reconstr Surg* 1974;54:585-98.
- O'Neill JV, Toomey JM, Snyder GG 3<sup>rd</sup>. Factors involved in the skin flap delay phenomenon. *Surg Forum* 1975;26:528-30.
- Cutting C, Bardach J, Rosewall D. Skin flap delay procedures: Proximal delay versus distal delay. *Ann Plast Surg* 1980;4:293-6.
- Isoda S, Yano Y, Jin Y, Walters HL 3<sup>rd</sup>, Kondo J, Matsumoto A, *et al.* Influence of a delay on latissimus dorsi muscle flap blood flow. *Ann Thorac Surg* 1995;59:632-7.
- Restifo RJ, Ahmed SS, Isenberg JS, Thomson JG. Timing, magnitude, and utility of surgical delay in the TRAM flap: I. Animal studies. *Plast Reconstr Surg* 1997;99:1211-6.
- Zhou J, Zhao Y, Wang J, Zhang S, Liu Z, Zhen M, *et al.* Therapeutic angiogenesis using basic fibroblast growth factor in combination with a collagen matrix in chronic hindlimb ischemia. *ScientificWorldJournal* 2012;2012:652794.
- Rao RR, Peterson AW, Ceccarelli J, Putnam AJ, Stegemann JP. Matrix composition regulates three-dimensional network formation by endothelial cells and mesenchymal stem cells in collagen/fibrin materials. *Angiogenesis* 2012;15:253-64.
- Hallock GG. The rat TRAM flap: A human analogue? *Plast Reconstr Surg* 1995;96:233-4.
- Callegari PR, Taylor GI, Caddy CM, Minabe T. An anatomic review of the delay phenomenon: I. Experimental studies. *Plast Reconstr Surg* 1992;89:397-407.
- Taylor GI, Palmer JH. The vascular territories (angiosomes) of the body: Experimental study and clinical applications. *Br J Plast Surg* 1987;40:113-41.
- Saint-Cyr M, Wong C, Schaverien M, Mojallal A, Rohrich RJ. The perforasome theory: Vascular anatomy and clinical implications. *Plast Reconstr Surg* 2009;124:1529-44.