Case Report

Recurrent Acute Pancreatitis with Anomalous Pancreaticobiliary Ductal Union (Komi's Type IIIc3) with Santorinocele in a Child: A Rare Case Report

Vineet Mishra, A. J. Joseph, Amit Kumar Dutta, Sudipto Dhar Chowdhury, Reuben Thomas Kurien, Saurabh Jaiswal, Itish Patnaik, Parika Kalra, Sayd Mohamad Rajeeb

Department of
Gastrointestinal Sciences,
Christian Medical College,
Vellore, Tamil Nadu, India

ABSTRACT

Anomalous pancreaticobiliary ductal union (APBDU) can cause recurrent acute pancreatitis. We describe the case of a 16-year-old boy with recurrent acute pancreatitis. The discussion provides a review of recent literature, supporting use of various diagnostic modalities and surgery as choice of treatment.

KEYWORDS: *Anomalous pancreaticobiliary ductal union, choledochal cyst, Komi classification, recurrent acute pancreatitis, santorinocele*

INTRODUCTION

Recurrent acute pancreatitis is defined as more than two discrete episodes of diagnosed acute pancreatitis at least 2 months apart with asymptomatic interval period. The most common causes are alcohol abuse and gallstone disease. Anomalous pancreaticobiliary ductal union (APBDU) has been described as cause for idiopathic recurrent acute pancreatitis in both pediatric and adult population. This anomaly can cause acute pancreatitis, recurrent acute pancreatitis, and recurrent cholangitis.^[1] It should be diagnosed and surgically treated before complications or carcinomas develop.^[1,2]

We report a case of an APBDU with choledochal cyst and santorinocele (with calculi) associated with probable choledocholithiasis presented with recurrent acute pancreatitis and complicated by acute cholangitis.

CASE REPORT

We admitted a 16-year-old boy with recurrent abdominal pain with a history of acute necrotizing pancreatitis (contrast-enhanced computed tomography [CECT] done elsewhere) 6 months before present admission. Etiological factors including trauma, drug, family history, and metabolic factors were excluded from the study. He was again admitted 2 months later with mild acute pancreatitis elsewhere.

On evaluation, magnetic resonance cholangiopancreatography (MRCP) [Figure 1 above] and

CECT showed that common hepatic duct was dilated till terminal portion (choledochal cyst) with T2 hyperintense focus in terminal common bile duct (CBD) near ampulla suggestive of calculi, mildly prominent pancreatic duct draining separately into the second part of duodenum (like classic pancreas divisum) with moderate bilobar intrahepatic biliary dilatation with liver function test (LFT) of bilirubin total and direct of 1.2/0.4 mg/dl, aspartate aminotransferase (AST)/alanine aminotransferase (ALT) of 112/294 (<40 IU/L) IU/L, and alkaline phosphatase (ALP) of 610 U/L (<125 IU/L) and normal amylase and lipase. Physical examination was normal.

He developed high-grade fever with jaundice in hospital (bilirubin total and direct 3.6/2.6 mg/dl, AST/ALT 204/369 IU/L and ALP 902 IU/L). Endoscopic retrograde cholangiopancreatography (ERCP) with sphincterotomy with plastic stenting was done; no CBD calculi noted. His fever resolved and LFT improved, but abdominal pain was persisting. Repeat ERCP for CBD clearance and minor papillotomy was performed. Cholangiogram revealed complex connection of CBD, main pancreatic duct (MPD), and accessory pancreatic duct (APD) with cystic dilatation in APD [Figure 1 below]. Cholangiogram revealed no

Address for correspondence: Dr. Vineet Mishra, Department of Gastrointestinal Sciences, Wellcome Trust Building, Christian Medical College and Hospitals, Vellore - 632 004, Tamil Nadu, India.
E-mail: vinegar2805@gmail.com

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Mishra V, Joseph AJ, Dutta AK, Chowdhury SD, Kurien RT, Jaiswal S, et al. Recurrent acute pancreatitis with anomalous pancreaticobiliary ductal union (Komi's Type IIIc3) with santorinocele in a child: A rare case report. J Dig Endosc 2017;8:196-8.

Access this article online

Quick Response Code:

Website: www.jdeonline.in

DOI: 10.4103/jde.JDE_22_17

Figure 1: Preprocedure magnetic resonance cholangiopancreatography image (above) and endoscopic retrograde cholangiopancreatography image - santorinicele (below)

CBD filling defect; contrast through major papilla also filled MPD and APD with a 7 mm filling defect in APD before its opening to minor papilla and pancreaticogram through minor papilla also filled MPD and CBD with both MPD and APD appeared to additionally communicate with CBD [Figure 2 above and below]. APD stenting was done following minor duct papillotomy and stent was removed 5 days later. The patient remained asymptomatic in terms of abdominal pain and there were no further episodes of fever or jaundice on 3-month follow-up. Need for cholecystectomy and pancreatoduodenectomy was discussed with relatives.

DISCUSSION

In APBDU, pancreatic and bile ducts join outside the duodenum and form a long common channel (usually >15 mm). The anomalous union is often associated with a choledochal cyst.^[3] Association of APBDU with choledochal cyst may be present in up to 90% of cases.^[4]

Gallbladder carcinomas develop in 15%–40%, with lesser frequency if associated with choledochal cyst.^[2]

APBDU is a rare anomaly with frequency of 1.5%–3.2%.^[4] Its diagnosis needs a high index of suspicion. Komi^[5] proposed a new classification of APBDU [Figure 3], dividing it into three types. In

Figure 2: Choledochal cyst and anomalous pancreaticobiliary ductal union in endoscopic retrograde cholangiopancreatography images

Type III, APBDU is complicated with a patent APD with or without complicated network of ducts and is subclassified into Types IIIa, IIIb, and IIIc.^[4] There are scarce data from India,^[6,7] and none of them provides the prevalence of APBDU, in particular the new Komi Type III.

APBDU and choledochal cyst association^[6] may be due to congenital weakness of CBD and reflux of the pancreatic juices in CBD. Subtype IIIC3 APBDU has been reported to be associated with ductal calculi in up to 43.8% of cases, but its mere presence causes symptoms is controversial.^[8]

Santorinicele is a focal cystic dilatation of (which may contain protein plug or calculi) the termination of APD at the minor papilla.^[9] This anomaly has been described in patients with pancreas divisum and recurrent acute pancreatitis.^[9] Pancreas divisum as a cause for acute recurrent pancreatitis is still debatable and undiagnosed santorinicele with calculi/plug or APBUD may be playing a role in those cases.

Patients with pancreas divisum who may get benefit from endoscopic therapy is still debatable.^[8] Santorinicele is due to stenotic accessory duct papilla or cause of relapsing dysfunction remains controversial. As per previous studies, the presence of santorinicele may help in selecting patients who might benefit from endoscopic treatment.^[9]

TYPES		DESCRIPTION
IA	Type I	Have single papilla & The Common Hepatic and pancreatic ducts join each other at a right angle with a non dilated common channel.
	A	
IB	Type I	Have single papilla & The Common Hepatic and pancreatic ducts join each other at a right angle with a dilated common channel.
	B	
IIA	Type II	Have single papilla & The Common Hepatic and pancreatic ducts join each other at An acute angle with a non dilated common channel.
	A	
IIB	Type II	Have single papilla & The Common Hepatic and pancreatic ducts join each other at An acute angle with a dilated common channel.
	B	
IIIA	Type III	Have two papilla & are equivalent to the classic pancreas divisum with biliary dilatation.
	A	
IIIB	Type III	Have two papilla & are characterized by the absence of the Wirsung's duct.
	B	
IIIC1	C1	Have two papilla & contain a tiny communicating duct between the main duct and the accessory ducts.
IIIC2	C2	Have two papilla with & characterized by a common channel made up of common and accessory ducts of equal caliber.
IIIC3	C3	Have two papilla with intricate network of dilated ducts that join each other by total or partial dilatation of the ductal system.

Figure 3: Komi's classification of anomalous pancreaticobiliary ductal union

ERCP is the investigation of choice;^[7] MRCP has up to 82% detection rate. Endoscopic ultrasound has been reported as a good method for the detection of APDU.^[10] Our patient had separate major and minor papillae, communicating with each other and with the dilated CBD and santorinocoele (with probable calculi) without obvious CBD obstruction (choledochocoele). After reviewing the literature, we found that this pattern was consistent with the Type IIIC3 anomaly.^[5] Type IB, IIB, and IIIC3 APBDU are particularly associated with recurrent pancreatitis and are treated surgically, pylorus-preserving pancreatoduodenectomy being the choice for Type IIIC3.^[6] In APBDU patient without dilated CBD, cancer risk is low, and hence, cholecystectomy without duct resection is recommended in few studies.

CONCLUSION

In patients with idiopathic recurrent acute pancreatitis, particularly young patients, APBDU should be a

differential. The complications and long-term sequelae of acute pancreatitis and risk of cancer in young patients warrant early and judicious management.^[6,7]

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Elnemr A, Ohta T, Kayahara M, Kitagawa H, Yoshimoto K, Tani T, *et al.* Anomalous pancreaticobiliary ductal junction without bile duct dilatation in gallbladder cancer. *Hepatogastroenterology* 2001;48:382-6.
2. Funabiki T, Matsubara T, Miyakawa S, Ishihara S. Pancreaticobiliary maljunction and carcinogenesis to biliary and pancreatic malignancy. *Langenbecks Arch Surg* 2009;394:159-69.
3. Todani T, Watanabe Y, Narusue M, Tabuchi K, Okajima K. Congenital bile duct cysts: Classification, operative procedures, and review of thirty-seven cases including cancer arising from choledochal cyst. *Am J Surg* 1977;134:263-9.
4. Kimura K. Studies in 28 cases of congenital cystic dilatation of the common bile duct in the adults. Roentgenological features and a union between the choledochus and the main pancreatic duct. *Jpn J Gastroenterol* 1976;73:61-74.
5. Komi N. New classification of anomalous arrangement of the pancreaticobiliary duct (APBD) in the choledochal cyst; A proposal of new Komi's classification of APBD. *J Jpn Pancreas Soc* 1991;6:234-43.
6. Kochhar R, Singhal M, Nagi B, Lal A, Dutta U, Singh K. Prevalence of type III anomalous pancreaticobiliary junction in a tertiary care hospital of North India. *JOP* 2009;10:383-6.
7. Misra SP, Gulati P, Thorat VK, Vij JC, Anand BS. Pancreaticobiliary ductal union in biliary diseases. An endoscopic retrograde cholangiopancreatographic study. *Gastroenterology* 1989;96:907-12.
8. Kaneko K, Ando H, Ito T, Watanabe Y, Seo T, Harada T, *et al.* Protein plugs cause symptoms in patients with choledochal cysts. *Am J Gastroenterol* 1997;92:1018-21.
9. Eisen G, Schutz S, Metzler D, Baillie J, Cotton PB. Santorinicele: New evidence for obstruction in pancreas divisum. *Gastrointest Endosc* 1994;40:73-6.
10. Sugiyama M, Atomi Y. Endoscopic ultrasonography for diagnosing anomalous pancreaticobiliary junction. *Gastrointest Endosc* 1997;45:261-7.