Commentary

With the rapid and widespread availability of neuroimaging, brain tumors are being more commonly diagnosed nowadays. The estimated annual incidence for primary and metastatic brain tumors is 9/100,000 and 8.3/100,000, respectively.^[1] Psychiatric problems can often be noted in patients with brain tumors, ranging from 10% to 40%.^[1-4] However, a study by Keschner *et al.* reported that 78% of 530 patients with brain tumors had psychiatric symptoms.^[5] These wide differences may be due to different study designs or questionnaires used. The psychiatric symptoms noted may be diverse, such as depression, anxiety, mood disorders, schizophrenia, psychosis, personality changes, abulia, and delusions and hallucinations.^[1,4,6-8]

Although commonly present concomitantly with neurologic features, psychiatric symptoms may be the only manifestation of brain tumors (so-called neurologically silent tumors).^[1,6] The most common brain neoplasms with neuropsychiatric symptoms include gliomas, meningiomas, and metastasis and rarely pituitary adenomas, epidermoids, craniopharyngiomas, and germ cell tumors.^[1,2,9-11] With regard to the association between tumor location and psychiatric symptoms, although few studies have reported higher incidence of depression and memory disturbances in left hemispheric tumors, many other studies have refuted this association.^[12-14] However, the association between specific areas in the brain and psychiatric manifestations can be seen, such as psychotic symptoms and pituitary tumors, anorexia, and hypothalamic tumors.^[1,15]

In an autopsy study, Furgal-Borzych *et al.* observed a higher incidence of pituitary adenomas in suicidal victims (44.7%) as compared to 18.7% in nonsuicidal group.^[16] Gupta and Kumar retrospectively analyzed 79 patients with brain tumors to examine the presence of psychiatric comorbidity in benign brain tumors. They included only those patients with psychiatric symptoms of several weeks to several months duration antedating a diagnosis of a brain tumor. Meningiomas constituted more than 90% of cases (72/79) and 15 (21%) meningioma cases (8 males and 7 females) presented with psychiatric symptoms in the absence of neurological symptoms. It was noted that affective disorders were the most common presentation. They concluded that psychiatric symptoms might be the only initial manifestations of meningiomas and such patients must be investigated by brain imaging studies even without neurological signs/symptoms.^[2]

It has also been observed that symptoms such as depression and anxiety are often underdiagnosed in brain tumor patients leading to reduced access to psychiatric medications.^[3,12] In an analysis of 598 high-grade glioma patients, Litofsky *et al.* found a very high rate (93%) of depressive symptoms reported by patients but only 15% of them were recognized by physicians. They concluded that physician recognition of depression is low and is associated with undertreatment of such depressed patients.^[12]

Quality of life also appears to be affected in brain tumor patients with psychiatric symptoms.^[4,12,14,17,18] Charkavliuk *et al.* analyzed 271 patients with brain tumors to evaluate the prevalence of psychiatric symptoms and psychotropic treatment and their relation to quality of life. They noted symptoms of psychiatric disorders in 8.1% of cases. Depression was the most common, followed by schizophrenia spectrum disorders, and only 4% of them were on medications. They concluded that psychiatric symptoms such as depression and anxiety may significantly hamper the quality of life.^[4] Further, it has also been observed that the longevity might also be decreased in brain tumor patients with depression.^[12,18]

In such brain tumor patients with psychiatric problems, symptoms may be misleading, often complicating clinical picture. Furthermore, in some instances, neurological syndromes may be confused with psychiatric conditions.^[19] Hence, it is recommended that in any patient above 40 years, a change in neurobehavioral statuses such as new-onset psychosis, mood or memory symptoms, occurrence of new or atypical symptoms, personality changes, and anorexia without body dysmorphic symptoms should have a neuroimaging workup. Similar recommendations also apply to patients with psychiatric presentations accompanied by specific neurologic changes or with poor response to psychotropic medications.^[1,6,7,20]

To conclude, I feel that psychiatrists and neurophysicians/neurosurgeons should work in liaison to avoid underdiagnosis and undertreatment of psychiatric disorders in brain tumor patients and to improve the quality of life and outcome of such patients.

G. Lakshmi Prasad

Department of Neurosurgery, Kasturba Medical College,
Manipal University, Manipal, Karnataka, India

Address for correspondence: Dr. G. Lakshmi Prasad,
Department of Neurosurgery, Kasturba Hospital, Room 12,
OPD Block, Manipal - 576 104, Karnataka, India.
E-mail: lakshmi.prasad@manipal.edu

REFERENCES

- Madhusoodanan S, Ting MB, Farah T, Ugur U. Psychiatric aspects of brain tumors: A review. *World J Psychiatry* 2015;5:273-85.
- Gupta RK, Kumar R. Benign brain tumours and psychiatric morbidity: A 5-years retrospective data analysis. *Aust N Z J Psychiatry* 2004;38:316-9.
- Arnold SD, Forman LM, Brigidi BD, Carter KE, Schweitzer HA, Quinn HE, *et al.* Evaluation and characterization of generalized anxiety and depression in patients with primary brain tumors. *Neuro Oncol* 2008;10:171-81.
- Charkavliuk A, Steibliene V, Tamasauskas S, Deltuva VP, Tamasauskas A, Bunevicius A. Psychiatric histories and treatment in patients with brain tumors: Association with psychological distress symptoms and health related quality of life. *Biol Psychiatry Psychopharmacol* 2014;16:79-84.
- Keschner M, Bender MB, Strauss I. Mental symptoms associated with brain tumor: A study of 530 verified cases. *JAMA* 1938;110:714-8.
- Betul O, Ipek M. Brain tumor presenting with psychiatric symptoms. *J Neuropsychiatry Clin Neurosci* 2011;23:E43-4.
- Moise D, Madhusoodanan S. Psychiatric symptoms associated with brain tumors: A clinical enigma. *CNS Spectr* 2006;11:28-31.
- Madhusoodanan S, Danan D, Moise D. Psychiatric manifestations of brain tumors: Diagnostic implications. *Expert Rev Neurother* 2007;7:343-9.
- Kar SK, Das KK, Jaiswal AK, Jaiswal S. Mood disorder as an early presentation of epidermoid of quadrigeminal cistern. *J Neurosci Rural Pract* 2017;8:443-5.
- Boele FW, Rooney AG, Grant R, Klein M. Psychiatric symptoms in glioma patients: From diagnosis to management. *Neuropsychiatr Dis Treat* 2015;11:1413-20.
- Malbari F, Gershon TR, Garvin JH, Allen JC, Khakoo Y, Levy AS, *et al.* Psychiatric manifestations as initial presentation for pediatric CNS germ cell tumors, a case series. *Childs Nerv Syst* 2016;32:1359-62.
- Litofsky NS, Farace E, Anderson F Jr., Meyers CA, Huang W, Laws ER Jr., Glioma Outcomes Project Investigators. Depression in patients with high-grade glioma: Results of the Glioma Outcomes Project. *Neurosurgery* 2004;54:358-66.
- Hahn CA, Dunn RH, Logue PE, King JH, Edwards CL, Halperin EC. Prospective study of neuropsychologic testing and quality-of-life assessment of adults with primary malignant brain tumors. *Int J Radiat Oncol Biol Phys* 2003;55:992-9.
- Pelletier G, Verhoef MJ, Khatri N, Hagen N. Quality of life in brain tumor patients: The relative contributions of depression, fatigue, emotional distress, and existential issues. *J Neurooncol* 2002;57:41-9.
- Madhusoodanan S, Opler MG, Moise D, Gordon J, Danan DM, Sinha A, *et al.* Brain tumor location and psychiatric symptoms: Is there any association? A meta-analysis of published case studies. *Expert Rev Neurother* 2010;10:1529-36.
- Furgal-Borzych A, Lis GJ, Litwin JA, Rzepecka-Wozniak E, Trela F, Cichocki T. Increased incidence of pituitary microadenomas in suicide victims. *Neuropsychobiology* 2007;55:163-6.
- Sherwood PR, Given BA, Donovan H, Baum A, Given CW, Bender CM, *et al.* Guiding research in family care: A new approach to oncology caregiving. *Psychooncology* 2008;17:986-96.
- Mainio A, Hakko H, Timonen M, Niemelä A, Koivukangas J, Räsänen P. Depression in relation to survival among neurosurgical patients with a primary brain tumor: A 5-year follow-up study. *Neurosurgery* 2005;56:1234-41.
- Butler C, Zeman AZ. Neurological syndromes which can be mistaken for psychiatric conditions. *J Neurol Neurosurg Psychiatry* 2005;76 Suppl 1:i31-8.
- Madhusoodanan S, Danan D, Brenner R, Bogunovic O. Brain tumor and psychiatric manifestations: A case report and brief review. *Ann Clin Psychiatry* 2004;16:111-3.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/jnpr.jnpr_132_17

How to cite this article: Prasad GL. Psychiatric disorders and brain tumors. *J Neurosci Rural Pract* 2017;8:446-7.

© 2017 Journal of Neurosciences in Rural Practice | Published by Wolters Kluwer - Medknow