Case Series

Osteolytic Skull Lesions—Our Experience

B. V. Sandeep, Manpreet Singh Banga, Suniti Kumar Saha, Kaushik Roy

Department of Neurosurgery,
Nil Ratan Sircar Medical
College and Hospital,
Kolkata, West Bengal, India

ABSTRACT

Objective: To present an overview of varied clinical presentations, investigations and treatment options for Osteolytic skull lesions. **Study Design:** It is a prospective study. **Materials and Methods:** We conducted this study from January 2013 to December 2015 in the Department of Neurosurgery, Nil Ratan Sircar Medical College and Hospital, Kolkata. During this period, 14 patients presented with osteolytic skull lesions through the outpatient department. All patients were thoroughly investigated with appropriate hematological and radiological investigations and treated following admission, and surgery was performed in the Neurosurgery Department. All were followed regularly in OPD. **Results:** Total 14 patients were included in the study. Amongst these 7 were male and 7 female. Age group of patients ranged from 5 to 72 years. Of 14 cases, three cases had dermoid cyst, four cases had metastasis, and one each case had epidermoid cyst, intradiploic meningioma, benign cystic lesion, tuberculosis, histiocytosis X, hemangioma, and osteomyelitis. All underwent diagnostic/therapeutic procedures and referred for Radio or chemotherapy where indicated. **Conclusion:** All scalp/skull lesions need careful clinical correlation, appropriate radiological investigations to establish diagnosis and subject them to suitable treatment.

KEYWORDS: Osteolytic skull lesion, skull lesions, skull tumors

INTRODUCTION

Osteolytic skull lesion is a rare entity among neurosurgical patients. It constitutes about 1% of all skull tumors. Differential diagnosis is broad which includes both benign and malignant lesions. Majority of these are benign. Lesions usually present as an enlarging hard or soft mass which may be painful or painless over the skull with or without tenderness. Lesions are broadly categorized into congenital, traumatic, inflammatory, and neoplastic lesions.^[1]

In children and young adults, osteolytic skull lesions include epidermoid/dermoid, eosinophilic granuloma, encephaloceles, hemangioma, osteoblastoma, and aneurysmal bone cyst. In adults, metastasis and multiple myeloma are the most common diagnosis. In adults, a source for primary lesion should always be searched with the suspicion of metastasis. However in all age groups, surgical defects, fibrous dysplasias, sarcomas, lymphomas, intraosseous

meningiomas, and intracranial tumor with erosion of skull are rare.^[2]

MATERIALS AND METHODS

We conducted a prospective study from January 2013 to December 2015 in the Department of Neurosurgery, Nil Ratan Sircar Medical College and Hospital, Kolkata. During this period, 14 patients presented with osteolytic skull lesions through the outpatient patient. They were treated following admission, and surgery was performed in the Neurosurgery Department.

Inclusion criteria

Patients of all ages and both genders were included in the study.

Address for correspondence: Dr. B. V. Sandeep,

Department of Neurosurgery, 6th Floor, U.N.B. Building, Nil Ratan Sircar Medical College and Hospital, 138, A J C Bose Road, Sealdah, Kolkata, West Bengal - 700 014, West Bengal, India.
E-mail: sandybv200@gmail.com

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/jnpr.jnpr_243_16

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Sandeep BV, Banga MS, Saha SK, Roy K. Osteolytic Skull Lesions—Our experience. J Neurosci Rural Pract 2017;8:434-8.

RESULTS

Gender and age group

Age group range of the patients in this study was wide (5–72 years), and the lesions were of same incidence in both male and female [Tables 1 and 2].

Clinical features

Of 14 cases, eight cases presented with painless swelling. The rest six cases presented as a painful swelling which included cases with metastasis, osteomyelitis, and tuberculosis and a case of benign cystic lesion.

Operative procedure and treatment

Planning of surgery and incisions was taken depending on the site of lesion.

All patients underwent complete excision of lesion including involved bone whenever necessary. Cranioplasty for calvarial defect following surgery was done in seven cases.

Four cases of metastasis after excision of the lesion were referred to the chemoradiotherapy department for further evaluation and management. One case of the tubercular lesion had excision of the lesion followed by

antitubercular therapy. One case of histiocytosis X after excision was referred to the hematology department for further management.

Histopathology

Of 14 cases, three cases had dermoid cyst, four cases had metastasis, and one each case had epidermoid cyst, intradiploic meningioma, benign cystic lesion, tuberculosis, histiocytosis X, hemangioma, and osteomyelitis [Table 3].

DISCUSSION

In a study by Anwar *et al.*, 12 cases of osteolytic skull lesions were treated. Of these, three cases were each of tuberculosis and malignancies, two cases of midline benign lesions, and one each of posttraumatic, infection, and metastasis.^[2]

In our study, 14 cases were treated which included three cases had dermoid cyst, four cases had metastasis, one each case had epidermoid cyst, intradiploic meningioma, benign cystic lesion, tuberculosis, histiocytosis X, hemangioma, and osteomyelitis.

Intracranial dermoid cysts are rare congenital lesions that account for 0.04%–0.7% of all intracranial tumors.^[3] Dermoid cysts originate from embryonic ectodermal cells that are involved in the process of primary neurulation. Dermoid cysts primarily occur in intradural regions; however, rarely, extradural dermoid cysts have been reported. Although the posterior fossa is the most common location for dermoid cysts, a few intradural dermoid cysts originating from skull base have been reported. Extradural dermoid cysts are observed in the posterior fossa, diploe of anterior fontanelle, and orbital region in children and young adult.^[4]

In our study, we encountered three cases of dermoid cysts, of which two patients were male and one was

Table 1: Sex distribution

Number of cases	Males	Females
14	9	5

Table 2: Age distribution

Age (years)	Number of cases
0-20	5
20-40	2
40-60	5
>60	2

Table 3: Histopathology of lesions

Serial number	Gender	Age (years)	Histopathology
1	Male	19	Dermoid cyst
2	Male	15	Benign cystic lesion
3	Male	62	Osteomyelitis
4	Female	45	Metastasis from thyroid follicular carcinoma
5	Female	23	Dermoid cyst
6	Male	27	Dermoid cyst
7	Female	50	Tuberculosis
8	Male	50	Metastasis from adenocarcinoma
9	Male	57	Metastasis from adenocarcinoma
10	Male	72	Metastasis from adenocarcinoma
11	Female	53	Intradiploic meningioma
12	Male	8	Histiocytosis X
13	Female	18	Epidermoid cyst
14	Male	12	Hemangioma

Figure 1: Dermoid cyst over the left frontal bone

female patient. Of three cases, one case was involving occipital midline skull while the other two was involving the left frontal and right temporoparietal skull [Figure 1].

An 18-year-old female presented with painless occipital swelling for 6 months. On excision, the histopathological report suggested epidermoid cyst.

Metastatic tumors to the skull are rare and originate mostly from the lung, breast, or prostate.^[5] Most metastatic skull lesions are asymptomatic.

Skull metastases from a well-differentiated thyroid carcinoma are very uncommon.^[6] In our study, we had a 45-year-old female patient who had undergone thyroidectomy for follicular thyroid malignancy. She presented with a painful mass over the left frontal region, which on excision turned out to be metastasis from the follicular malignancy. We also encountered three male patients who presented with a painful skull lesion in the left frontal bone, occipital bone, and right temporal bone each. The lesion was excised and histopathological examination report stated metastasis from adenocarcinoma. The patient was referred to the chemoradiotherapy department for further evaluation and treatment [Figure 2].

Intraosseous hemangioma is a rare bone tumor accounting for 0.7%–1.0% of all bone tumors. It can occur in all ages but is most common in the fourth and fifth decades. It has a female preponderance (3:1).^[7]

Intraosseous hemangiomas are usually seen in the vertebral column and rarely in the calvarium. Among the calvarial bones, the parietal bone is the most commonly involved, followed by the frontal bone and less frequently by the occipital and temporal bones.^[8]

In our study, we had a 12-year-old male who presented with a painless swelling in the left parietal region which

was excised and cranioplasty was done. Histopathology came out to be hemangioma.

The meningiomas that originate in the skull have been referred to as calvarial, intraosseous, or intradiploic. Winkler, in 1904, first described a meningioma originating in an extradural location. There are several reported cases of intraosseous intradiploic meningiomas although it is a rare entity. These tumors are mainly osteoblastic and few cases reported are osteolytic in nature. Of these, osteolytic type is more malignant than osteoblastic type of meningioma.^[9]

In our study, we encountered a 53-year-old female patient referred a slowly growing painless lump in the right frontal region for 1 year and 6 months with a history of trauma. On excision, histopathological report was intradiploic meningioma [Figure 3].

Langerhans cell histiocytosis (LCH) is a rare condition that usually affects the pediatric population with an estimated annual incidence between 1 and 7 cases per million of the population. The incidence of LCH is greater in males than in females, with a male-to-female ratio of 2:1. LCH usually affects patients under 10 years of age, patients mainly present with localized bone pain, dyspnea, and malaise, and 75% have nondisseminated disease. The skull, femur, pelvis, and ribs are most commonly involved. With skull lesions, the orbit and the cranial base are frequently involved.^[10]

In our study, we encountered an 8-year-old male with a history of head trauma 2 years back presenting with the left frontal painless swelling for 6 months. The lesion was involving the left frontal bone and roof of left orbit. The lesion was excised. Histopathological report suggested histiocytosis X. The patient was referred to the hematology department for further evaluation and treatment.

Reid *et al.*^[11] reported the first case of calvarial tuberculosis in 1842. Before the advent of effective chemotherapy,

Figure 2: Metastatic lesion involving occipital bone

Figure 3: Intradiploic meningioma

Figure 4: Calvarial tuberculosis

calvarial tuberculosis was estimated to represent 0.2%–1.3% of all cases of skeletal tuberculosis. About 50% of the cases reported in the literature were in patients younger than 10 years, and 70%–90% were younger than 20 years. Increased vascularity, decreased resistance, and unmasking of a latent infection secondary to trauma are thought to be the predisposing factors. In most cases discussed in the literature, patients present with painless scalp swelling and at times a discharging sinus in the scalp.^[1]

In our study, we encountered a 50-year-old female with no predisposing factors presenting with a painful swelling in the left temporal region with no evidence of discharging sinus. The lesion was excised and histopathology report suggested tuberculosis. The patient was started on antitubercular drugs [Figure 4].

Osteomyelitis was first described in 1852 by a French surgeon, Edouard Chassaignac. Pott, in 1778, postulated that osteomyelitis of the skull was caused by a bone contusion and extradural hemorrhage. However, it was not until 100 years later that Van Launelongue classified cases into two types, primary or hematogenous osteomyelitis and secondary or contiguous osteomyelitis. Etiology may result from trauma, bone surgery, bacteremia, or a contiguous infectious focus and is further influenced by diseases that affect the vascularity of bone, as well as by systemic diseases that produce an alteration of host defenses.^[12]

In our study, we encountered a 62-year-old diabetic male patient with painful swelling in the left frontal region with discharging sinus for 3 months. The lesion was excised and culture report was suggestive of *Staphylococcus aureus*. Postoperative antibiotics were given in accordance to the culture report.

In our study, we encountered a 15-year-old male patient with a history of painless swelling in the

Figure 5: Benign cystic lesion of the skull

right frontoparietal region for 2 years. On excision, the histopathological report suggested benign cystic swelling. We did not find a similar case during an extensive search of articles [Figure 5].

CONCLUSION

Osteolytic skull lesions may be the presentation of many pathologies which may vary from infective, benign and malignant tumors. Proper clinical evaluation of the patient must be done including systemic examination for optimum treatment.

In the present study, patients with age more than 40 years had malignant lesions and infective lesions.

Surgical procedure was planned so that complete excision of the lesion must be done whenever possible. Further management of the patient mainly depends on the histopathological diagnosis.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Raut AA, Nagar AM, Muzumdar D, Chawla AJ, Narlawar RS, Fattepurkar S, et al. Imaging features of calvarial tuberculosis: A study of 42 cases. *AJNR Am J Neuroradiol* 2004;25:409-14.
2. Anwar M, Rizwan, Akmal M, Mehmood K. Osteolytic skull lesions rare but important pathology in neurosurgical department. *Pak J Neurol Surg* 2014;18:1.
3. Rubin G, Scienza R, Pasqualin A, Rosta L, Da Pian R. Craniocerebral epidermoids and dermoids. A review of 44 cases. *Acta Neurochir (Wien)* 1989;97:1-16.
4. Ko SJ, Park KJ, Park DH, Kang SH. Extradural dermoid cyst located in the lateral sphenoid ridge. *Brain Tumor Res Treat* 2014;2:39-42.
5. Stark AM, Eichmann T, Mehdorn HM. Skull metastases: Clinical

- features, differential diagnosis, and review of the literature. *Surg Neurol* 2003;60:219-25.
6. Houra K, Vucic M, Cupic H, Kovac D, Cerina V. Osteolytic skull metastasis with dural involvement from a papillary thyroid carcinoma. *Coll Antropol* 2011;35:919-22.
 7. Heckl S, Aschoff A, Kunze S. Cavernomas of the skull: Review of the literature 1975-2000. *Neurosurg Rev* 2002;25:56-62.
 8. Park BH, Hwang E, Kim CH. Primary intraosseous hemangioma in the frontal bone. *Arch Plast Surg* 2013;40:283-5.
 9. Monteiro JT, Baptista AE, Trabulo AS, Almeida C. Intradiploic meningioma of the skull: Case report and review of the literature. *Neurocirugia* 1996;7:129-32.
 10. Chiong C, Jayachandra S, D Eslick G, Al-Khawaja D, Casikar V. A rare case of langerhans cell histiocytosis of the skull in an adult: A systematic review. *Rare Tumors* 2013;5:e38.
 11. Reid E. *Medizinisches Correspondenzblatt Bayerischer Erlangen*. 33;1842.
 12. Pincus DJ, Armstrong MB, Thaller SR. Osteomyelitis of the craniofacial skeleton. *Semin Plast Surg* 2009;23:73-9.