Letters to the Editor

CD4 Count, Magnetic Resonance-white Matter Signals, and Magnetic Resonance Spectroscopy in Human Immunodeficiency Virus Patients

Sir,

Human deficiency virus (HIV) causes acquired immune deficiency syndrome (AIDS). The immune system of the body is vulnerable to attack by this virus and leads to multiple focal and diffuse lesions. Reduction in CD4 cells is seen. CD4 cells are type of immune cells, and they are also called “T-cells” or “helper cells.” Centers for Disease Control and Prevention define a person having AIDS if they have a CD4 count below 200, with or without any opportunistic infections or cancers.^[1] Patients were mainly from School of tropical Medicine the largest tertiary center in eastern India, and from Medical College and Hospitals. Altogether 32 patients who were seropositive for HIV were examined in our center. Detailed history covering all relevant points including CD4 count and CD4% was recorded. Magnetic resonance (MR) spectroscopy study was done in all patients putting the voxel in white matter area. The study was done at 1.5 Tesla GE Signa, and the results were compiled and data analyzed.

The data were accumulated in a tabular form as shown in Table 1 and Table 2. The majority of the patients were males about, and they were about 67.75% of the total population. Most of the patients were in the age group of 20–40 years and they comprised 93.75% of our study group. Twenty-four patients had CD4 count <200 cells/mm³. Eight patients had CD4 count above 200 cells/mm³. CD4% below 5% was seen in 9 patients comprising about 28.0%. CD4% between 5% and 10% was seen in 10 patients comprising about 31.25%. Six patients had CD4% between 20% and 40%. White matter changes were seen in 18 patients comprising about 56% of the study population as shown in Figure 1. MR Spectroscopy showing choline peaks which was seen in 16 patients [voxel placed as seen in Figures 2 and 3 is showing choline peaks] who were having white matter changes and comprised about 89% of them. The finding was that absolute risk of having white matter lesions in AIDs patient was 56.25%. Absolute Risk of choline peaks in patients with white matter lesions was 89.20%. Absolute risk of choline peaks in patients with CD4 Count below 200 was 58.33%.

HIV infection leads to multiple and diffuse lesions in the brain. Noninfective and nonmitotic focal lesions can also be small or diffuse white matter changes. CD4 cells are type of white cells that fight infection. CD4 cells move throughout the body destroying germ cells such as bacteria

Table 1: Number of persons with CD4 count less and more than 200/mm³ and with choline peaks

CD4 count	Choline peaks		Total number of persons
	Yes	No	
<200/mm ³	14	10	24
>200/mm ³	2	4	8
No of persons ¹	16	16	32

Table 2: The percentage and absolute risk in different group

	No of persons and value
Gender	
Male	22
Female	10
Age groups (years)	
20-40	30
Above 40	2
CD4 count	
<100	20
100-200	4
200-300	4
Above 300	6
CD4%	
Below 5	9
5-10	10
10-20	6
20-40	7
White matter changes	
18 out of 32 patients	-
MR spectroscopy showing choline peaks	
No of patient out of 18 patients having white matter changes	16
Absolute risk of white matter lesions (%)	56.25
Absolute risk of choline peaks in patients with white matter lesions (%)	88.90
Absolute risk of choline peaks in patients with CD4 count below 200 (%)	58.33
MR: Magnetic resonance	

and viruses. A normal CD4 count is from 500 to 1500 cells cubic millimeter. Some patients can have CD4 countless but not so severe infections while the reverse can also be seen. CD4% or the rate of its change is a better predictor of clinical progression. Although the CD4+ absolute number that identifies a specific level of immune suppression changes with age,^[2] the CD4+ percentage that defines each immunologic category does not. Thus, a change in CD4+ percentage, not number, may be a better marker to

Figure 1: T2 fluid-attenuated inversion recovery images are showing small white matter hyperintensities in the right subcortical parietal region and in periventricular region

Figure 2: Magnetic resonance spectroscopy done in posterior periventricular lesion

Figure 3: Magnetic resonance spectroscopy showed increase in choline peaks at that level

identify disease progression.^[3] HIV infects and destroys CD + T Lymphocytes.^[4]

HIV encephalopathy leads to myelin pallor leading to hyperintense signal changes as seen in T2 weighted imaging.^[5] These lesions do not enhance. They do not have any mass effect. Increase in these lesions is seen as the patient condition deteriorates or CD4 or CD4% decreases. MR Spectroscopy is showing increase in choline peaks^[6] and choline/creatine ratio.^[7,8] Choline which resonates at 3.2 ppm is basically phosphatidylcholine and is a membrane constituent.^[4] and is a marker of white matter lesions.^[9,10] The study shows a correlation between CD4 count, white matter changes and choline peaks in MR Spectroscopy. Choline peaks are seen in about 89% of patients with white matter changes, which is a significant value. White matter changes are there CD4 count and CD% decreases. Significant increases in the choline/creatine ratios were seen in patients with low CD4 count.^[9] The advantage of this study is that it will help in monitoring the treatment and cliniconeurological and radiological findings in HIV patient. By having CD4 count possibility of white matter changes and its associated importance can be known.

Acknowledgment

We would like to thank all the staff that has greatly supported me in this project.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

M. A. Hashmi, Shyam Baboo¹, Bibhuti Saha²

Medical College and Hospitals Campus, MRI Section,

¹Department of Radiology, SKMCH, Muzaffarpur, Bihar,

²Department of Medicine, School of Tropical Medicine, Kolkata, West Bengal, India

Address for correspondence: Dr. M. A. Hashmi, Mri-Section, EKO Diagnostics Medical College and Hospitals Campus, 88 College St., Kolkata - 700 073, West Bengal, India. E-mail: ahashmidradd@yahoo.co.in

REFERENCES

1. Yu LM, Philippa J. Easterbrook, tom marshall. *Int J Epidemiol* 1997;26:1367-72.
2. Khilnani P, Rajdev SK, Uttam R. Pediatric aids – Part 1. *Indian J Crit Care Med* 2003;7:257-72.
3. Kannangai R, Kandathil AJ, Ebenezer DL, Nithyanandam G, Samuel P, Abraham OC, *et al.* Evidence for lower CD4+T cell and higher viral load in asymptomatic HIV-1 infected individuals of India: Implications for therapy initiation. *Indian J Med Microbiol* 2008;26:217-21.
4. Elston JW, Thaker HK. Co-infection with human immunodeficiency virus and tuberculosis. *Indian J Dermatol Venerol Leprol* 2008;74:194-9.
5. Hurley RA, Ernst T, Khalili K, Del Valle L, Simone IL, Taber KH, *et al.* Identification of HIV-associated progressive

- multifocal leukoencephalopathy: Magnetic resonance imaging and spectroscopy. *J Neuropsychiatry Clin Neurosci* 2003;15:1-6.
6. Jarvik JG, Lenkinski RE, Grossman RI, Gomori JM, Schnall MD, Frank I, *et al.* Proton MR spectroscopy of HIV-infected patients: Characterization of abnormalities with imaging and clinical correlation. *Radiology* 1993;186:739-44.
 7. Simone IL, Federico F, Tortorella C, Andreola CF, Zimatore GB, Giannini P, *et al.* Localised 1H-MR spectroscopy for metabolic characterisation of diffuse and focal brain lesions in patients infected with HIV. *J Neurol Neurosurg Psychiatry* 1998;64:516-23.
 8. Offiah CE, Turnbull IW. The imaging appearances of intracranial CNS infections in adult HIV and AIDS patients. *Clin Radiol* 2006;61:393-401.
 9. Chong WK, Sweeney B, Wilkinson ID, Paley M, Hall-Craggs MA, Kendall BE, *et al.* Proton spectroscopy of the brain in HIV infection: Correlation with clinical, immunologic, and MR imaging findings. *Radiology* 1993;188:119-24.
 10. Paul RH, Yiannoutsos CT, Miller EN, Chang L, Marra CM, Schifitto G, *et al.* Proton MRS and neuropsychological correlates in AIDS dementia complex: Evidence of subcortical specificity. *J Neuropsychiatry Clin Neurosci* 2007;19:283-92.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/jnnp.jnnp_33_18

How to cite this article: Hashmi MA, Baboo S, Saha B. CD4 count, magnetic resonance-white matter signals, and magnetic resonance spectroscopy in human immunodeficiency virus patients. *J Neurosci Rural Pract* 2018;9:437-9.

© 2018 Journal of Neurosciences in Rural Practice | Published by Wolters Kluwer - Medknow