Case Series

Surgically Treated Primary Spinal Cord Neoplasms in Southeastern Nigeria

Mark Chukwunweike Chikani, Okwuoma Okwunodulu¹, Mathew Mesi, Wilfred C. Mezue, Samuel C. Ohaegbulam¹, Chika C. Ndubuisi¹

Department of Surgery,
Division of Neurosurgery,
University of Nigeria
Teaching Hospital,
¹Department of Surgery,
Memfy's Hospital for
Neurosurgery, Enugu, Nigeria

ABSTRACT

Background: Primary spinal cord tumors are not as frequently encountered as their cranial counterparts. They could present in such an indolent manner that requires a reasonable index of suspicion for their diagnosis to be considered. **Objective:** The objective of this study is to analyze the incidence and pattern of primary spinal cord and appendage neoplasms in patients surgically treated in our institution over a decade of practice. **Materials and Methods:** A retrospective review of clinical, radiological, and histopathology profiles of patients surgically treated for primary spinal cord tumor from 2006 to 2016 was carried out. Retrieved data were analyzed using SPSS version 21. **Results:** Out of 472 spine procedures were performed within the study period 17 (3.6%) cases of histologically proven primary spinal cord tumors were identified. The age of patients ranged between 17 and 77 years with a mean age was 45 years. The male: female ratio was 1:1.1. Motor deficit and pain were the most common presenting symptoms seen in 35.3% and 29.4% of patients, respectively. Meningiomas are the most common histological diagnosis (70.6%), distantly followed by Schwannoma (17.6%). The most common location of the tumors was intradural extramedullary (70.6%). All patients had gross total resection of tumor with no perioperative mortality. **Conclusion:** Meningioma is the most common surgically treated primary spinal cord tumor in our setting. Surgery is associated with good outcome.

KEYWORDS: Primary, spinal cord, surgery, tumors

INTRODUCTION

Since Sir Victor Horsley performed the first successful resection of a spinal intradural extramedullary (IDEM) tumor by means of a laminectomy in 1887, surgical management of these lesions has continued to evolve over time.^[1] Primary spinal cord tumors are 10 times less common than their cranial counterparts, although they are histopathologically similar.^[2] Based on their location with respect to the dural sac and spinal cord, the tumors can be classified into extradural; IDEM; or intramedullary with most primary tumors located in the IDEM area.^[3] The presentation could be nonspecific, but pain and motor dysfunction are important symptoms of spinal cord tumors.^[1,3-5] Therefore, most of the patients are wrongly diagnosed with cervical spondylopathy or intervertebral disk herniation.^[6]

Magnetic resonance imaging (MRI) is currently the gold standard for imaging as it provides accurate diagnoses and delineates the soft-tissue components.^[5] Intraspinous tumors are relatively uncommon lesions but can cause significant morbidity and can be associated with mortality as well.^[7] Complete surgical extirpation is the goal where feasible. The use of intraoperative monitoring has been shown to improve the extent of tumor resection with increased safety^[8] particularly for IDEM. Prognosis for patient longevity, which can be determined by the patient's age at diagnosis and the extent of the disease,

Address for correspondence: Dr. Mark Chukwunweike Chikani, Department of Surgery, Neurosurgery Unit, University of Nigeria Teaching Hospital, Enugu, Nigeria.
E-mail: markochikani@yahoo.com

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Chikani MC, Okwunodulu O, Mesi M, Mezue WC, Ohaegbulam SC, Ndubuisi CC. Surgically treated primary spinal cord neoplasms in Southeastern Nigeria. J Neurosci Rural Pract 2018;9:137-9.

Access this article online

Quick Response Code:

Website:
www.ruralneuropractice.com

DOI:
10.4103/jnpr.jnpr_391_17

is the major factor in deciding which surgical approach would prove suitable.^[9]

No previous or similar work has been done in this region on surgically treated primary spinal cord neoplasm and is imperative to understand the behavior of these tumors for an appropriate, well-timed intervention to be made.

MATERIALS AND METHODS

This is a descriptive cross-sectional retrospective analysis of patients with primary spinal cord and appendage neoplasms surgically managed in our institution from 2006 to October 2016. All patients had diagnostic preoperative MRI and surgical intervention. Data on demographic variables, clinical features, surgical approach, pathology, location, and association with syrinx were extracted. Primary cord tumors were classified according to anatomical location.

Retrieved data were analyzed using SPSS for windows version 21 (IBM). $P < 0.05$ was considered statistically significant.

Tumors without histological diagnosis were excluded from the analysis.

RESULTS

Out of 472 spine procedures performed within the study, 17 cases of histologically proven primary spinal cord tumors were identified. This represents 3.6% of spine all procedures. The age of patients ranged between 17 and 77 years with a mean age was 45 years. The disease peaks at the fourth decade of life [Table 1]. The male:female ratio was 1:1.1. Nine patients (52.9%) presented within 1 year of onset of symptoms. The mean duration of symptoms before presentation was 3.4 years.

About 94% of patients, the symptoms were insidious in onset, while the remaining 6% of patients had a sudden onset. Motor deficit and pain were the most common presenting symptoms seen in 35.3% and 29.4% of patients, respectively [Figure 1].

Tumor predominantly affects the cervical region [Figure 2]. The IDEM area is the site of tumor

location in 12 patients (70.6%). Meningiomas are the most common histological diagnosis (70.6%), and it has a female preponderance male:female of 1:2 [Figure 3 and Table 2]. The ratio of benign to malignant lesions is 16:1. No histologically confirmed intramedullary astrocytoma/ependymomas were documented in our cohort to our utmost surprise.

Laminectomy and gross total resection (GTR) was done in 53% of cases, hemilaminectomy + GTR in 35%, and subligamentous laminectomy + GTR in 12% of patients. All patients were discharged home. No mortality was noted.

DISCUSSION

Primary spine tumor surgeries are not particularly common in our environment as it accounts for only 3.6% of spine procedures. However, the fact that it peaks at the fifth decade and affected cohorts with a mean age of 45

Figure 1: Distribution of the clinical features (n=17)

Figure 2: Region of the spine affected (n=17)

Figure 3: Tumor location by histology (n=17)

Table 1: Age distribution of patients

Age (years)	Frequency (%)
11-20	2 (11.8)
21-30	1 (5.9)
31-40	5 (29.4)
41-50	3 (17.6)
51-60	3 (17.6)
61-70	2 (11.8)
71-80	1 (5.9)
Total	17 (100)

Table 2: Distribution of spinal meningioma

	Frequency (%)
Sex	
Male	4 (33.3)
Female	8 (66.7)
Location	
Cervical spine	7 (58.3)
Thoracic spine	5 (41.7)

suggests implications on the workforce. Furthermore, the tumors were predominantly IDEM in location. In the work by Moein *et al.* among 102 patients in Iran, they reported a mean age for primary tumors of 40.2 years with most tumors located in the intradural area.^[10] Our findings corroborate with this report. However, whereas meningioma was the most common tumor in our series, they reported nerve sheath tumors. Most of the tumors (94.1%) were benign and exhibit a long latency (time to presentation) averaging 3.4 years among our cohorts.

Adeolu *et al.* found metastasis to be the most common histology followed by meningioma in Ibadan, southwest Nigeria.^[11] This pattern is understandable as we excluded metastasis from our study. When this is applied to their findings, meningioma will be the most common tumor as we also found among our patients. There is a clear female preponderance in meningiomas among our patients with all the lesions distributed in the cervical and thoracic regions only. Our findings were also in line with reports from other literature where female predominance was also reported; though the thoracic spine was the most reported site, it could occur in other regions of the spine.^[12-14] In another small-sized study on 14 patients over a 5-year period, the authors reported the most common IDEM tumors as meningiomas and Schwannomas.^[15] In our series, schwannoma was also the second most common histological diagnosis seen in 3 patients (17.6%), and all were located in the IDEM space. Motor deficits and local pain are important clinical features observed in our patients.

In a review of 35 surgically managed cervical spine primary tumours, Zileli *et al.* concluded that complete tumor resection is the oncologically best surgical strategy and should be attempted whenever possible.^[16] Although different forms of laminectomy were used, we were able to achieve GTR in all our patients. No perioperative mortality was observed.

CONCLUSION

Primary spinal cord tumors commonly affect young- and middle-aged patients. A triad of motor dysfunction, regional pain, and sphincteric dysfunction were the common clinical features. Meningioma is the most

common primary tumor. Overall surgical outcome is good.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Sellin JN, Tatsui CE, Rhines LD. Assessment and treatment of benign intradural extramedullary tumors. In: Winn HR, editor. Youmans & Winn Neurological Surgery. 7th ed. Philadelphia: Elsevier; 2017. p. 2428-34.e2.
- Samartzis D, Gillis CC, Shih P, O'Toole JE, Fessler RG. Intradural spinal cord tumors: Part I-epidemiology, pathophysiology, and diagnosis. *Global Spine J* 2015;5:425-35.
- Quiles AM, Roselló EG, Laguillo G, García R, Caro JL, Pérez F, *et al.* A Comprehensive Review of Intraspinal Tumors : Diagnostic, Classification and Radio-Pathologic Correlation. European Society of Radiology; 2013. p. 1-50. Available from: <http://www.myESR.org>. [Last accessed on 2017 May 22].
- Tobin MK, Geraghty JR, Engelhard HH, Linninger AA, Mehta AI. Intradural spinal cord tumors: A review of current and future treatment strategies. *Neurosurg Focus* 2015;39:E14.
- Bruneau M, Lefranc F, Balériaux D, Brotchi J. Intradural extramedullary and intramedullary spinal cord tumors. In: Ellenbogen RG, Abdulrauf SI, Sekhar LN, editors. Principles of Neurological Surgery. 3rd ed. Philadelphia: Elsevier Saunders; 2012. p. 422-36.e3.
- Govind M, Radheyshyam M, Achal S, Ashok G. Intradural extramedullary spinal cord tumors : A retrospective study at tertiary referral hospital. *Rom Neurosurg* 2016;XXX: 106-12.
- Ravi N, Manjappa BH, Nagaraj BR, Naveen KG, Lakshmeesha MT, Ramesh V, *et al.* MRI evaluation of different spectrum of spinal tumors. *SSRG Int J Med Sci* 2014;1:14-30.
- Ghadirpour R, Nasi D, Iaccarino C, Giraldo D, Sabadini R, Motti L, *et al.* Intraoperative neurophysiological monitoring for intradural extramedullary tumors: Why not? *Clin Neurol Neurosurg* 2015;130:140-9.
- Willey JO, Larkins MV. Editors Surgical removal of spinal cord tumors: SurgTechnol 1996. Available from: www.ast.org/pdf/WSeries/1005.pdf. [Cited 2017 Aug 02].
- Moein P, Behnamfar O, Khalighinejad N, Farajzadegan Z, Fard SA, Razavi M, *et al.* A 12-year epidemiologic study on primary spinal cord tumors in Isfahan, Iran. *J Res Med Sci* 2013;18:17-21.
- Adeolu AA, Oyemolade TA, Salami AA, Adigun TA, Malomo AO, Akang EA, *et al.* Features and outcome of surgical management of spinal tumors in a cohort of Nigerian patients. *World Neurosurg* 2015;84:1090-4.
- Bydon M, Gokaslan ZL. Spinal meningioma resection. *World Neurosurg* 2015;83:1032-3.
- Riad H, Knafo S, Segnarbieux F, Lonjon N. Spinal meningiomas: Surgical outcome and literature review. *Neurochirurgie* 2013;59:30-4.
- Saraceni C, Harrop JS. Spinal meningioma: Chronicles of contemporary neurosurgical diagnosis and management. *Clin Neurol Neurosurg* 2009;111:221-6.
- Arnautovic K, Arnautovic A. Extramedullary intradural spinal tumors: A review of modern diagnostic and treatment options and a report of a series. *Bosn J Basic Med Sci* 2009;9 Suppl 1:40-5.
- Zileli M, Kilinçer C, Ersahin Y, Cagli S. Primary tumors of the cervical spine: A retrospective review of 35 surgically managed cases. *Spine J* 2007;7:165-73.