Case Series

Multiple, Primary Brain Tumors with Diverse Origins and Different Localizations: Case Series and Review of the Literature

Thara Tunthanathip, Kanet Kanjanapradit¹, Sanguansin Ratanalert, Nakornchai Phuenpathom, Thakul Oearsakul, Anukoon Kaewborisutsakul

Department of Surgery,
Neurosurgical Unit, Faculty
of Medicine, Prince
of Songkla University,

¹Department of Pathology,
Faculty of Medicine, Prince
of Songkla University,
Hat Yai, Songkhla, 90110,
Thailand

ABSTRACT

Background: Multiple, primary brain tumors with different histological types occurring in the same patient are extremely rare. Several hypotheses have been proposed, and the pathophysiology of coexisting tumors has long been debated; however, due to low incidence, standard practices for this scenario are still inconclusive. **Case Description:** The authors describe 6 cases of coexisting tumors. By conducting a literature research focused on the computed tomography (CT) era and patients without prior radiation or phakomatoses. Sixty-five such reported cases were identified. In addition, the authors summarize their experience in 6 patients including histopathological features, chronological presentations, outcomes, mortality, and management from their series as well as from previous cases from the reported literature. **Conclusion:** The coexistence of multiple, primary brain tumors is an interesting condition. Surgical management remains the major treatment; malignant histology has a poor prognostic factor.

KEYWORDS: Coexistence of brain tumor, diffuse astrocytoma, glioblastoma, meningioma, multiple primary brain tumor

INTRODUCTION

The association of multiple, primary brain tumors with different histology occurring simultaneously in the same patient is an extremely rare condition. It has been frequently associated with previous radiotherapy or phakomatoses^[1,2] but is still exceptionally rare as a primary observation. Before the era of computed tomographies (CTs), Cushing and Eisenhardt reported, in 1938, a patient who harbored both a glioblastoma and a meningioma.^[3] Only a few cases of this scenario have since been reported. The authors present their experience gained from a series of such primary brain tumors with different histopathology types and in different intracranial localizations. They discuss these cases in the context of the currently available literature.

The most frequently reported primary brain tumor in the Central Brain Tumor Registry of the United States is nonmalignant meningioma (36.1%), followed by glioblastoma (15.4%). In addition, the nonmalignant pituitary adenoma and nerve sheath tumors account for 15.1% and 8% of all tumors, respectively. In

gender-specific distribution, the incidence rate of tumor of the meninges is higher in females (10.51/100,000) rather than in males (4.85/100,000), while the incidence rate of the tumor of the neuroepithelial tissue is higher in males (7.77/100,000) rather than in females (5.61/100,000).^[4] Moreover, the incidental, primary brain tumor can be found in general population. Asymptomatic meningiomas were the most frequent in 0.9%–1% while pituitary adenomas were 0.3% within the population.^[5,6] However, the condition where coexisting, primary brain tumors occur in the same patient is truly scarce.

We reviewed these from the relevant literature in MEDLINE, EMBASE, and Scopus. The selection criteria included those in the English language and

Address for correspondence: Dr. Thara Tunthanathip,

Department of Surgery, Neurosurgical Unit, Faculty of Medicine,
Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.
E-mail: tsus4@hotmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Tunthanathip T, Kanjanapradit K, Ratanalert S, Phuenpathom N, Oearsakul T, Kaewborisutsakul A. Multiple, primary brain tumors with diverse origins and different localizations: Case series and review of the literature. J. Neurosci Rural Pract 2018;9:593-607.

Access this article online

Quick Response Code:

Website:

www.ruralneuropractice.com

DOI:

10.4103/jnpr.jnpr_82_18

illustrated neuroimaging such as CT or magnetic resonance imaging (MRI) brain scans. The authors also do not consider these tumors to be associated with previous radiotherapy (Cahan's criteria) or phakomatosis.^[7] According to the criteria, 65 cases were reported in the literature and the authors added 6 cases in Appendix 1.^[8-39]

CASE SERIES

Patient 1: Meningioma and diffuse astrocytoma

A 68-year-old, right-handed woman suffering from dementia on the right arm coupled with suffering leg weakness in October 2013. A clinical examination showed right hemiparesis and right facial nerve palsy with upper motor neuron type. She had neither café-au-lait spots nor Lisch nodules, which are found in neurofibromatosis Type 1 (NF1). There was also an absence of first-degree relatives of NF1.

MRI of the brain showed a homogeneously enhancing mass with dural tail sign located in the left frontal lobe as seen on T1-weighted (T1W) gadolinium-enhanced images. Furthermore, another lesion was observed at the right insular lobe that showed an ill-defined area of mixed hypo- and hyperintense but nonenhancing lesion on T1W, fluid-attenuated inversion recovery, and T1W gadolinium-enhanced images, respectively [Figure 1a]. In addition, the T1-weight inversion recovery MRI showed hyperintensity of the right insular lesion [Figure 1b].

A left craniotomy for tumor removal was performed to address the frontal lesion first. The tumor was found attached to the dura mater and measuring 4.5 cm in diameter by 4.5 cm. Both tumor and attached dura mater were completely removed. After surgery, this patient made an uneventful recovery without any new neurological impairment. The pathology of the frontal lesion was fibrous meningioma (WHO II) [Figure 1d].

One year after the first surgery, the patient suffered a generalized tonic-clonic seizure. A follow-up MRI brain scan showed no evidence of recurrence of the left frontal meningioma, and there was a stable size of the right insular lesion. In addition, no decreased N-acetylaspartate (NAA) and no choline peak were noted in magnetic resonance spectroscopy [Figure 1c]. Taking the clinical seizure as a sign of progression, we had the indications that an operation was required to obtain a diagnosis. A right-sided craniotomy for a transsylvian approach was performed. Discoloration of the insular cortex was encountered, and the extent of resection was 90% in the complex location. Thus, surgical specimens were sent for analysis.

Pathological examination of the specimens found hypercellularity of glial cells without endothelial

proliferation and necrosis. Furthermore, the specimen stained positive for glial fibrillary acidic protein (GFAP), which indicated a glial cell tumor. Due to the lack of diagnostic molecular testing, a final diagnosis of tumor was designated as those not otherwise specified diffuse astrocytoma (WHO II) according to the 2016 WHO classification [Figure 1 e and f]. After surgery, this patient again made an uneventful recovery without any new neurological impairment. At follow-up, she maintained independent daily activities.

Chromosome analysis

Chromosomal analysis from peripheral blood cells of the patient was performed for screening germline mutation. The karyotype was 44, XX and an absence of chromosome 17p deletions.

Patient 2: Glioblastoma multiforme with pituitary adenoma

A 63-year-old male patient was referred to our hospital due to an alteration in his consciousness. Two weeks prior, he had suffered from a change in mentation and left hemiparesis, so the patient had been admitted to a provincial hospital. CT head scans demonstrated two sites with intracranial tumors: an irregular rim-enhancing mass at the right frontoparietal lobe and a sellar tumor. During hospitalization, the patient showed progressive neurological deterioration. Therefore, the patient underwent a decompressive craniectomy and was transferred to our hospital. At the tertiary hospital, he was fully oriented but had hemiplegia on the left side. Cranial T1W MRI with intravenous (I/V) gadolinium revealed an irregular, heterogeneously contrast-enhancing mass at the right frontoparietal lobe measuring 52 mm × 68 mm × 54 mm with central hypointensity indicative of necrosis [Figure 2a].

In addition, another tumor was noted in the sellar region with suprasellar extension, measuring 20 mm × 21 mm × 29 mm [Figure 2d and e]. Pituitary hormones were found to be within normal values. An extended right frontoparietal craniotomy was performed to approach both tumors. However, the patients' position was suboptimal to access the sellar region limiting the resection to a partial one at this site.

Histopathology of the right frontoparietal lesion revealed a glioblastoma multiforme [Figure 2b and c] while the sellar tumor was identified as pituitary adenoma [Figure 2f]. Since his hormonal profiles were normal, nonfunctioning pituitary adenoma was designated. The postoperative period was uneventful, and the patient was sent for adjuvant therapy. Unfortunately, the patient refused adjuvant therapy and was lost to for any further follow-up.

Figure 1: Case 1 – Magnetic resonance imaging showing an enhanced mass with a dural tail sign at the left frontal convexity (a). T1-weight inversion recovery magnetic resonance imaging showing hyperintense lesion at the right insular lobe (b). Magnetic resonance spectroscopy showing decreased N-acetylaspartate without elevated choline at right temporal lesion (c). Histological findings of a fibrous meningioma demonstrating a proliferation of generally spindled cells (d). Histological findings of diffused astrocytoma demonstrating increased cellular density of glial cell without necrosis (e). Immunoreactivity of astrocytoma was positive for glial fibrillary acidic protein (f)

Figure 2: Case 2 – Magnetic resonance imaging showing an enhanced mass with central necrosis at the right frontoparietal lobe (a). Palisading tumor necrosis was observed (b). The neoplastic glial cells have nuclear enlargement with hyperchromatic nuclei, irregular nuclear membrane, and abnormal endothelial cell proliferation (c). Magnetic resonance imaging showing enhanced suprasellar mass (d). Sagittal magnetic resonance imaging showing enhanced sellar-suprasellar mass compressed chiasm (e). Pituitary tumor showing proliferation of homogeneous round cell with sheeting pattern. The tumor cells have eosinophilic granular cytoplasm with round nucleus (f)

Patient 3: Planum sphenoidale meningioma and pituitary adenoma

A 62-year-old female patient presented with progressively blurred vision in her right eye, which had occurred over the course of 1 year. On examination, she had normal visual acuity in her left eye but only 20/200 in her right eye. A cranial T1W MRI with I/V gadolinium revealed a homogeneously enhancing, board-based tumor at the planum sphenoidale with a dural tail sign, measuring 37 mm × 30 mm × 18 mm. Interestingly, a second

enhancing sellar tumor with suprasellar extension was observed posterior to the former [Figure 3a and b]. Therefore, the patient underwent a subfrontal approach for microsurgical resection. The anterior tumor showed a good arachnoidal plane for dissection but was attached to the dura of the planum sphenoidale. This tumor was subtotally removed due to high vascularity.

Histological characteristics were those of a meningothelial meningioma. Tumor cells formed lobules where they appeared to form a syncytium with oval

nuclei, occasionally showing central clearing [Figure 3c]. The tumor was immunostained and found to be positive for epithelial membrane antigen. After surgery, the patient developed Gram-negative septicemia and required antibiotic administration for 2 weeks. She recovered well and her vision, in the right eye, gradually improved too. At follow-up, she maintained independent daily activities. She does not wish to pursue a second surgery for the tumor residual.

Patient 4: Intraosseous meningioma and splenium anaplastic astrocytoma

A 61-year-old female patient presented with progressive headaches. Ten years prior, she had been diagnosed with an intraosseous meningioma at the right frontal base [Figure 4a and b] for which she underwent a right-sided craniotomy for resection. Follow-ups with MRIs were obtained annually and showed stable residual disease. However, over the course of 3 months, before admission, she had begun to develop progressively worse headaches. After a repeat of an MRI, a newly infiltrative heterogeneously enhancing mass involving bilateral splenium of the corpus callosum was observed. To obtain a pathological diagnosis, an image-guided biopsy was carried out [Figure 4c and d].

Microscopic examination revealed hyperchromatic nuclei and mitotic figures. However, no areas of necrosis, hemorrhage, or endothelial hyperplasia were observed. A diagnosis of anaplastic astrocytoma WHO Grade III was made [Figure 4e]. Her hospital course was uneventful. Later, she was given a course of adjuvant external beam radiotherapy. She did not receive chemotherapy due to financial reasons. She has been followed up since, and she was still able to manage daily activities at the last follow-up.

Patient 5: Pituitary adenoma with multiple meningiomas

A 50-year-old female patient presented with a chronic headache for 3 months and with progressively worsening visual field defects. On examination, a dense bilateral temporal hemianopia with normal visual acuity was detected. Her hormonal laboratory demonstrated mildly elevated prolactin (64.5 ng/ml) although all other hormones were within normal range. Cranial MRI revealed a heterogeneously enhancing sellar tumor with suprasellar extension and chiasma compression [Figure 5a]. Two other hemispheric lesions were identified, which imposed as homogeneously enhancing solid nodules (1.1 cm on the right posterior parietal lobe and 1 cm on the left vertex with hyperostosis of adjusted skull). These were interpreted as likely meningiomas [Figure 5c and d]. The patient underwent an endoscopic endonasal transsphenoidal approach for successful sellar tumor removal.

Microscopic examination of this surgical specimen showed epithelioid cells with a sheet-like growth pattern. A histological diagnosis of pituitary adenoma was made [Figure 5b]. The postoperative period was without incident. The patient is being followed clinically and expectantly for her convexity lesions. Her visual field defects have improved since.

Patient 6: Pineal epidermoid cyst with vestibular schwannoma

A 37-year-old female was referred from a provincial hospital due to a rapidly, worsening headache over a period of 2 weeks. According to her history, she had been suffering from progressive headaches for 5 years as well as from hearing disturbances on the right side for 1 year. Two weeks before admission, her headaches got rapidly more severe and she developed a bilateral papilledema.

Figure 3: Case 3 – Sagittal magnetic resonance imaging showing enhanced extra-axial mass at planum sphenoidale and the mass had more extension into the suprasellar region. The pituitary gland was enlarged and heterogeneous enhancement (a). Axial magnetic resonance imaging showing the pituitary tumor is left posteriorly displaced from the planum sphenoidale tumor (b). Histological photomicrograph of the planum sphenoidale tumor showing proliferation of meningothelial cells that arrange in a whirling pattern. The tumor cells have an elongated shape with a homogeneous round nucleus (c)

Figure 4: Case 4 – Bone window axial computed tomography brain showing 1.7 cm thickening of bone involving the right sphenoid bone and superior and lateral orbital wall (a). Magnetic resonance imaging showing enhanced en plaque soft tissue and dural enhancement in the right middle cranial fossa. Severe right exophthalmos was observed (b). Magnetic resonance imaging showing 3.3 cm × 4.6 cm heterogeneously enhancing mass involving bilateral splenium of corpus callosum (c and d). Histological photomicrograph of splenium anaplastic astrocytoma showing aggregation of oval-to-elongated shape glial cell with and dysplastic nuclei (e)

Cranial MRI revealed two tumors located in the pineal region and at the cerebellopontine angle (CPA), causing obstructive hydrocephalus.

A cranial T1W MRI revealed a nonenhancing large tumor at the pineal region with third ventricle extension. This tumor displayed restricted diffusion on diffusion-weighted MRI [Figure 6a and b]. The second lesion imposed as a homogeneously enhancing cone-shaped tumor at CPA at the level of the right internal acoustic canal. The findings were compatible with vestibular schwannoma [Figure 6d].

The patient first underwent ventriculoperitoneal shunting for diversion of her cerebrospinal fluid. Later, an occipital transtentorial approach was chosen for pineal tumor removal. Near total tumor removal was achieved and the postoperative period was uneventful.

Histologically, the microscopic examination showed epithelium and lamellated keratin [Figure 6c], and the final diagnosis was epidermoid tumor. At last follow-up, she had recovered well and decided to postpone any CPA tumor surgery.

DISCUSSION

In this case series, the authors report these six cases of coexisting, primary brain tumors observed in individual patients. They then reviewed the available literature for similar reported cases of coexistence between various tumor entities.

Figure 5: Case 5 – Axial magnetic resonance imaging showing a heterogeneous enhanced pituitary tumor with suprasellar extension (a). Histological photomicrograph of pituitary tumor showing proliferation of round-to-oval cell in a sheeting pattern. The tumor cells have homogeneous round nucleus fine granular chromatin (b). Two extra-axial enhanced nodules at the left vertex and posterior parietal lobe (c and d)

Clinical features

The clinical characteristics of the patient cohort are summarized in Table 1. In addition, the summary table of all cases that have been reportedly been performed is in Appendix 1. This rare scenario is more frequently observed in the fourth-to-sixth decades of life, and we found a female predominance. The most frequently encountered combination is meningioma and pituitary adenoma.

The majority of coexisting tumors occurred simultaneously and in distinct locations. Almost a half of the tumors found under these circumstances were meningiomas. In cases of pituitary lesions, they were nonfunctioning adenomas.

Other associations with meningioma were described as glioma, primary cerebral lymphoma, schwannoma, and craniopharyngioma. In cases in which gliomas were found in this condition these were usually high grade glioma. Interestingly, an unusual combination between an epidermoid cyst and a schwannoma was found in our own series.

From pooled analysis, the mortality rate has been reported to be 21.1% (15/71). Mean survival is 10.36 months; however, the reason of death was restricted from secondary citation. Notice, 86.6% (13/15) of lethal cases are a combination between glioma and meningioma and 66.7% (10/15) are malignant glioma.

Pathogenesis

Various hypothesis of an association between the “tumor pairs” has been advocated in the past. Local tissue irritation from perilesional edema caused by

Figure 6: Case 6 – Sagittal magnetic resonance imaging revealing a lobulated extra-axial cystic lesion about 6.3 cm × 6.6 cm × 4.2 cm located at suprapineal recess and superior cerebellar recess. The lesion insinuates into posterior aspect of the third ventricle (a). Diffusion-weighted imaging showing restricted diffusion pineal lesion (b). The tumor shows proliferation of benign squamous epithelium with keratin material (c). Extra-axial heterogeneous enhancing mass is located at the right cerebellopontine angle with right internal acoustic canal invasion. Widened right internal auditory canal is observed (d)

the first tumor has been considered as a factor in inducing astrocyte or arachnoid cell transformation, causing neoplastic proliferation in the future.^[16,33] However, this hypothesis cannot completely explain the observations made since more than half of the simultaneously existing tumors are topographically disconnected (spatially remote) lesions. Therefore, another hypotheses have been proposed where a common genetic pathway is suggested, including the p53 disruption and receptor tyrosine kinase signaling molecules that have an expression of platelet-derived growth factor receptors (PDGFRs).^[34]

PDGF is one of the numerous growth factors that appears to play a significant role in angiogenesis and that can lead to tumorigenesis.^[40,41] Suzuki *et al.* reported an association of glioblastoma simultaneously presenting with meningioma. In their study, both tumors were strongly positive for PDGFR expression levels as seen through immunohistochemistry.^[34] PDGFR subunits were classified in two types: alpha- and beta-subunits.

The alpha-subunit of the human PDGFR-alpha (PDGFRA) gene maps on q11-q12 of the human chromosome 4^[42] and gene encoding the beta-subunit of the human PDGFR-beta (PDGFRB) maps on human chromosome 5 at region 5q31-32.^[43] Theoretically, coexisting tumors should therefore carry abnormalities of chromosome 4 or 5, which PDGFRA and PDGFRB are transcribed.^[40-43]

Nevertheless, relevant studies did not demonstrate concordant evidence. Nestler *et al.* undertook

Table 1: Characteristics of coexisting glioma and meningioma in the computed tomography era (n=71)

Characteristics	n (%)
Sex (n=68)	
Male	24 (35.3)
Female	43 (64.7)
Mean age-year (SD)	
Mean	55.8
SD	12.2
Range	33-82
Coexisting tumors (n=142)*	
Meningioma	67 (46.5)
Glioma	30 (20.8)
Pituitary adenoma	30 (20.8)
Schwannoma	6 (4.2)
Lymphoma	6 (4.2)
Craniopharyngioma	3 (2.1)
Epidermoid	1 (0.7)
Chordoma	1 (0.7)
Chronological presentation (n=71)	
Simultaneous	59 (83.1)
Sequential	12 (16.9)
Interval of second tumor presentation-month (n=12)	
Median	29
Median	50.58
SD	56.17
Range	1-156
Location of both tumors (n=71)	
Adjacent	34 (47.9)
Separate	37 (52.1)
Meningioma	
Coexisting tumor with meningioma (n=67)	
Pituitary adenoma	27 (40.3)
Glioma	26 (38.8)
Other	13 (19.4)
Schwannoma	5 (7.5)
Lymphoma	6 (8.9)
Craniopharyngioma	3 (4.5)
Location of meningioma (n=67)	
Frontal	14 (20.9)
Parasagittal/flax	13 (19.7)
Parietal	9 (13.4)
Sphenoid	8 (12.1)
Sellar/tuber	7 (10.4)
Temporal	4 (6.0)
Tentorium cerebelli	2 (3.0)
Planum sphenoidale	2 (3.0)
Intraventricular	2 (3.0)
Frontoparietotemporal (convexity)	3 (4.5)
Occipital	1 (1.5)
Cerebellopontine angle	1 (1.5)
Cerebellum	1 (1.5)
Site (n=66)	

Contd...

Table 1: Contd...

Characteristics	n (%)
Right	26 (38.8)
Left	25 (37.3)
Midline	15 (22.4)
Both sides	1 (1.5)
Type of meningioma (n=32)	
Meningothelial	14 (43.8)
Fibrous	7 (21.9)
Transitional	5 (15.6)
Psammomatous	5 (15.6)
Angioblastic	1 (3.1)
Glioma	
Coexisting tumor of glioma (n=28)	
Meningioma	26 (92.9)
Pituitary adenoma	2 (7.1)
Location (n=30)*	
Frontal	9 (30.0)
Parietal	6 (20.0)
Occipital	1 (3.3)
Temporal	10 (33.4)
Corpus callosum	1 (3.3)
Brainstem*	2 (6.7)
Optic glioma	1 (3.3)
Site (n=28)	
Right	11 (35.7)
Left	15 (53.6)
Midline	3 (7.4)
Type of glioma (n=26)	
Diffuse astrocytoma	5 (19.2)
Anaplastic astrocytoma	5 (19.2)
Glioblastoma multiforme	14 (53.8)
Oligodendroglioma	1 (3.8)
Pituitary adenoma	
Coexisting tumor of pituitary adenoma (n=30)	
Meningioma	27 (90.0)
Glioma	2 (6.7)
Chordoma	1 (3.3)
Type of pituitary adenoma (n=25)	
Nonfunctioning	11 (44.0)
Acromegaly	7 (28.0)
Prolactinoma	6 (24.0)
Cushing	1 (4.0)
Lymphoma†	16 (59.2)
Site (n=6)	
Right	4 (66.7)
Midline	2 (33.3)
Location (n=6)	
Frontal	2 (33.3)
Parietal	1 (16.7)
Temporal	2 (33.3)
Periventricular	1 (16.7)
Treatment (n=63)	
Conservative treatment/medication	5 (7.9)

Contd...

Table 1: Contd...

Characteristics	n (%)
One-surgery	25 (39.7)
Two-stage	22 (34.9)
Resection single tumor	11 (17.5)
Reported death (n=71)	15 (21.1)
Survival time-months	Median 10.00
	Mean 10.36
	SD 10.53
	Range 2 wk-42

*The original paper reported triple primary brain tumor in a patient.^[19,23] †All coexisting tumors of lymphoma are meningioma. SD: Standard deviation

chromosomal and genetic investigations in an assumed association between glioblastoma multiforme and meningioma. However, chromosomal analysis from peripheral blood cells of their patient was normal, with an absence of chromosome 17p deletions (p53). Beyond that, genetic analysis of tumor cells, using comparative genomic hybridization (CGH), found monosomy of chromosome 22 in meningioma cells and amplification of chromosome 7 and loss on chromosome 10 in glioblastoma cells.^[30] Similarly, Ohba *et al.* recently reported the results of CGH of tumor cells in which losses on 22q and gain 3, 5, 7, 11, 12, 13q, 15q, 17q, 20, and 21q in meningioma cell versus losses on 1p35, 9pter-21, 10, 11q23, 13q, 14q, 20q, and 22q and gain on 7 in glioblastoma cells.^[31] Since molecular variations isolated each tumor and were not linked with chromosome 4 or 5 alterations, there has been inconclusive common pathways, which carried out etiology of coexisting tumors.

We, therefore, believe that the concept of an association of multiple, primary brain tumors is incompletely supported and the current evidence does not allow us to conclude the existence of a common genetic pathway to explain pathogenesis of this condition. However, such investigations are severely limited because of this being such a scarce scenario, with only a few patients reported in the literature as of now.

In epidemiology, those of the female gender coupled with increasing age are at greater risk for meningioma while glioma is frequently observed in the mid-30s of male gender. Vernooij *et al.* conducted MRIs of the brain in 2000 people to determine the prevalence of incidental brain findings. The prevalence of incidental brain tumor was 1.6%. Meningioma was the most common tumor (0.9%) while pituitary adenoma could incidentally be found in 0.3%.^[6] The “Swiss cheese model” of accident causation explained that various errors fortuitously become accidents and incidents.^[44,45]

Adaptively, the concept of truly incidental events should be considered like a model.

Management issues

According to the literature, surgery is usually the treatment of choice for this scenario, but the decision to operate on both tumors at the same time is debatable. Simultaneous and adjacent tumors are found in only 42.2% (30/71) of cases. A single operation could be accepted in cases when both tumors are spatially contiguous.^[21] However, 49.2% (29/59) of stimulus cases show discontinuous tumors. Hence, the question will come up: “what tumor should be removed first?” We propose the idea of management for this scenario focus on the natural history of the disease.

Decision-making based on the presumed histopathology is an option. If a glioma is assumed, it will be the key determinant of the patients’ prognosis in this scenario because 73% (19/26) of gliomas are malignant while almost all of the other coexisting tumors are benign. Planning of any therapy would, therefore, focus on the glioma. This correlates well with the observation that 86.6% (13/15) of reported mortality occurred in patients in whom one of the lesions was a glioma. It is, therefore, our opinion that the malignant tumor should be removed first and treatment should center around the best, adjuvant management.

Hence, a second operation for the removal of any remaining lesions can be delayed or such lesions can also be observed only when these secondary tumors are likely benign, asymptomatic, small sized, or slow to grow.

CONCLUSION

The authors described a case series of rare patients, who presented with coexisting, primary brain tumors. The authors presented their patients in detail and summarized other clinical features found in the literature.

The pathomechanism of this rare disease scenario has been inconclusive. Further genetic investigations performed on tumors found in this scenario will help to answer the question whether there is a common underlying genetic pathway for both (coexisting) tumors. As of now, surgical management remains a mainstay of treatment. In cases in which high-grade glioma had been found as one of the lesions, it is evident that it conveys a poor prognosis.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal.

The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Acknowledgment

The authors would like to offer their special thanks to Dr. Pornprot Limprasert and Kanut Thapasathitsin for their advice on information in regard to human genetics.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Salvati M, D’Elia A, Melone GA, Brogna C, Frati A, Raco A, *et al.* Radio-induced gliomas: 20-year experience and critical review of the pathology. *J Neurooncol* 2008;89:169-77.
2. Schoenberg BS. Multiple primary neoplasms and the nervous system. *Cancer* 1977;40:1961-7.
3. Cushing H, Eisenhardt L. Meningiomas. Their Classification, Regional, Behavior, Life History and Surgical End Results. Ch. 22. Baltimore: Charles C Thomas; 1938. p. 506-7.
4. Ostrom QT, Gittleman H, Liao P, Rouse C, Chen Y, Dowling J, *et al.* CBTRUS statistical report: Primary brain and central nervous system tumors diagnosed in the united states in 2007-2011. *Neuro Oncol* 2014;16 Suppl 4:iv1-63.
5. Haberg AK, Hammer TA, Kvistad KA, Rydland J, Müller TB, Eikenes L, *et al.* Incidental intracranial findings and their clinical impact; The HUNT MRI study in a general population of 1006 participants between 50-66 years. *PLoS One* 2016;11:e0151080.
6. Vernooij MW, Ikram MA, Tanghe HL, Vincent AJ, Hofman A, Krestin GP, *et al.* Incidental findings on brain MRI in the general population. *N Engl J Med* 2007;357:1821-8.
7. Cahan WG, Woodard HQ, Higinbotham NL, Stewart FW, Coley BL. Sarcoma arising in irradiated bone: Report of eleven cases 1948. *Cancer* 1998;82:8-34.
8. Abs R, Parizel PM, Willems PJ, Van de Kelft E, Verlooy J, Mahler C, *et al.* The association of meningioma and pituitary adenoma: Report of seven cases and review of the literature. *Eur Neurol* 1993;33:416-22.
9. Arnetoli G, Nencioni L, Sottini M, Ammanati F. Simultaneous meningioma and glioma. Difficulties of neuroradiological diagnosis. Report of a case. *Ital J Neurol Sci* 1983;4:481-3.
10. Buccoliero AM, Taddei GL, Caldarella A, Mennonna P, Ammannati F, Taddei A, *et al.* Meningioma-primary brain lymphoma association. *Neuropathology* 2004;24:336-40.
11. Cannavò S, Curtò L, Fazio R, Paterniti S, Blandino A, Marafioti T, *et al.* Coexistence of growth hormone-secreting pituitary adenoma and intracranial meningioma: A case report and review of the literature. *J Endocrinol Invest* 1993;16:703-8.
12. Chandra PS, Hegde T. A case of coexisting cerebellopontine angle meningioma and schwannoma. *Neurol India* 2000;48:198.
13. Curto L, Squadrito S, Almoto B, Longo M, Granata F, Salpietro F, *et al.* MRI finding of simultaneous coexistence of growth hormone-secreting pituitary adenoma with intracranial meningioma and carotid artery aneurysms: Report of a case. *Pituitary* 2007;10:299-305.
14. da Costa LB, Riva-Cambrin J, Tandon A, Tymianski M. Pituitary

- adenoma associated with intraventricular meningioma: Case report. *Skull Base* 2007;17:347-51.
15. Dario A, Marra A, Cerati M, Scamoni C, Dorizzi A. Intracranial meningioma and astrocytoma in the same patient. Case report and review of the literature. *J Neurosurg Sci* 1995;39:27-35.
16. Davis GA, Fabinyi GC, Kalnins RM, Brazenor GA, Rogers MA. Concurrent adjacent meningioma and astrocytoma: A report of three cases and review of the literature. *Neurosurgery* 1995;36:599-604.
17. Deen HG Jr, Laws ER Jr. Multiple primary brain tumors of different cell types. *Neurosurgery* 1981;8:20-5.
18. George B, Kumar R, Johns P, Bartlett RJ, O'Brien D. Contiguous synchronous occurrence of primary cerebral lymphoma and meningioma. *Br J Neurosurg* 2007;21:35-8.
19. Gökalp HZ, Erdoğan A, Egemen N, Naderi S. Multiple intracranial tumors of different cell types. *Neurosurgery* 1990;27:463-5.
20. Gordon AS, Fallon KE, Riley KO. Meningioma interdigitated with primary central nervous system B-cell lymphoma: A case report and literature review. *Surg Neurol Int* 2011;2:181.
21. Goyal A, Singh AK, Sinha S, Tatke M, Singh D, Gupta V, *et al.* Simultaneous occurrence of meningioma and glioma in brain: Report of two cases. *J Clin Neurosci* 2003;10:252-4.
22. Honegger J, Buchfelder M, Schrell U, Adams EF, Fahlbusch R. The coexistence of pituitary adenomas and meningiomas: Three case reports and a review of the literature. *Br J Neurosurg* 1989;3:59-69.
23. Ildan F, Bağdatoğlu H, Boyar B, Hacıyakupoğlu S, Gönülşen G, Tunali N, *et al.* Combined occurrence of primary cerebral lymphoma and meningioma. *Neurosurg Rev* 1995;18:45-8.
24. Iyer VR, Sanghvi DA, Shenoy A, Goel A. Three distinct co-existent primary brain tumors in a patient. *J Cancer Res Ther* 2009;5:293-6.
25. Kuroiwa T, Ohta T, Kobata H, Yamamoto H, Kimura N. Coexistence of intracranial meningioma and primary malignant lymphoma – case report. *Neurol Med Chir (Tokyo)* 1990;30:268-71.
26. Lee EJ, Chang CH, Wang LC, Hung YC, Chen HH. Two primary brain tumors, meningioma and glioblastoma multiforme, in opposite hemispheres of the same patient. *J Clin Neurosci* 2002;9:589-91.
27. Mahvash M, Igressa A, Pechlivanis I, Weber F, Charalampaki P. Endoscopic endonasal transsphenoidal approach for resection of a coexistent pituitary macroadenoma and a tuberculum sellae meningioma. *Asian J Neurosurg* 2014;9:236.
28. Maiuri F, Cappabianca P, Iaconetta G, Esposito F, Messina A. Simultaneous presentation of meningiomas with other intracranial tumors. *Br J Neurosurg* 2005;19:368-75.
29. Mathuriya SN, Vasishtha RK, Dash RJ, Kak VK. Pituitary adenoma and parasagittal meningioma: An unusual association. *Neurol India* 2000;48:72-4.
30. Nestler U, Schmidinger A, Schulz C, Huegens-Penzel M, Gamberdinger UA, Koehler A, *et al.* Glioblastoma simultaneously present with meningioma – Report of three cases. *Zentralbl Neurochir* 2007;68:145-50.
31. Ohba S, Shimizu K, Shibao S, Miwa T, Nakagawa T, Sasaki H, *et al.* A glioblastoma arising from the attached region where a meningioma had been totally removed. *Neuropathology* 2011;31:606-11.
32. Raco A, Cervoni L, Salvati M. Two distinct intracranial tumors of different cell types in a single patient. Case report and review of the literature. *Neurosurg Rev* 1994;17:305-8.
33. Spallone A, Santoro A, Palatinsky E, Giunta F. Intracranial meningiomas associated with glial tumours: A review based on 54 selected literature cases from the literature and 3 additional personal cases. *Acta Neurochir (Wien)* 1991;110:133-9.
34. Suzuki K, Momota H, Tonooka A, Noguchi H, Yamamoto K, Wanibuchi M, *et al.* Glioblastoma simultaneously present with adjacent meningioma: Case report and review of the literature. *J Neurooncol* 2010;99:147-53.
35. Tugcu B, Kepoglu U, Gunal M, Gunaldi O, Karakaya B, Demirgil BT, *et al.* Two distinct primary brain tumors, in same region of the same patient: A case report. *J Neurooncol* 2006;79:219-20.
36. Vaquero J, Coca S, Martínez R, Jiménez C. Convexity meningioma and glioblastoma in collision. *Surg Neurol* 1990;33:139-41.
37. Yamada K, Hatayama T, Ohta M, Sakoda K, Uozumi T. Coincidental pituitary adenoma and parasellar meningioma: Case report. *Neurosurgery* 1986;19:267-70.
38. Yarman S, Minareci O. Value of petrosal sinus sampling: Coexisting acromegaly, empty sella and meningioma. *Neuroradiology* 2004;46:1027-30.
39. Zentner J, Gilsbach J. Pituitary adenoma and meningioma in the same patient. Report of three cases. *Eur Arch Psychiatry Neurol Sci* 1989;238:144-8.
40. Black PM, Carroll R, Glowacka D, Riley K, Dashner K. Platelet-derived growth factor expression and stimulation in human meningiomas. *J Neurosurg* 1994;81:388-93.
41. Hermanson M, Funa K, Hartman M, Claesson-Welsh L, Heldin CH, Westermark B, *et al.* Platelet-derived growth factor and its receptors in human glioma tissue: Expression of messenger RNA and protein suggests the presence of autocrine and paracrine loops. *Cancer Res* 1992;52:3213-9.
42. Gronwald RG, Adler DA, Kelly JD, Distecche CM, Bowen-Pope DE. The human PDGF receptor alpha-subunit gene maps to chromosome 4 in close proximity to c-kit. *Hum Genet* 1990;85:383-5.
43. Enrique P. Handbook of Growth Factors. Hematopoietic Growth Factors and Cytokines. Vol. 3. Florida: CRC press; 1994. p. 297-78.
44. Perneger TV. The swiss cheese model of safety incidents: Are there holes in the metaphor? *BMC Health Serv Res* 2005;5:71.
45. Reason J. Human error: Models and management. *BMJ* 2000;320:768-70.

Appendix 1: Summary of 71 cases of coexisting glioma and meningioma without previous radiotherapy or phakomatosis in English literatures

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
1	Deen <i>et al.</i> , 1981 ^{[17]*}	65 male	Craniopharyngioma		Simultaneous	Adjacent	Alive and well after 2 nd operation
2		67 male	Intrasellar meningioma				
3		54 female	Left frontal meningioma	Right frontal Grade 3 astrocytoma	Sequential	Nonadjacent	He died for 2 months after 2 nd operation
4	Arnetoli <i>et al.</i> , 1983 ^[9]	44 male	Left parietal Grade 2 oligodendroglioma		Simultaneous	Adjacent	He died for 10 months after 2 nd operation
5	Spallone <i>et al.</i> , 1991 ^[33]	47 male	Left parietal meningioma				
6		44 male	Right frontal meningioma		Simultaneous	Adjacent	He is in good health after surgery 2 years
7		47 male	Right frontal protoplasmic astrocytoma				
8		47 male	Right parieto-occipital meningioma	Right temporoparietal diffuse astrocytoma	Simultaneous	Nonadjacent	Death from recurrence of glioma at 18 months after surgery
9		33 male	Right frontal parasagittal meningioma	Left temporo-occipital GBM	Simultaneous	Nonadjacent	Death from recurrence of GBM at 12 months after surgery
10		48 male	Left insular malignant glioma		Simultaneous	Nonadjacent	Death at 2 months after operation
11		50 male	Retrosellar meningioma				
12	Davis <i>et al.</i> , 1995 ^{[16]†}	50 male	Left medial parietal meningioma	Left medial parietal diffuse astrocytoma	Sequential (10 months later)	Adjacent	Improved from seizure at 7 months after 2 nd operation
13		67 female	Right parietal meningioma	Right parietal anaplastic astrocytoma	Simultaneous	Adjacent	NA
14		58 male	Right parietal parasagittal meningioma	Right temporal anaplastic astrocytoma	Sequential	Nonadjacent	NA
15	Dario <i>et al.</i> , 1995 ^[15]	58 male	Right parietal parasagittal meningioma	Right temporal anaplastic astrocytoma	Sequential	Nonadjacent	NA
16	Lee <i>et al.</i> , 2002 ^[26]	70 female	Left temporo-occipital GBM	Right frontal meningioma	Simultaneous	Nonadjacent	Death from recurrence of GBM at 18 months after surgery
17		34 male	Right frontal anaplastic astrocytoma	Tentorial meningioma	Simultaneous	Nonadjacent	NA
18	Goyal <i>et al.</i> , 2003 ^[21]	34 male	Right frontal anaplastic astrocytoma	Tentorial meningioma	Simultaneous	Nonadjacent	NA
19		72 male	Right sphenoid meningioma		Simultaneous	Adjacent	NA
20		72 male	Right temporal GBM				

Contd...

Appendix 1: Contd....

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
14	Maiuri <i>et al.</i> , 2005 ^[28]	49 male	Left parietal meningioma		Simultaneous	Nonadjacent	Cured
15		63 female	GH-secreting pituitary adenoma		Simultaneous	Nonadjacent	Cured
16		61 female	Right frontotemporal meningioma		Simultaneous	Adjacent	Dead postoperatively
17		68 female	Nonfunctioning pituitary adenoma		Simultaneous	Nonadjacent	Death at 60 months from recurrence of glioma
18		65 male	Tuberculum sellae meningioma		Simultaneous	Adjacent	Death at 42 months after 2 nd operation
19		66 male	Third ventricle craniopharyngioma		Simultaneous	Adjacent	Death at 14 months after 2 nd operation
20		41 female	Left posterior third flax meningioma		Simultaneous	Adjacent	Cured at 9 years
21		60 female	Right temporal diffuse astrocytoma		Simultaneous	Nonadjacent	Cured at 11 years
22		64 male	Left anterior third flax meningioma		Simultaneous	Adjacent	16 months survival
23		42 male	Left frontal diffuse astrocytoma		Simultaneous	Adjacent	NA
24	Tugcu <i>et al.</i> , 2006 ^[35]	42 male	Left middle third parasagittal meningioma		Simultaneous	Adjacent	NA
25		72 female	Left parietal GBM		Simultaneous	Nonadjacent	Death at 13 months after operation
26		49 male	Left tentorial meningioma		Simultaneous	Adjacent	Death at 4 months after operation
27		68 female	Left CPA schwannoma		Simultaneous	Nonadjacent	Death at 13 months after operation
28	Nestler <i>et al.</i> , 2007 ^[30]	68 female	Left sphenoidal meningioma		Simultaneous	Nonadjacent	Death at 13 months after operation
29		72 female	Left CPA schwannoma		Simultaneous	Adjacent	Death at 4 months after operation
30		49 male	Right parasagittal meningioma		Simultaneous	Adjacent	Death at 14 months after operation
31		49 male	Right parietal lymphoma		Simultaneous	Adjacent	Death at 14 months after operation

Contd...

Appendix 1: Contd....

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
27	Iyer <i>et al.</i> , 2009 ^{[24]‡}	55 male	Olfactory groove meningioma Left temporal GBM Brainstem glioma		Simultaneous	Nonadjacent	Death from acute illness at 2 months after operation
28	Suzuki <i>et al.</i> , 2010 ^[34]	75 female	Left temporal meningioma Left temporal GBM		Simultaneous	Adjacent	NA
29	Ohba <i>et al.</i> , 2011 ^[31]	72 male	Right frontal meningioma	Right frontal GBM	Sequentially (4 years later)	Adjacent	GBM regrowth was detected at 7 months after operation
30	da Costa <i>et al.</i> , 2007 ^[14]	45 male	PRL-secreting pituitary adenoma Intraventricular meningioma		Simultaneous	Nonadjacent	NA
31	Raco <i>et al.</i> , 1994 ^[32]	78 female	Anterior flax meningioma	Frontal paramedian GBM	Sequentially (9 months later)	Adjacent	Death on GBM diagnosed admission
32	Curto <i>et al.</i> , 2007 ^[13]	61 female	GH-secreting pituitary adenoma	Right frontal meningioma	Sequentially (1 year later)	Nonadjacent	
33	Cannavò <i>et al.</i> , 1993 ^[11]	47 female	GH-secreting pituitary adenoma Retrosellar meningioma		Simultaneous	Adjacent	NA
34	Abs <i>et al.</i> , 1993 ^[8]	47 female	PRL-secreting pituitary adenoma Tuberculum sellae meningioma		Simultaneous	Adjacent	
35		61 female	ACTH-secreting adenoma Lt frontal meningioma		Simultaneous	Nonadjacent	
36		45 female	Prolactinoma Right temporal meningioma		Simultaneous	Nonadjacent	
37		45 female	Acromegaly Left parietal meningioma		Simultaneous	Nonadjacent	
38		82 female	Nonfunctioning pituitary adenoma Right sphenoidal meningioma		Simultaneous	Adjacent	
39		61 female	PRL-secreting pituitary adenoma	Right choroid plexus meningioma	Sequentially (12 years later)	Nonadjacent	
40		51 female	Nonfunctioning pituitary adenoma Right sphenoidal meningioma		Simultaneous	Nonadjacent	

Contd...

Appendix 1: Contd....

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
41	Ildan <i>et al.</i> , 1995 ^[23]	38 female	Right periventricular lymphoma		Simultaneous	Nonadjacent	
42	Honegger <i>et al.</i> , 1989 ^[22]	37 female	Left temporoparietal meningioma		Simultaneous	Nonadjacent	Postoperatively course was uneventful
43		49 female	PRL-secreting pituitary adenoma		Simultaneous	Nonadjacent	
44		74 male	Right temporal meningioma		Simultaneous	Nonadjacent	
45	Yamada <i>et al.</i> , 1986 ^[37]	52 female	Acromegaly		Simultaneous	Adjacent	Postoperatively course was uneventful
46	Zentner <i>et al.</i> , 1989 ^[39]	46	Left flax meningioma		Simultaneous	Adjacent	Postoperatively course was uneventful
47		63	Nonfunctioning pituitary adenoma		Simultaneous	Adjacent	Postoperatively course was uneventful
48		61	Right parasellar (medial sphenoidal) meningioma		Simultaneous	Adjacent	Postoperatively course was uneventful
49	Mahvash <i>et al.</i> , 2014 ^[27]	36 female	Nonfunctioning pituitary adenoma		Simultaneous	Adjacent	
50	Mathuriya <i>et al.</i> , 2000 ^[29]	58 female	Planum meningioma		Simultaneous	Nonadjacent	Patient had improved at follow up after 6 months
51	Yarman <i>et al.</i> , 2004 ^[38]	40 male	GH-secreting pituitary adenoma		Simultaneous	Nonadjacent	
52	Gordon <i>et al.</i> , 2011 ^[20]	65 female	Right parasagittal meningioma		Simultaneous	Adjacent	18 months follow-up, patient remain neurologically intracc and Karnofsky score is 90% and imaging is stable without new/enlarging mass

Contd...

Appendix 1: Contd....

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
53	George <i>et al.</i> ^[18]	71 male	Right temporal meningioma		Simultaneous	Nonadjacent	
54	Kuroiwa <i>et al.</i> , 1990 ^[25]	62 female	Right temporal lymphoma Right temporal meningioma		Simultaneous	Nonadjacent	185 days died after surgery
55	Buccoliero <i>et al.</i> , 2004 ^[10]	67 female	Olfactory groove meningioma	Bifrontal lymphoma	Sequentially (1 months later)	Adjacent	Patient alived 6 month after 1 st admission
56	Chandra and Hegde, 2000 ^[12]	35 female	Right CPA meningioma		Simultaneous	Adjacent	
57	Gökalp <i>et al.</i> , 1990 ^{[19]‡}	36 female	Right CPA schwannoma Pituitary adenoma		Simultaneous	Adjacent	
58		43 female	Parasellar chordoma Pituitary adenoma		Simultaneous	Nonadjacent	
59		50 female	Left frontal meningioma Pituitary adenoma	Left temporal and pons astrocytoma	Sequentially (3 years later)	Nonadjacent	
60		56 female	Left temporal meningioma	Left cerebellopontine angle neurinoma	Sequentially (11 months later)	Nonadjacent	
61		58 male	Right frontal meningioma		Simultaneous	Nonadjacent	
62		38 female	Pituitary adenoma		Simultaneous	Adjacent	
63		59 female	Optic glioma Right CPA neurinoma		Simultaneous	Nonadjacent	
64		62 female	Right occipital meningioma Right Frontal meningioma	Craniopharyngioma	Sequentially (1.5 years later)	Nonadjacent	
65	Vaquero <i>et al.</i> , 1990 ^[36]	75 female	Left parietal meningioma Left parietal GBM		Simultaneous	Adjacent	Death from pulmonary complication after surgery 2 weeks
66	Current series	68 female	Lt frontal meningioma Right insular diffuse astrocytoma		Simultaneous	Nonadjacent	Live and good performance (3 years follow-up)
67		63 male	Right frontoparietal GBM		Simultaneous	Nonadjacent	Loss to follow-up
68		62 female	Pituitary adenoma Planum sphenoidale meningioma Pituitary adenoma		Simultaneous	Adjacent	Live and good performance (2 years follow-up)

Contd...

Appendix 1: Contd....

Case number	Authors and year	Age and sex	1 st tumor	2 nd tumor	Chronological presentation	Location	Outcome
69		61 female	Right frontal intraosseous meningioma	Splenium anaplastic astrocytoma	Sequentially (10 years later)	Nonadjacent	Live and good performance (2 years follow-up)
70		50 female	Pituitary adenoma Bilateral small meningioma		Simultaneous	Nonadjacent	Live and good performance (1 years follow-up)
71		37 female	Pineal epidermoid tumor Right CPA schwannoma		Simultaneous	Nonadjacent	Live and good performance (1 years follow-up)

*Five cases enrolled were 14th-18th case of the original paper, [†]3rd case of the original paper was excluded due to previous radiotherapy, [‡]The original paper reported triple primary brain tumor in a patient, 3rd tumor is brainstem glioma diagnosed from MRI. CPA: Cerebellopontine angle, Si: Simultaneous, Sq: Sequential, NA: Not available, GBM: Glioblastoma multiforme, GH: Genomic hybridization, PRL: Prolactin, ACTH: Adrenocorticotrophic hormone