Letter to the Editor

Profile of non-Hodgkin lymphoma: An Indian perspective

DOI: 10.4103/sajc.sajc_60_18

Dear Editor,

Non-Hodgkin lymphoma (NHL) subtype, pattern of presentation as well as patient population, varies with geographical regions. The World Health Organization (WHO) 2016 lymphoma classification clarifies the diagnosis and management of NHL in relation to the stages of lymphomagenesis. It refines the diagnostic criteria to incorporate the expanding genetic/molecular landscape of NHL.^[1] In view of comparative data regarding the distribution of NHL subtypes in India is scarce in the literature, we did this retrospective analysis of newly diagnosed patients with NHL treated in a tertiary care

center. A total of 390 cases of adult (>18 years) NHL over a period of 27 months (May 1, 2013 and July 31, 2015) were registered in the Department of Medical Oncology at our institute (AIIMS, New Delhi). The individual NHL cases were retrospectively reviewed according to the WHO lymphoma classification 2016 revision, immunophenotypic expression and morphology. B-cell lymphomas formed 347 (89%) whereas T-cell lymphomas formed 43 (11%) of the NHLs. Diffuse large B-cell lymphoma (DLBCL) was the most common subtype which was present in 267 (68.5%) cases. Follicular lymphoma (FL), mantle cell lymphoma (MCL), marginal zone B-cell lymphoma, small lymphocytic lymphoma, and Burkitt's lymphoma amounted to 35 (9%), 20 (5%), 9 (2.3%), 5 (1.3%), and 5 (1.3%) of all NHLs cases, respectively. Among the T-cell lymphomas, peripheral T-cell lymphoma not otherwise specified (PTCL-NOS) was the most common subtype 15 (3.85%), followed by anaplastic large-cell lymphomas, T-cell lymphoblastic lymphoma, NK/T-cell lymphoma, and angioimmunoblastic T-cell lymphoma which accounted for 9 (2.3%), 7 (1.8%), 5 (1.3%), and 2 (0.75%) of all NHL cases, respectively. Details of all NHL are given in Table 1.

The present study of North Indian population shows key differences in the presentation as compared to the developing country and other parts of India. Details of epidemiological studies are summarized in Table 2.

Classifying NHL according to B- and T-cell type has therapeutic and prognostic significance. Epidemiology of the Indian subcontinent is marked different from that of the Western literature in view of marked preponderance of high-grade lymphoma especially DLBCL. In this study, 68.5% patients were DLBCL, which is significantly higher as compared to previous reported study from India and the West.^[2-6] FL and MCL were the second and third most common subtype of B-NHL, and PTCL-NOS is the most common T-cell lymphoma in adult. The younger average age (median 50 years) of our patients is consistent with the pattern seen in most other malignancies in India, due to the effect of a younger population

Table 1: Non-Hodgkin lymphoma subtypes in adults

Non-Hodgkin lymphoma (n=390)	n (%)
B-cell lymphoma	347 (89)
DLBCL	267 (68.5)
FL	35 (9.0)
MCL	20 (5.0)
Marginal zone (nodal and extranodal)	9 (2.3)
SLL	5 (1.3)
NHL-others (BL, SLVL, PMBCL, etc.)	11 (2.8)
T-cell lymphoma	43 (11)
PTCL-NOS	15 (3.85)
ALCL	9 (2.3)
T-cell lymphoblastic lymphoma	7 (1.8)
NK/T-cell lymphoma	5 (1.3)
AITL	2 (0.75)
Others	5 (1.3)

DLBCL: Diffuse large B-cell lymphoma, FL: Follicular lymphoma, MCL: Mantle cell lymphoma, SLL: Small lymphocytic lymphoma, BL: Burkitt's lymphoma, ALCL: Anaplastic large-cell lymphoma, PTCL-NOS: Peripheral T-cell lymphoma not otherwise specified, AITL: Angioimmunoblastic T-cell lymphoma, SLVL: Small splenic lymphoma with villous lymphocytes, PMBCL: Primary mediastinal B-cell lymphoma, NK: Natural killer

Table 2: Subtype distribution of lymphoma across India and the west (USA)

	Nimmagadda <i>et al.</i> ^[2]	Arora <i>et al.</i> ^[3]	Naresh <i>et al.</i> ^[4]	Sahni and Desai ^[5]	SEER database (USA) ^[6]	Present study
Number of patients	1431 (total 1723, 16.8% are HL)	4026	2773	935	77,490	390
Type of lymphoma (%)						
DLBCL	55	46.85	33.8	50.2	31.67	68.5
FL	11	10.51	12.6	13.1	32.81	9
ALCL	3	5.04	4.1	4.8	1.11	2.3
PTCL	2.7	5.91	1.9	4.6	3.27	3.9
BL	2.5	3.38	1.8	3.0	1.42	1.3
MCL	1.8	1.59	3.4	2.1	2.18	5
Others	24*	26.7*	36.8*	21.2*	27.5*	10*

*Others includes: Angioimmunoblastic T-cell lymphoma; adult T-cell lymphoma/leukemia; enteropathy-associated T-cell lymphoma; extranodal NK/T-cell lymphoma, nasal type; hairy cell leukemia; hepatosplenic T-cell lymphoma; lymphoblastic lymphoma; nodal marginal zone B-cell lymphoma; subcutaneous panniculitis-like T-cell lymphoma; lymphoplasmacytic lymphoma; primary cutaneous CD30-positive lymphoproliferative disorders; splenic marginal zone B-cell lymphoma; mycosis fungoides/Sézary syndrome; PMBCL. Small lymphocytic lymphoma. DLBCL: Diffuse large B-cell lymphoma, FL: Follicular lymphoma, ALCL: Anaplastic large-cell lymphoma, PTCL-NOS: Peripheral T-cell lymphoma not otherwise specified, BL: Burkitt's lymphoma, MCL: Mantle-cell lymphoma, CLL/SLL: Chronic lymphocytic leukemia/small lymphocytic lymphoma, NK: Natural killer, PMBCL: Primary mediastinal B-cell lymphoma

(Continue on page 166...)

(Letter to the editor continue from page 162...)

pyramid in our country.^[7,8] The present study of North Indian population shows key differences in the NHL subtypes as compared to the developed world and other parts of India.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

**Ajay Gogia, Chandan K. Das, Lalit Kumar, Atul Sharma,
M. C. Sharma¹, Soumya Mallick¹**

Departments of Medical Oncology and ¹Pathology, All India Institute of Medical Sciences, New Delhi, India

Correspondence to: Dr. Ajay Gogia,
E-mail: ajaygogia@gmail.com

References

1. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, *et al.* The 2016 revision of the World Health Organization (WHO) classification of lymphoid neoplasms. *Blood* 2016;6:643-569.
2. Nimmagadda RB, Digumarti R, Nair R, Bhurani D, Raina V, Aggarwal S, *et al.* Histopathological pattern of lymphomas and clinical presentation and outcomes of diffuse large B cell lymphoma: A multicenter registry based study from India. *Indian J Med Paediatr Oncol* 2013;34:299-304.
3. Arora N, Manipadam MT, Nair S. Frequency and distribution of lymphoma types in a tertiary care hospital in South India: Analysis of 5115 cases using the World Health Organization 2008 classification and comparison with world literature. *Leuk Lymphoma* 2013;54:1004-11.

(Continue on page 170...)

(Letter to the editor continue from page 166...)

4. Naresh KN, Srinivas V, Soman CS. Distribution of various subtypes of non-Hodgkin's lymphoma in India: A study of 2773 lymphomas using R.E.A.L. and WHO classifications. *Ann Oncol* 2000;11 Suppl 1:63-7.
5. Sahni CS, Desai SB. Distribution and clinicopathologic characteristics of non-Hodgkin's lymphoma in India: A study of 935 cases using WHO classification of lymphoid neoplasms (2000). *Leuk Lymphoma* 2007;48:122-33.
6. Nair R, Arora N, Mallath MK. Epidemiology of Non-Hodgkin's lymphoma in India. *Oncology* 2016;91 Suppl 1:18-25.
7. Gogia A, Raina V, Kumar L, Sharma A, Sharma MC, Mallick SR, *et al.* Follicular lymphoma: An institutional analysis *Asian Pac J Cancer Prev* 2017;18:681-5.
8. Das CH, Gogia A, Kumar L, Sharma A, Sharma MC, Mallick SR, *et al.* Mantle cell lymphoma: A North Indian tertiary care centre experience *Asian Pac J Cancer Prev* 2016;17:4583-6.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article: Gogia A, Das CK, Kumar L, Sharma A, Sharma MC, Mallick S. Profile of non-Hodgkin lymphoma: An Indian perspective. *South Asian J Cancer* 2018;7:162-70.

© 2018 The South Asian Journal of Cancer | Published by Wolters Kluwer - Medknow