

## Letter to Editor

## Cardiac volumes and left ventricular ejection fraction on myocardial perfusion scintigraphy

Dear Editor,

Our understanding of cardiac volumes – end-diastolic volume (EDV) and end-systolic volume (ESV) – and left ventricular ejection fraction (LVEF) was constructively and productively enhanced from reading the work of Mardanshahi *et al.*<sup>[1]</sup> It was very thought-provoking that there could be such a degree of difference from varying the filter cutoffs during reconstruction in myocardial perfusion scintigraphy (MPS) with single-photon emission computed tomography (SPECT). Being able to provide information such as ESV, EDV, and LVEF is a great advantage of MPS over some other noninvasive cardiac imaging modalities and should be regarded as a possible unintended benefit as it is far from the primary objective.<sup>[2]</sup>

We wonder if the values differ – and would the interpretation be different – using different software packages to calculate the EDV, ESV, and LVEF values. These can have significant differences. Just this week, we found that one patient was assessed by four Dimension Myocardial SPECT (Vital Images, Minnetonka, MN, USA) as having EDV, ESV, and LVEF of 43ml, 26ml, and 60%, respectively. By contrast, Quantitative Perfusion SPECT (Cedars-Sinai, Los Angeles, CA, USA) assessed these as 38ml, 15ml and 40%. The referring clinician would have a vastly different impression based on the differences in the information provided.

Beyond the anecdote, it has been discussed in greater detail in the literature. While some studies and articles suggest reasonable correlation such as Schaefer *et al.*,<sup>[3]</sup> others<sup>[4,5]</sup> have differed. In this context, the questions that we pose are:

1. How can we apply the findings to the different software?
2. How much weight can we place on this set of data given the discrepancies shown between the software most commonly used?

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

**JOSEPH C. LEE<sup>1,2</sup>, JIA WEN CHONG<sup>2</sup>**

<sup>1</sup>Department of Medical Imaging, The Prince Charles Hospital, Chermside, <sup>2</sup>Faculty of Medicine, University of Queensland, Herston, Queensland, Australia

## Address for correspondence:


Dr. Joseph C. Lee,  
Department of Medical Imaging, The Prince Charles Hospital,  
Chermside, Queensland 4032, Australia.  
E-mail: joseph.lee@health.qld.gov.au

**Submitted:** 15-Jan-2020, **Revised:** 06-Mar-2020,  
**Accepted:** 18-Mar-2020, **Published:** 14-Sep-2020

## REFERENCES

1. Mardanshahi AR, Alavi A, Yazdani J, Hosseinimehr SJ, Khoshakhlagh M, Dabirian M, *et al.* The correlation between myocardial perfusion scintigraphy and three-dimensional echocardiography in ejection fraction and cardiac volumes for determination of the nearest filtering parameters. *World J Nucl Med* 2019;18:373-7.
2. Lee JC, West MJ, Khafagi FA. Myocardial perfusion scans. *Aust Fam Physician* 2013;42:564-7.
3. Schaefer WM, Lipke CS, Standke D, Kühl HP, Nowak B, Kaiser HJ, *et al.* Quantification of left ventricular volumes and ejection fraction from gated 99mTc-MIBI SPECT: MRI validation and comparison of the Emory Cardiac Tool Box with QGS and 4D-MSPECT. *J Nucl Med* 2005;46:1256-63.
4. Ather S, Iqbal F, Gulotta J, Aljaroudi W, Heo J, Iskandrian AE, *et al.* Comparison of three commercially available softwares for measuring left ventricular perfusion and function by gated SPECT myocardial perfusion imaging. *J Nucl Cardiol* 2014;21:673-81.
5. Lavender FM, Meades RT, Al-Nahhas A, Nijran KS. Factors affecting the measurement of left ventricular ejection fraction in myocardial perfusion imaging. *Nucl Med Commun* 2009;30:350-5.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

| Access this article online            |   |
|---------------------------------------|---|
| <b>Website:</b><br>www.wjnm.org       | <b>Quick Response Code</b><br> |
| <b>DOI:</b><br>10.4103/wjnm.WJNM_8_20 |   |

**How to cite this article:** Lee JC, Chong JW. Cardiac volumes and left ventricular ejection fraction on myocardial perfusion scintigraphy. *World J Nucl Med* 2020;19:446.

© 2020 World Journal of Nuclear Medicine | Published by Wolters Kluwer - Medknow