Patient Self-Management of Asthma Using Mobile Health Applications: A Systematic Review of the Functionalities and Effects

Mehrdad Farzandipour^{1,2} Ehsan Nabovati^{1,2} Reihane Sharif^{2,3} Marzieh Heidarzadeh Arani⁴
Shima Anvari^{2,3}

¹Health Information Management Research Center, Kashan University of Medical Sciences, Kashan, Iran

²Department of Health Information Management and Technology, School of Allied Health Professions, Kashan University of Medical Sciences, Kashan, Iran

³Student Research Committee, Kashan University of Medical Sciences, Kashan, Iran

⁴Department of Pediatrics, Faculty of Medicine, Kashan University of Medical Sciences, Kashan, Iran

Address for correspondence Ehsan Nabovati, MSc, PhD, Qotbe Ravandi Boulevard, Pardis Daneshgah, Health Information Management Research Center, Kashan University of Medical Sciences, Kashan, Iran (e-mail: Nabovati@kaums.ac.ir).

Appl Clin Inform 2017;8:1068–1081.

Abstract

Objective The aim of this systematic review was to summarize the evidence regarding the effects of mobile health applications (mHealth apps) for self-management outcomes in patients with asthma and to assess the functionalities of effective interventions.

Methods We systematically searched Medline, Scopus, and the Cochrane Central Register of Controlled Trials. We included English-language studies that evaluated the effects of smartphone or tablet computer apps on self-management outcomes in asthmatic patients. The characteristics of these studies, effects of interventions, and features of mHealth apps were extracted.

Results A total of 10 studies met all the inclusion criteria. Outcomes that were assessed in the included studies were categorized into three groups (clinical, patient-reported, and economic). mHealth apps improved asthma control (five studies) and lung function (two studies) from the clinical outcomes. From the patient-reported outcomes, quality of life (three studies) was statistically significantly improved, while there was no significant impact on self-efficacy scores (two studies). Effects on economic outcomes were equivocal, so that the number of visits (in two studies) and admission and hospitalization-relevant outcomes (in one study) statistically significantly improved; and in four other studies, these outcomes did not improve significantly. mHealth apps features were categorized into seven categories (inform, instruct, record, display, guide, remind/alert, and communicate). Eight of the 10 mHealth apps included more than one functionality. Nearly all interventions had the functionality of recording user-entered data and half of them had the functionality of providing educational information and reminders to patients.

Conclusion Multifunctional mHealth apps have good potential in the control of asthma and in improving the quality of life in such patients compared with traditional interventions. Further studies are needed to identify the effectiveness of these interventions on outcomes related to medication adherence and costs.

Keywords

- ▶ asthma
- ▶ chronic diseases
- ▶ patient self-care
- ▶ e-health
- ▶ smartphone

received
July 8, 2017
accepted after revision
September 16, 2017

Copyright © 2017 Schattauer

DOI <https://doi.org/10.4338/ACI-2017-07-R-0116>
ISSN 1869-0327.

Background and Significance

Asthma is a common, chronic, noncommunicable disorder of the respiratory tract, affecting more than 334 million people of all ages in all parts of the world¹ and it is predicted that the number of patients with asthma will increase by 100 million more by 2025.² Asthma affects negatively the different domains of quality of life of patients due to long-term treatment and multiple attacks. Asthma is a growing cause of morbidity and mortality³ and imposes a significant burden on patients, families, and healthcare systems such that it is known as the third-ranking cause of hospitalization among children.⁴ It was estimated that a mean cost per year for each asthmatic patient in Europe and United States is \$USD 1,900 and 3,100, respectively.^{5–7}

An effective way to reduce costs and the harm caused by asthma, like most chronic diseases, is the management of the disease by the patients (self-management).⁸ Asthma-related outcomes such as unscheduled care (e.g., admissions, hospitalizations, emergency department attendances, and doctor consultations), markers of asthma control (e.g., days of restricted activity, night disturbance, and school absence), exacerbations, and quality of life can be improved with optimal self-management.^{9–11} According to the guidelines on asthma management,^{12–14} patient education in the treatment of asthma is essential and the main emphasis is to increase patients' knowledge and to promote patients' adherence to treatment regimens.^{15,16} Some problems such as nonadherence to medications and the lack of information about self-management have led to unwillingness of asthmatic patients to disease management activities.¹⁷

Due to the poor implementation of asthma self-management programs and their underutilization by patients, there is a need for innovative patient self-management methods.^{18–20} Information technology-based (IT-based) interventions such as mobile health (mHealth) applications (apps) have potential to enhance self-management outcomes through the provision of supports (e.g., information, education, and reminders) to patients.²¹ Estimates show that more than a billion people around the world will use mHealth apps on their smartphones by 2018.²² Moreover, mHealth apps are regarded as a multifunctional media to communicate information, share experiences, and collect patients' data, and they are highly customizable, low cost, and easily available. Therefore, this type of intervention has the potential to improve self-management for patients with chronic diseases.^{23–27}

So far, several studies have been conducted to evaluate the effects of mHealth apps on self-management outcomes in asthmatic patients, but all these studies reported different results.^{28–32} For example, a study showed that a mobile phone-based interactive self-care intervention improves pulmonary function, quality of life, asthma symptoms, and medications used and reduces rate of acute exacerbations.³² However, another study showed that the use of a smart phone application does not have a significant impact on asthma control, self-efficacy, quality of life, number of acute attacks, and admissions.²⁹ Therefore, to determine the over-

all impact of these types of interventions, it seems necessary that the effects reported by individual studies aggregate in a systematic review.

According to our knowledge, several systematic reviews have been conducted on the use of IT-based interventions to promote self-management in patients with asthma.^{23,33,34} In 2013, a systematic review was conducted to assess the effects of smartphone-based apps on self-management outcomes in patients with asthma. Only two randomized controlled trial (RCT) studies were included in that study; and finally, due to the small number of studies conducted in this field, it did not achieve clear and definitive results.²³ Then in 2014, a meta-review was conducted with the aim of evaluating the effectiveness of digital or online interventions on the outcomes of self-management in patients with asthma. This study demonstrated the positive impact of such interventions on some self-management outcomes, including quality of life and adherence to medications.³³ A systematic review was also recently conducted in 2016 on 12 studies with a wide range of IT-based interventions with a focus on the features of these interventions (e.g., education, monitoring/electronic diary, action plans, medication reminders/prompts, facilitating professional support, raising patient awareness of asthma control, and decision support for professionals). That review showed that the most successful interventions included multiple features, but effects on health-related outcomes in asthma patients were inconsistent.³⁴

The aforementioned review studies show that the number of studies performed regarding the effects of mHealth apps on the outcomes of patients with asthma in recent years has increased. According to our knowledge, since 2013 no new systematic review has been focused on the effects of mHealth apps on self-management outcomes in asthma patients. Understanding the functionalities of asthma apps helps researchers to develop a tailored content for their interventions. Therefore, to determine these functionalities and to support the widespread effective use in clinical practice, in addition to identifying information needs for asthmatic patients,³⁵ potential effects and the features of mHealth apps should be regularly reviewed.

Objective

The aim of this systematic review was to summarize the evidence regarding the effects of mHealth apps for a variety of self-management outcomes in patients with asthma and assess the functionalities of this type of interventions.

Methods

Data Sources and Search Strategies

The electronic databases searched were Medline (via PubMed), Scopus, and the Cochrane Central Register of Controlled Trials. A combination of MeSH terms and keywords related to asthma and mHealth (e.g., mobile phone, smartphone, apps, and ehealth) were used in the search strategy. The details of search strategies are described in ►Appendix. The search language was English, and the

search period was from January 2000 to July 2016 (because technologies existing prior to that time are unlikely to be representative of contemporary technologies, especially mobile health applications).³⁶ The literature search was updated on August 2, 2016. The reference list of included studies and relevant reviews were also examined to identify additional studies.

Inclusion and Exclusion Criteria

Studies were included if they met the following criteria:

1. Participants included people with asthma at any age, their caregivers, physicians, or other healthcare providers.
2. The studies concerned the feasibility assessment, pilot testing, or evaluation of an mHealth app intervention using a smart phone or a tablet computer.
3. The studies published in English.
4. Studies used at least one approach to the promotion of self-management skills (self-management refers to the behaviors that asthmatic patients and their family members perform to reduce the effects of this disease³⁷).

Studies were excluded if they:

1. Were conducted on patients with other chronic respiratory and allergic diseases.
2. Used mobile devices only to send text messages, access Web sites, or make phone calls.
3. Were reviews, conference proceedings, letters to the editor, protocols, or theses.
4. Focused only on the assessment of participants' attitude toward the mHealth apps (without intervention evaluation results).
5. Only designed and developed mHealth apps.

Screening and Data Extraction

Titles, abstracts, and full-texts were screened by two researchers independently (M.F. and R.Sh.). Full-text articles were retrieved from the searched databases and other available databases. Data were extracted from included studies using a structured form. Disagreements were resolved by discussion with a third researcher (E.N.), if necessary. For each included study, intervention characteristics, targeted outcomes, and effects of the intervention were extracted. The studies in this review were classified into four categories by the framework developed by the University of California San Francisco–Stanford Evidence-Based Practice Center: RCTs (level 1), non-RCTs (NRCTs, level 2), observational studies with controls (OWCs, level 3), and observational studies without controls (level 4).^{38,39}

Data Synthesis and Analysis

Due to the heterogeneity of studies in terms of methodology, statistical analyses, and outcomes, meta-analysis was not appropriate. To conduct narrative synthesis, the included studies were categorized based on different characteristics including intervention features, outcome categories, and effects of interventions on outcomes. The intervention features were classified into seven groups (inform, instruct, record, display, guide, remind/alert, and

communicate) according to technology functionality framework (→Appendix).⁴⁰ This framework was deductively derived using several external frameworks and has been used in another study (2016).⁴¹

Outcomes that were assessed in the included studies were categorized into three groups (clinical, patient-reported, and economic outcomes) based on the American Medical Association (AMA) reports for Performance Improvement (→Appendix).⁴² Similar to the systematic review by Nabovati et al.,⁴³ the effect of interventions was classified as being statistically significantly positive, positive without statistical argument, not effective (not statistically significant), negative without statistical argument, or statistically significantly negative. This systematic review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.⁴⁴

Quality Assessment of Studies

The quality of the included studies was assessed using the Effective Public Health Practice Project (EPHPP) quality assessment tool.^{45–47} The EPHPP is an appropriate instrument for the evaluation of the different study designs such as RCTs, NRCTs, and OWCs.⁴⁸ It contains six criteria to rate the studies: selection bias, study design, confounders, blinding, data collection methods, and withdrawals and dropouts. Each criterion was rated as strong, moderate, or weak, and then the overall quality score (global ratings) was calculated for each study. Studies with no weak ratings were given a global rating of “strong,” studies with one weak rating were given a global rating of “moderate,” and studies with two or more weak ratings were given a global rating of “weak.” As with the data extraction stage, each study was scored independently by two reviewers (M.F. and R.Sh.), and any disagreements were resolved through discussion with a third researcher (E.N.).

Results

Literature Search Results

The search results and study selection process were summarized in the PRISMA flow diagram (→Fig. 1). In total, 3,370 studies were found. After removal of duplicates and screening of titles and abstracts, 27 articles were eligible for full-text review. Finally, a total of 10 studies met all inclusion criteria.

Characteristics of the Included Studies

Key characteristics of the included studies are summarized in →Table 1. A total of 50% (5/10) of the studies were conducted in the United States, and one study in each of the following countries—United Kingdom, Australia, Taiwan, Turkey, and Canada. All the studies began to be published from 2011 onward. The duration of the interventions varied from 7 days to 6 months (median: 3.5 months, Q1: 1.46, Q3: 6).

Of the 10 included studies, 4 studies were RCTs, 3 were NRCTs, and the remaining 3 studies were observational designs without controls. The sample sizes of the studies ranged from 20 to 288 participants (median: 42.5, Q1: 20, Q3: 124).

Fig. 1 Flow diagram of the literature search and study selection.

In total, seven studies involved adults and adolescents (age range: 12–82 years), one study involved teenagers (age range: 5–18 years), one study involved pregnant women (older than 18 years), and the one remaining study did not report participants' ages. In none of the included studies, participants were not children, physicians, or other health-care providers. In seven studies, participants were patients with persistent asthma, two studies were patients with uncontrolled asthma, and in one study the type of asthma was not specified.

Quality Assessment of the Included Studies

The quality of the included studies is presented in ▶ **Table 2**. The total sum shows that 50 and 70% of the studies scored moderate concerning selection bias and blinding, and 40, 60, 70, and 40% scored strong regarding study design, confounders, data collection, and dropouts, respectively. Concerning

the global quality score, 40% of the studies were of strong quality, 30% moderate, and 30% weak.

Interventions Description

According to the inclusion criteria, type of interventions in all studies was mHealth apps. In total, three studies (3/10, 30%) used both Android and iOS platforms, two studies (2/10, 20%) used iOS platform, one study (1/10, 10%) used Android platform, only one study used an application on the Black-Berry platform (1/10, 10%), and in other studies platform type did not specify. The functionalities of mHealth apps for patients were categorized into seven groups: inform, instruct, record, display, guide, remind/alert, and communicate. Details of evaluating the apps' functionalities by the technology functionality framework are presented in Appendix 1. Most studies (8/10, 80%) employed more than one functionality. Nearly all interventions had the

Table 1 Main characteristics of the included studies (ranked according to year of publication)

Author, year, country	Study design	Duration	Sample size	Setting	Intervention description	Control group	Outcomes
Liu et al, 2011, ³² Taiwan	RCT	3 and 6 mo	120 patients	Outpatient clinics	The mobile phone-based interactive self-care software provided an electronic diary to record patients' daily asthma symptom score (sleep quality, the severity of coughing, difficulty in breathing, and daily activities affected by asthma), use of relievers, PEFR and PEFR variability	The control group participants were given an asthma symptom diary booklet and were asked to record their PEFR regularly, preferably daily	• Pulmonary function (FEV₁, FVC) • Quality of life (SF-12) • Episodes of exacerbation • Medications used for asthma control • Unscheduled visits to emergency department • Hospitalizations • Adherent to the intervention
Ryan et al, 2012, United Kingdom ²⁹	RCT (multicenter)	6 mo	288 patients (32 practices)	Primary care	Mobile phone-monitoring system: the t+ Asthma application, running on the web-enabled handset, allowed the patient to record symptoms, drug use, and peak flow from Piko meter	Patients in the control group were asked to keep a paper diary, recording the same data as the intervention group (symptoms, drug use, and peak flow readings) twice daily	• Asthma control (ACQ) • Self-efficacy (KASE-AQ) • Quality of life (mini-AQLQ) • Enablement (mPEI) • Acute exacerbations • Steroid courses prescribed • Unscheduled consultations • Out of hours visits • Emergency department visits • Admissions • Healthcare costs
Haze and Lynaugh, 2013, United States ⁵⁵	Observational without control group (pilot study)	3 to 6 mo	25 patients	Primary care	The smartphone application included four elements: (1) a questionnaire assessing asthma control (2) View a personalized asthma action plan (3) Select to view short education modules of either a slide or a video clip on a selected asthma topic (4) Select to send a text message using short message service	No control group for comparison	• The feasibility and efficacy of using a smartphone (patients and registered nurse care coordinators' perception) to communicate between a patient and his/her registered nurse care coordinator
Licskai et al, 2013, ⁴⁹ Canada	NRCT (pilot preintervention-postintervention study)	81 d	24 patients	Primary care	An agnostic web browser-based asthma action plan smartphone application: Patients securely transmitted symptoms and peak flow data daily, and received automated control assessment,	Before the intervention (self-management education and a written action plan)	• Quality of life (AQLQ) • Adherence to self-management and risk reduction recommendations • Knowledge (action plan and the AQHI) • Asthma symptom control

Table 1 (Continued)

Author, year, country	Study design	Duration	Sample size	Setting	Intervention description	Control group	Outcomes
Burbank et al, 2015, United States ⁵⁰	NRCT (preintervention–postintervention study)	8 wk	20 patients	Pediatric specialty clinics	A mobile application asthma action plan: participants were prompted by the mobile application to record either daily symptoms or peak flow measurements and to record medications. Also, the mobile application provided weekly asthma education	Before the intervention	- Healthcare utilization - Satisfaction (the ease of use, timeliness, simplicity, and clarity of the smartphone application, mobile health system)
Cingi et al, 2015, Turkey ³¹	RCT (multicenter)	3 mo	136 patients	Outpatient clinics	A mobile phone application (“physician on call patient engagement trial” [POPET]): enabling patients to communicate with their physician, and record their health status and send and receive messages, and ask for immediate assistance with an urgent message option, track their medicine use with a diary that sent automated reminders according to their prescribed treatment plan and complete the ACT	The control groups received an application that allowed completion of the ACT only at the beginning and end of the trial and did not include the communication, health status, or medication usage tracking	- ACT (frequency of shortness of breath and general asthma symptoms, use of rescue medications, the effects of asthma on daily functioning, and overall self-assessment of asthma control) - Medication adherence - Follow-up visits - Emergency visits - Status updates and patient satisfaction scores - Usage
Carpenter et al, 2016, United States ⁵⁶	Observational without control group	1 wk	20 patients	Pediatric practices	After their medical visit, the research assistant gave the adolescent an iPod Touch that contained two asthma self-management apps; one app was targeted to adults (features: peak flow monitoring, trigger and symptom tracking, diary/notes, chart and doctor report) and one was targeted to children (features: diary/notes,	No control group for comparison	- Self-observation - Self-judgment - Self-reaction

(Continued)

Table 1 (Continued)

Author, year, country	Study design	Duration	Sample size	Setting	Intervention description	Control group	Outcomes
					medication reminder, self-check, doctor appointment reminder, school form and my allergies and emergency plan). Adolescents were asked about their initial impressions of the apps including ease of use, perceived usefulness, and suggestions for improvement		
Cook et al, 2016, United States ²⁸	NRCT (before and after intervention)	4 mo	60 patients	Multispecialty clinic	Mobile app: patients interacted with the smartphone app when prompted by alerts and by patient-initiated actions such as information inquiries and data entry. Alerts included requests for self-assessment of asthma control, assessment of patient knowledge regarding asthma self-care, and resulting individualized coaching based on their previous entries	Before the intervention	• Asthma control (ACT score) • Lung function (FEV1) • Prescribed courses of systemic steroids • User experience, engagement, and feedback (patient satisfaction)
Roberts et al, 2016, United States ⁵⁷	Observational without control group (multicenter)	1 wk	20 patients	Two suburban pediatric practices	Adolescents and caregivers provided app feedback during a semistructured interview at a regularly scheduled clinic appointment and during a telephone interview 1 wk later. Friends completed one telephone interview	No control group for comparison	• Adolescent asthma patients' preferences for communicating with their caregivers and medical provider • Caregivers' preferences for communication with asthma patients
Zairina et al, 2016, Australia ³⁰	RCT (multicenter)	3 and 6 mo	72 patients	Antenatal clinics of two large maternity hospitals	A telehealth program supported by a handheld respiratory device and an Android smartphone application (Breathe-easy®): to record asthma symptoms and asthma medication usage weekly, also an automated feedback message regarding asthma status was sent weekly	Control group received usual medical care from their health professionals	• Asthma control (ACQ) • Quality of life (mini-AQLQ) • Lung function (FEV1 and FEV6) • Self-reported exacerbations • Unscheduled health-care visits • Days off work/study • Oral corticosteroid use

Abbreviations: ACQ, asthma control questionnaire; ACT, asthma control test; AQH, air quality health index; AQLQ, asthma quality of life questionnaire; FEV, forced expiratory volume; FVC, forced vital capacity; KASE-AQ, knowledge, attitude, and self-efficacy asthma questionnaire; mPEI, modified patient enablement instrument; NRCT, nonrandomized controlled trial; PEFR, peak expiratory flow rate; RCT, randomized controlled trial.

Table 2 Quality of the included studies

Study	Selection bias	Study design	Confounders	Blinding	Data collection	Drop-outs	Global rating
32	3	3	3	2	1	2	2
29	3	3	3	2	3	3	3
55	2	1	1	2	1	2	1
49	2	2	1	1	3	3	1
50	2	2	2	1	3	3	2
31	3	3	3	3	3	2	3
56	2	1	3	2	3	3	2
28	2	2	2	2	3	3	3
57	1	1	3	2	1	3	1
30	3	3	3	2	3	3	3
Sum (n, %)							
Weak	1, 10%	3, 30%	2, 20%	2, 20%	3, 30%	0, 0%	3, 30%
Moderate	5, 50%	3, 30%	2, 20%	7, 70%	0, 0%	3, 30%	3, 30%
Strong	4, 40%	4, 40%	6, 60%	1, 10%	7, 70%	7, 70%	4, 40%

Note: 3 = strong; 2 = moderate; 1 = weak.

functionality of recording user-entered data (9/10, 90%), half of them had the functionality of providing educational information and reminders to patients, 40% of them graphically displayed user-entered data and used apps to communicate between healthcare providers and patients, and 30% of them also provided instructions and guidance based on user-entered information.

Effects of Interventions on Outcomes

The categories of asthma outcomes and effects of interventions on them are shown in ▶Table 3. Among clinical outcomes, asthma control and lung function improved significantly in most studies which examined them. However, it should be noted that in two high-quality RCTs,^{29,30} these two outcomes did not change, whereas the positive results on these outcomes were from a mixture of RCTs and NRCTs of more variable quality.

In terms of patient-reported outcomes, the quality of life was significantly improved in the majority of studies that were assessed. This outcome was improved in multiple studies including a high-quality RCT,³⁰ while no improvement was found in another high-quality RCT.²⁹

There is limited evidence concerning the effects of interventions on economic outcomes. While the number of visits (in two studies) and admission and hospitalization (in one study) statistically significantly improved, economic outcomes in four other studies did not improve significantly.

▶Table 4 shows the outcomes that were assessed only in intervention groups and has not been compared with control groups. These outcomes were categorized into the patient-reported outcome category. mHealth apps had a positive impact on measured outcomes and there were no studies showing negative effects.

Discussion

In this systematic review, 10 studies (4 RCTs, 3 NRCTs, and 3 OWCs) were included regarding the effect of mHealth apps on patient self-management of asthma. These studies evaluated a wide range of outcomes related to asthma self-management, which were classified into three main categories: clinical (e.g., asthma control), patient-reported (e.g., medication adherence and quality of life), and economic outcomes (e.g., healthcare visits). In general, most of the studies (9/10, 90%) were indicative of the positive impact of mHealth apps on most outcomes assessed. In contrast, only one of the studies²⁹ did not report the positive impact on any of the outcomes assessed (including asthma control, asthma attacks, and adherence to treatment). Regarding the functionalities of the mHealth apps, nearly all interventions had the functionality of recording user-entered data and half of them had the functionality of providing educational information and reminders to patients.

Regarding the clinical outcomes (including control of asthma symptoms, lung function, and asthma attacks), 7 of the 10 studies assessed the impact of mHealth apps. Among the clinical outcomes, asthma control was investigated more than the others (lung function and asthma attacks), which indicates the importance of controlling this clinical outcome in patients with asthma. Five studies showed a positive, statistically significant improvement of this outcome,^{28,30,31,49,50} including two “strong” RCTs^{30,31} and one “strong” NRCTs.²⁸ These findings were similar to the results of previous systematic reviews in some outcomes, but were different in others. The study by Marcano et al did not report conclusive evidence about the impact of smartphone apps on asthma self-management outcomes.²³ But Hui and colleagues, by performing a meta-analysis of the three clinical trials, showed that

Table 3 The categories of asthma outcomes and effects of mHealth apps on them in intervention groups compared with control

Outcome category	Outcomes	Effect		
		Positive effect (statistically significant)	Positive effect (not stated about significance)	Not statistically significant
Clinical outcomes	Lung/Pulmonary function	28,32	–	30
	Symptom/Asthma control	28,30,31,49,50	–	29
	Acute attack/Exacerbation of asthma	50	30,32	29
Patient-reported outcomes	Medications use/adherence	28,32	31	29,30
	Self-efficacy	–	–	29,50
	Quality of life	30,32,49	–	29
	Patient enablement	–	–	29
Economic outcomes	Healthcare visits	31,32	–	29,30,49
	Days off work/study	–	–	30
	Hospitalization/Admissions for asthma	31	–	29,32,49
	Healthcare costs	–	–	29

apps-based interventions have a positive impact on the outcome of asthma control.³⁴ In contrast to the study by Marcano et al, which included only two studies (about the impact of mHealth apps on asthma self-management outcomes), in our study and the study by Hui et al, more studies were included that showed the positive impact of mHealth apps on improvement of this clinical outcome. In fact, this represents an

increase in the number of studies in recent years and the importance of this issue.

In the case of patient-reported outcomes, while three studies showed a significant positive impact on quality of life,^{30,32,49} one study showed no impact.²⁹ Similarly, Morrison and colleagues, by performing a meta-review, showed that digital interventions can improve the quality of life in patients with asthma.³³ Hui et al also³⁴ reported that in 50% of the interventions, quality of life improved over 6 to 30 months. The results of these three reviews show that IT-based interventions, especially smartphone apps, are appropriate interventions to improve the quality of life for asthma patients. Medication adherence is another patient-reported outcome and the results of our study and those by Hui et al³⁴ and Morrison et al³³ similarly demonstrated that this outcome improved in some studies and did not improve in others. Although mHealth apps, by providing reminders, have the potential to improve medication adherence in patients with chronic diseases,^{51,52} their effects on asthma patients is mixed and consequently there is a need for further studies in this area.

mHealth apps can help lower costs by facilitating the delivery of care and connecting patients to their healthcare providers.⁵³ In general, the results of this study showed that the number of visits, admissions, and hospitalization from economic outcomes improved in some studies and did not improve in others. Other economic outcomes, such as days off work or school and healthcare costs, did not improve in any study. Hui and colleagues in their systematic review showed that none of the smartphone apps had nonsignificant improvement in outcomes such as hospitalizations and emergency department visits.³⁴ Also, it was reported by Marcano et al that information about the effectiveness and

Table 4 Outcomes in intervention groups without control

Outcomes	Study
Patient satisfaction	28,31,49,50
Adherence to the intervention	32,49
Mobile app usage	31,50
Adherence to self-management and risk reduction recommendations	49
Knowledge	49
Patients' perception of smartphone app-supporting communication	55
Registered nurse/care coordinators' perception of smartphone app-supporting communication and accuracy of assessment data	55
Adolescent asthma patients' preferences for communicating with their caregivers and medical provider	57
Caregivers' preferences for communication with asthma patients	57
Self-management parameters (self-observation/self-judgment/self-reaction)	56

cost-effectiveness of smartphone and tablet self-management apps is conflicting.²³ In some studies, the short duration of the study and the small sample size were reported as the causes of inefficiency of IT-based interventions on outcomes related to asthma self-management.^{23,33,54} Therefore, future studies on the effectiveness of mHealth app interventions on the economic outcomes of asthma should consider the items listed above.

The results of this study showed the functionalities of mHealth apps for patients with asthma and were divided into seven groups (inform, instruct, record, display, guide, remind/alert, and communicate). Eight of the 10 interventions had more than one functionality. Similarly, Hui et al³⁴ classified the features of mobile-based interventions into seven categories (education, monitoring/electronic diary, action plans, medication reminders/prompts, facilitating professional support, raising patient awareness of asthma control, and decision support for professionals) and they concluded that the most successful interventions were those that had multiple functionalities. Due to numerous features provided by mHealth apps (such as recording patient-entered data and providing educational information and reminders) for patients and their relatively good accessibility, it can be expected that they have more effectiveness than other IT-based interventions (such as short message, phone call, email, and Web site) which have either limited features or low accessibility. Therefore, it is suggested that the designers apply multiple functionalities/features in designing these types of apps.

Strengths and Limitations

There are strengths and limitations in this study. The strengths included the use of broad search strategy which led to the identification of a large number of studies (3,370 articles). Two authors independently extracted data and assessed the quality of studies. A valid and comprehensive tool was used to assess the quality of the included studies. However, this study also has four limitations. First, studies were identified based on searches in only three databases. It is possible that this failed to identify some eligible studies. To address this in part, reference list of included studies and relevant reviews were hand searched. The second limitation is related to differences in objectives, methods, outcomes, and quality of the included studies that caused more difficult comparisons. The third limitation is that non-English language articles and conference proceedings (due to lack of full-text) were excluded; so, we may have lost some valuable studies in this field. The last limitation is that, due to the emerging mHealth apps, the results of unpublished and in-progress studies related to this field were not available. As a result, our review does not show experiences drawn from non-English and unpublished studies. However, among electronically accessible English-language studies, we applied a comprehensive and robust search strategy.

Implications for Practice

According to the evidence found in this systematic review on the effectiveness of mHealth app-based interventions to

improve self-management in patients with asthma, it is recommended to do the following to improve the acceptance of this type of interventions: use new models of self-care, including smartphone apps for better control of asthma, and enhance patients' quality of life, as well as use apps that provide multiple functionalities to patients.

Implications for Future Research

Given that the majority of the included studies (7/10, 70%) were conducted on adults with asthma, few studies were done on special populations, including adolescents and pregnant women with asthma. No studies assessed the effectiveness of mHealth apps on children and healthcare providers, and only two studies were conducted on patients with uncontrolled asthma.^{28,29} So, it is suggested that future studies focus more on these populations. Since most of the included studies (8/10, 80%) were conducted in developed countries, it is suggested that researchers conduct further investigations in developing and low-income countries. Considering that the results of this study showed that the effects of mHealth apps for patients with asthma on the outcomes related to adherence to medication/treatment and cost-effectiveness are not clear, further studies in this regard are needed.

Conclusion

Nearly all evaluated asthma apps have the functionality of recording patients' data and use more than one functionality. Multifunctional mHealth apps have good potential for improving control of asthma and quality of life in suffering patients. However, the effects of these interventions on outcomes related to healthcare costs and medication adherence are inconsistent.

Multiple Choice Question

In mHealth applications for patients with asthma, nearly all interventions have which of the following functionality?

- A. Recording user-entered data
- B. Providing educational information and reminders to patients
- C. Displaying user-entered data
- D. Providing instructions based on user-entered information

Correct Answer: The correct answer is A, Recording user-entered data. The results of this systematic review showed that nearly all mobile health applications for patients with asthma had the functionality of recording user-entered data (9/10, 90%).

Protection of Human and Animal Subjects

Human and/or animal subjects were not included in the study.

Funding

This study was supported by a grant from Kashan University of Medical Sciences Research Council (number: 9606).

Conflict of Interest

None.

References

- Global Asthma Network. The Global Asthma Report 2014. Auckland, New Zealand 2014
- Masoli M, Fabian D, Holt S, Beasley R; Global Initiative for Asthma (GINA) Program. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004; 59(05):469–478
- World Health Organization. Chronic obstructive pulmonary disease (COPD). Geneva, Switzerland: WHO. Available at: <http://www.who.int/respiratory/copd/en/>. Accessed September 2017
- Hall MJ, DeFrances CJ, Williams SN, Golosinskiy A, Schwartzman A. National hospital discharge survey: 2007 summary. 2010: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics
- Accordini S, Corsico AG, Braggion M, et al. The cost of persistent asthma in Europe: an international population-based study in adults. *Int Arch Allergy Immunol* 2013;160(01):93–101
- Ehteshami-Afshar S, FitzGerald JM, Doyle-Waters MM, Sadatsafavi M. The global economic burden of asthma and chronic obstructive pulmonary disease. *Int J Tuberc Lung Dis* 2016;20(01):11–23
- Nunes C, Pereira AM, Morais-Almeida M. Asthma costs and social impact. *Asthma Res Pract* 2017;3:1
- Itchev P, Śliwczyński A, Szykiewicz P, Marczak M. Mobile Health Applications Assisting Patients with Chronic Diseases: Examples from Asthma Care, in *M-Health Innovations for Patient-Centered Care*. IGI Global; 2016:170–196
- Andrews KL, Jones SC, Mullan J. Asthma self management in adults: a review of current literature. *Collegian* 2014;21(01):33–41
- Smith JR, Mugford M, Holland R, et al. A systematic review to examine the impact of psycho-educational interventions on health outcomes and costs in adults and children with difficult asthma. *Health Technol Assess* 2005;9(23):iii–iv, 1–167
- Welsh EJ, Hasan M, Li P. Home-based educational interventions for children with asthma. *Cochrane Database Syst Rev* 2011;(10): CD008469
- British Thoracic Society Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma: SIGN 153: Scottish Intercollegiate Guidelines Network; 2016. September 2016. Available at: <https://www.brit-thoracic.org.uk/publication-library/bts-guidelines/>
- National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol* 2007;120(5, Suppl):S94–S138
- GINA. Global Strategy for Asthma Management and Prevention (2017 Update). Global Initiative for Asthma; 2017
- Razi SP, Piroozmand N, Zolfaghari M, Kazemnejad A, Firoozbakhsh S. Education of how-to-use peak flow meter and following up via SMS on asthma self-management. *J Hayat* 2013;18(04):19–27
- López-Viña A, del Castillo-Arévalo E. Influence of peak expiratory flow monitoring on an asthma self-management education programme. *Respir Med* 2000;94(08):760–766
- Morowatisharifabad M, Nadrian H, Falahi A, Mohammadi M. Predictors of self-management behaviors in patients with asthma based on Green's Model of Health Education Planning. *J Sch Public Health Inst Public Health Res* 2009;7(03):37–49
- Ring N, Jepson R, Hoskins G, et al. Understanding what helps or hinders asthma action plan use: a systematic review and synthesis of the qualitative literature. *Patient Educ Couns* 2011;85(02):e131–e143
- Roberts NJ, Younis I, Kidd L, Partridge MR. Barriers to the implementation of self management support in long term lung conditions. *London J Prim Care (Abingdon)* 2012;5(01):35–47
- Kaya Z, Erkan F, Ozkan M, et al. Self-management plans for asthma control and predictors of patient compliance. *J Asthma* 2009; 46(03):270–275
- Schnall R, Mosley JP, Iribarren SJ, Bakken S, Carballo-Diéguez A, Brown Iii W. Comparison of a user-centered design, self-management app to existing mHealth apps for persons living with HIV. *JMIR Mhealth Uhealth* 2015;3(03):e91
- Jahns R, Houck P. Mobile Health Market Report 2013–2017. 2013
- Marcano Belisario JS, Huckvale K, Greenfield G, Car J, Gunn LH. Smartphone and tablet self management apps for asthma. *Cochrane Database Syst Rev* 2013;11(11):CD010013
- Beratarrechea A, Lee AG, Willner JM, Jahangir E, Ciapponi A, Rubinstein A. The impact of mobile health interventions on chronic disease outcomes in developing countries: a systematic review. *Telemed J E Health* 2014;20(01):75–82
- Cafazzo JA, Casselman M, Hamming N, Katzman DK, Palmert MR. Design of an mHealth app for the self-management of adolescent type 1 diabetes: a pilot study. *J Med Internet Res* 2012;14(03):e70
- El-Gayar O, Timsina P, Nawar N, Eid W. Mobile applications for diabetes self-management: status and potential. *J Diabetes Sci Technol* 2013;7(01):247–262
- Hollenbach JP, Cushing A, Melvin E, McGowan B, Cloutier MM, Manice M. Understanding clinicians' attitudes toward a mobile health strategy to childhood asthma management: a qualitative study. *J Asthma* 2017;54(07):754–760
- Cook KA, Modena BD, Simon RA. Improvement in asthma control using a minimally burdensome and proactive smartphone application. *J Allergy Clin Immunol Pract* 2016;4(04):730–737.e1
- Ryan D, Price D, Musgrave SD, et al. Clinical and cost effectiveness of mobile phone supported self monitoring of asthma: multi-centre randomised controlled trial. *BMJ* 2012;344:e1756
- Zairina E, Abramson MJ, McDonald CF, et al. Telehealth to improve asthma control in pregnancy: a randomized controlled trial. *Respirology* 2016;21(05):867–874
- Cingi C, Yorgancioglu A, Cingi CC, et al. The “physician on call patient engagement trial”(POPET): measuring the impact of a mobile patient engagement application on health outcomes and quality of life in allergic rhinitis and asthma patients. *International Forum of Allergy and Rhinology—Wiley Online Library*; 2015. Wiley Online Library
- Liu WT, Huang CD, Wang CH, Lee KY, Lin SM, Kuo HP. A mobile telephone-based interactive self-care system improves asthma control. *Eur Respir J* 2011;37(02):310–317
- Morrison D, Wyke S, Agur K, et al. Digital asthma self-management interventions: a systematic review. *J Med Internet Res* 2014; 16(02):e51
- Hui CY, Walton R, McKinstry B, Jackson T, Parker R, Pinnock H. The use of mobile applications to support self-management for people with asthma: a systematic review of controlled studies to identify features associated with clinical effectiveness and adherence. *J Am Med Inform Assoc* 2017;24(03):619–632
- Ansari R, Ahmadian L, Bazargan Harandi N, Mirzai M. Determining the content of a pediatric asthma website from parents' perspective: the Internet use and information needs. *Int J Pediatr* 2017;5 (06):5113–5123
- Huang X, Matricardi PM. Allergy and asthma care in the mobile phone era. *Clin Rev Allergy Immunol* 2016. Doi: 10.1007/s12016-016-8542-y
- Shegog R, Bartholomew LK, Parcel GS, Sockrider MM, Mâsse L, Abramson SL. Impact of a computer-assisted education program on factors related to asthma self-management behavior. *J Am Med Inform Assoc* 2001;8(01):49–61
- Kaushal R, Shojania KG, Bates DW. Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review. *Arch Intern Med* 2003; 163(12):1409–1416
- Garg AX, Adhikari NK, McDonald H, et al. Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review. *JAMA* 2005;293(10): 1223–1238

- 40 Aitken M, Gauntlett C. Patient Apps for Improved Healthcare: From Novelty to Mainstream. Parsippany, NJ: IMS Institute for Healthcare Informatics; 2013
- 41 Chaet AV, Morshedi B, Wells KJ, Barnes LE, Valdez R. Spanish-language consumer health information technology interventions: a systematic review. *J Med Internet Res* 2016;18(08):e214
- 42 American Medical Association. Measures Development, Methodology, and Oversight Advisory Committee: Recommendations to PCPI Work Groups on Composite Measures. Chicago, IL: American Medical Association; 2010
- 43 Nabovati E, Vakili-Arki H, Taherzadeh Z, et al. Information technology-based interventions to improve drug-drug interaction outcomes: a systematic review on features and effects. *J Med Syst* 2017;41(01):12
- 44 Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(07):e1000097
- 45 Thomas B, Ciliska D, Dobbins M, Micucci S. Quality Assessment Tool for Quantitative Studies Dictionary: The Effective Public Health Practice Project (EPHPP). McMaster University; 2008
- 46 National Collaborating Centre for Methods and Tools. Quality Assessment Tool for Quantitative Studies. Ontario: McMaster University Hamilton; 2008
- 47 Armijo-Olivo S, Stiles CR, Hagen NA, Biondo PD, Cummings GG. Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research. *J Eval Clin Pract* 2012;18(01):12–18
- 48 Effective Public Health Practice Project. Quality assessment tool for quantitative studies. 2010; Available at: <http://www.ephpp.ca/tools.html>
- 49 Licskai C, Sands TW, Ferrone M. Development and pilot testing of a mobile health solution for asthma self-management: asthma action plan smartphone application pilot study. *Can Respir J* 2013;20(04):301–306
- 50 Burbank AJ, Lewis SD, Hewes M, et al. Mobile-based asthma action plans for adolescents. *J Asthma* 2015;52(06):583–586
- 51 Dayer L, Heldenbrand S, Anderson P, Gubbins PO, Martin BC. Smartphone medication adherence apps: potential benefits to patients and providers. *J Am Pharm Assoc* (2003) 2013;53(02):172–181
- 52 Hayakawa M, Uchimura Y, Omae K, Waki K, Fujita H, Ohe K. A smartphone-based medication self-management system with realtime medication monitoring. *Appl Clin Inform* 2013;4(01):37–52
- 53 West DM. Improving health care through mobile medical devices and sensors. Brookings Institution Policy Report 2013;10:1–13
- 54 Mosnaim GS, Powell LH, Rathkopf M. A review of published studies using interactive Internet tools or mobile devices to improve asthma knowledge or health outcomes. *Pediatr Allergy Immunol Pulmonol* 2012;25(02):55–63
- 55 Haze KA, Lynaugh J. Building patient relationships: a smartphone application supporting communication between teenagers with asthma and the RN care coordinator. *Comput Inform Nurs* 2013;31(06):266–271, quiz 272–273
- 56 Carpenter DM, Geryk LL, Sage A, Arrindell C, Sleath BL. Exploring the theoretical pathways through which asthma app features can promote adolescent self-management. *Transl Behav Med* 2016;6(04):509–518
- 57 Roberts CA, Geryk LL, Sage AJ, Sleath BL, Tate DF, Carpenter DM. Adolescent, caregiver, and friend preferences for integrating social support and communication features into an asthma self-management app. *J Asthma* 2016;53(09):948–954

Appendix Database search strategy

PubMed

(Asthma* OR wheez* OR "asthma"[MeSH Terms]) AND (telephone[Title/Abstract] OR mobile[Title/Abstract] OR phone[Title/Abstract] OR smartphone[Title/Abstract] OR smart-phone[Title/Abstract] OR mhealth[Title/Abstract] OR m-health[Title/Abstract] OR "m health"[Title/Abstract] OR e-health[Title/Abstract] OR ehealth[Title/Abstract] OR "e health"[Title/Abstract] OR telecare[Title/Abstract] OR tele-care[Title/Abstract] OR telemedicine[Title/Abstract] OR tele-med[Title/Abstract] OR telehealth[Title/Abstract] OR tele-health[Title/Abstract] OR telenursing[Title/Abstract] OR tele-nursing[Title/Abstract] OR android[Title/Abstract] OR samsung[Title/Abstract] OR sony[Title/Abstract] OR ios[Title/Abstract] OR iphone*[Title/Abstract] OR ipad*[Title/Abstract] OR ipod*[Title/Abstract] OR "tablet device"[Title/Abstract] OR app[Title/Abstract] OR apps[Title/Abstract] OR handheld[Title/Abstract] OR hand-held[Title/Abstract] OR "personal digital"[Title/Abstract] OR PDA[Title/Abstract] OR blackberry[Title/Abstract] OR nokia[Title/Abstract] OR symbian[Title/Abstract] OR HTC[Title/Abstract] OR Palm[Title/Abstract] OR INQ[Title/Abstract] OR "tablet computer"[Title/Abstract] OR telemedicine[MeSH Terms] OR telephone[MeSH Terms] OR Computers, Handheld[MeSH Terms] OR Mobile Applications[MeSH Terms])

Scopus

TITLE-ABS KEY ((asthma* OR wheez*) AND (telephone OR mobile OR phone OR smartphone OR smart-phone OR mhealth OR m-health OR "m health" OR e-health OR ehealth OR "e health" OR telecare OR tele care OR telemedicine OR telemed OR telehealth OR tele-health OR telenursing OR tele-nursing OR android OR samsung OR sony OR ios OR iphone* OR ipad* OR ipod* OR "tablet device" OR app OR apps OR handheld OR hand-held OR "personal digital" OR pda OR blackberry OR nokia OR symbian OR htc OR palm OR inq OR "tablet computer")) AND (LIMIT-TO (LANGUAGE, "English")) AND (LIMIT-TO (SRCTYPE, "j")) AND (LIMIT-TO (PUBYEAR, 2016) OR LIMIT-TO (PUBYEAR, 2015) OR LIMIT-TO (PUBYEAR, 2014) OR LIMIT-TO (PUBYEAR, 2013) OR LIMIT-TO (PUBYEAR, 2012) OR LIMIT-TO (PUBYEAR, 2011) OR LIMIT-TO (PUBYEAR, 2010) OR LIMIT-TO (PUBYEAR, 2009) OR LIMIT-TO (PUBYEAR, 2008) OR LIMIT-TO (PUBYEAR, 2007) OR LIMIT-TO (PUBYEAR, 2006) OR LIMIT-TO (PUBYEAR, 2005) OR LIMIT-TO (PUBYEAR, 2004) OR LIMIT-TO (PUBYEAR, 2003) OR LIMIT-TO (PUBYEAR, 2002) OR LIMIT-TO (PUBYEAR, 2001) OR LIMIT-TO (PUBYEAR, 2000)) AND (LIMIT-TO (DOCTYPE, "ar") OR LIMIT-TO (DOCTYPE, "ip"))

Cochrane

(Asthma* OR wheez*)AND(telephone OR mobile OR phone OR smartphone OR smart-phone OR mhealth OR m-health OR "m health" OR e-health OR ehealth OR "e health" OR telecare OR tele-care OR telemedicine OR telemed OR telehealth OR tele-health OR telenursing OR tele-nursing OR android OR samsung OR sony OR ios OR iphone* OR ipad* OR ipod* OR "tablet device" OR app OR apps OR handheld OR hand-held OR

"personal digital" OR PDA OR blackberry OR nokia OR symbian OR HTC OR Palm OR INQ OR "tablet computer"

Table A1 Technology functionality framework⁴⁰

Functionality subcategory	Definition
Inform	Provide information in a variety of formats (text, photo, video)
Instruct	Provide instructions to the user
Record	Capture user-entered data
Display	Graphically display user-entered data/output user-entered data
Guide	Provide guidance based on user-entered information (e.g., recommend a physician consultation or course of treatment)
Remind/Alert	Provide reminders to the user
Communicate	Provide communication with health care provider/patients and/or provide links to social networks

Table A2 Categories of outcome measures⁴²

Types of outcome measures	Examples
Clinical outcomes	
Mortality	Infant death rates
Morbidity	Reduced rates of preeclampsia; reduced rates of gestational diabetes
Intermediate clinical outcomes	Blood pressure levels, blood glucose levels
Symptoms	Reduction in chest pain; reduced depression episodes
Clinical events	Stroke; Cancer; adverse events during childbirth
Patient-reported outcomes	
Health status	Quality of life—health status as perceived by the individual • Functional measures—SF-36, PROMIS • Pre- and posttreatment physical function • Pre- and posttreatment mental health (e.g., depression severity) • Pre- and posttreatment social/role function • Other measures of health status such as pain, vitality, perceived well-being, health risk status, etc.

Table A2 (Continued)

Types of outcome measures	Examples
Clinical outcomes	
Patient experiences with care (patient satisfaction, patient engagement/patient preference, patient education)	Consumer assessment (CAHPS); quality of life, compliance with treatment regimen; provider retention; shared decision making; engagement of family and friends; patient knowledge and understanding
Economic outcomes	
Resource use/costs	Healthcare service utilization; cost per episode of care

Table A3 Evaluation of intervention features

Functionality	Study									
	32	29	55	49	50	31	56	28	57	30
Inform	–	–	✓	✓	✓	✓	–	✓	–	–
Instruct	–	–	–	–	✓	–	–	✓	–	✓
Record	✓	✓	✓	✓	✓	✓	✓	✓	–	✓
Display	–	✓	✓	✓	–	–	✓	–	–	–
Guide	–	–	✓	✓	–	–	–	–	–	✓
Remind/Alert	–	–	–	✓	✓	✓	✓	✓	–	–
Communicate	–	–	✓	–	–	✓	✓	–	✓	–