


Supplementary Table S1 Platelet reactivity among study sites

Initial	<i>n</i>	Singapore	Malaysia	Hong Kong	New Zealand	<i>p</i> -Value
ASP	921	179 ± 156	285 ± 138	206 ± 187	283 ± 235	<0.001
ADP	926	427 ± 287	625 ± 335	453 ± 290	560 ± 323	<0.001
Final	<i>n</i>	Singapore	Malaysia	Hong Kong	New Zealand	<i>p</i> -Value
ASP	925	134 ± 101	216 ± 136	159 ± 151	201 ± 196	<0.001
ADP	928	248 ± 161	417 ± 252	300 ± 274	340 ± 222	<0.001

Supplementary Table S2 Platelet reactivity tests performed between 08:00–12:00H and 12:00–18:00H

Initial ^a	<i>n</i>	08:00 to 12:00	12:00 to 18:00	<i>p</i> -Value
ASP	921	220 ± 177	164 ± 138	<0.001
ADP	926	492 ± 314	304 ± 220	<0.001

^a97% of the samples were obtained primarily during office hours (08:00–18:00H).

**Fig. S1** Kaplan–Meier curve of 12-month MACCE.

Supplementary Table S3 Analysis of cut-off value of 203 AU*min and 300 AU*min for the ASPI test

	Intraprocedural thrombotic events OR (95% CI)	MACCE at 30 d OR (95% CI)
Initial ASPI		
≤203	Reference	Reference
>203	2.21 (0.64–7.61)	2.80 (0.76–10.38)
Initial ASPI		
≤300	Reference	Reference
>300	7.29 (2.43–21.89) ^a	2.35 (0.99–5.60)

^a $p < 0.05$.