

Intravenous fluid Considerations in the Resuscitation of a Head Injured Patient

Prakash Singh M Ch

Army Hospital (R&R), Delhi Cantt - 10

Abstract: Resuscitation of the head injured patients with shock poses a dilemma to the treating neurosurgeon. Whereas achieving hemodynamic stability is the objective in shock and in head injury, there is apprehension that aggressive fluid resuscitation may lead to aggravation of brain edema, especially if fluids used in resuscitation are hyposmolar. Instituting cerebral decongestants such as mannitol, frusemide to treat raised intracranial pressure may cause hypotension due to obligatory diuresis. This review article examines the literature on the resuscitation of head injured patient and closely looks into the suitability of various intravenous fluids available today for this emergency situation.

Keywords: cerebral edema, colloids, crystalloids, head injury, intravenous fluids

INTRODUCTION

Damage suffered by the brain from traumatic injury is not complete at the time of impact but continues to evolve during the ensuing period and contributes significantly to the poor outcome. Hypotension singularly is a significant factor for this secondary damage^{1,2}. In 717 patients of closed head injury in traumatic coma data bank, 35% had systolic pressure of less than 90mm of Hg and THIS was associated with 150% increase in mortality. Hypotension is poorly tolerated by injured brain. Because brain injury usually disrupts normal cerebral auto regulation, cerebral blood flow rises and falls with systolic blood pressure³. Hypotension on arrival in emergency, in the intensive care unit (ICU) and during surgery has been associated with increased mortality and poor outcome^{4,5}. It is therefore pertinent to avoid hypotension and treat it promptly in head injured patient. However there is apprehension that intravenous fluids may worsen cerebral edema and thus increase intracranial pressure (ICP)⁶. In a recent prevalence survey of 3147 patients in 200 ICUs, it was found that fluid balance was the primary factor that influenced the outcome⁷. It therefore becomes important, what degree of resuscitation and what kind of fluids should be given to head injured patient during resuscitation so as not to exacerbate cerebral edema and increase ICP, thus improving their outcome.

Degree of Resuscitation

In patients having head injury and hypotension, there is dichotomy of the views on the degree of resuscitation and

type of intravenous fluids. In patient without head injury a systolic blood pressure of 90 mm Hg (controlled hypertension) is considered adequate to reduce the chances of hemorrhage⁸. This may not be appropriate for a head injured patient, as it may not provide adequate cerebral perfusion pressure in the setting of impaired cerebral auto regulation and raised ICP. It is desirable that a cerebral perfusion pressure (CPP) of >70 mm of Hg is provided to the injured brain which is possible if mean arterial pressure of >90 mm of Hg is achieved. Thus the practice of controlled hypertension is not desirable in the setting of head injury⁹.

At the same time aggressive fluid resuscitation may lead to cerebral edema formation and poor outcome. This is particularly so if hypo-osmolar fluids are used for resuscitation^{1,6,8}. Fluid restriction in the head injury which was in vogue in the past has been found to be harmful as hypotension is associated with devastating secondary ischaemic injury to traumatic brain^{9,10,11}. Therefore aim, in such a situation, should be to achieve euvolemia so as to ensure adequate substrate delivery (Oxygen & glucose) to the injured brain. Use of aggressive fluid resuscitation has shown good results and even CPP based management has been advocated by some^{12, 13}.

What kind of fluids?

What kind of fluids should be used in head injured patients? The answers are not straightforward. A fluid which maintains or augments intravascular volume and doesn't increase cerebral edema is ideal and there are efforts towards that end. But let us have a look on the intravenous fluids that are available to us and generally used.

Crystalloids are generally used intravenous fluids for the resuscitation of hypovolemic patient. Ringer's lactate

Address for correspondence: Col Prakash Singh M Ch
Department of Neurosurgery
Army Hospital (R&R), Delhi Cantt 110010

has been found to be good¹⁴, but in the setting of head injury it is slightly hypo-osmolar with an osmolality of 270 mOsmol/L and thus may increase cerebral edema^{1,14}. Isotonic glucose (5% glucose) rapidly becomes hyposmolar after the utilization of glucose and will aggravate cerebral edema, thus raising ICP¹⁵. It may also contribute to the deleterious effects of hyperglycemia in a severely head injured patient.

Isotonic saline is good choice for head injured patient as it has an osmolality of 308 mOsmol/L and therefore very little potential for exacerbation of brain edema^{1,16,17}.

However the amount of crystalloids required is 3 times the amount of loss (3:1 rule) to amount for interstitial fluid deficit. But since these fluids are cheap and easily available and amongst them isotonic saline is good choice^{1,18}.

Hypertonic saline of late has been found to be effective fluid for the resuscitation of traumatic brain injury both in experimental and clinical studies^{8,19,20,21}. Because if its higher osmolality (514 mOsmol/L) it withdraws fluid from the tissues and increases intravascular volume and may help in preventing brain edema and rise in ICP by creating a favorable gradient for free water from extra vascular to the intravascular compartment^{22,23,24}. It is also known to improve cardiac output, thus increasing systolic arterial pressure resulting in better perfusion of the brain²³. Due to its vasodilator effect it also improves microcirculation. It is also known to have modulatory effect on the immune trauma response, which may also be contributory to the better outcome with hypertonic saline²⁶.

Hypertonic saline has been used as single bolus therapy and as continuous as well as in various concentrations (from 1.8-7.5%) with good outcome^{27,28}. However 3% saline is commonly available and thus can be easily used. Despite its good qualities for resuscitation of head injury patient certain concern remains with hypernatremia, hyperosmolality and its related renal failure^{28,29}. Therefore a close monitoring of serum sodium, osmolality and renal functions is mandatory. It should be seen that serum sodium does not rise beyond 155 mEq/L and serum osmolality does not exceed 320 mOsmol though even higher values have also been reported without adverse effects^{19,24,27,29}. Central pontine demyelination, which has been reported with rapid correction of hyponatremia³⁰, has not been reported with hypertonic saline resuscitation in head injured patients^{19,24,27,29}, yet this still remains a serious concern.

Colloids don't move out of intra vascular compartment due large size of their molecule, thus look appealing for resuscitation of head injured patient as they will help in stabilization of systolic blood pressure by with drawing

fluid into the intravascular compartment and at the same time will not aggravate cerebral edema^{18,31}. Volume required is also small (1:1 equal to amount of loss). But they are 40-80 times costlier than crystalloids. In addition have some potential adverse effects such as allergic reactions, anti coagulation side effects^{18,32}. These are less likely with hydroxyl starch than dextan³³. Despite theoretically appearing superiority of colloids limited available scientific data provide no definitive conclusions. A meta-analysis of several studies suggests rather a slightly lower mortality with crystalloids³⁴.

Recently albumen has also been looked into as a resuscitation fluid for head injury patients in SAFE study (saline versus Albumen fluid evaluation). It is supposed to draw fluid into intravascular compartment like colloids. However it did not show any benefit and was rather found to increase mortality even when its cost is prohibitive.³⁵

Blood has been ideal substitute for blood loss, however it takes some time for its availability. It is a well-known fact that blood loss upto 20% of blood volume can be treated with crystalloids and colloids but a loss of 30% or more requires blood replacement. A blood loss of 40% more is life threatening and requires immediate transfusion³⁶. Hemoglobin of less than 10 gm% also lowers oxygen carrying capacity of the blood and thus will impairs substrate delivery (oxygen) to the brain. Therefore a blood loss of more than 20% of blood volume should be corrected to improve oxygen delivery to brain³⁷.

CONCLUSIONS

In the era of molecular medicine when we have started looking at neurochemical and cellular mediators of secondary brain damage it is more pertinent that have closer look in the fluid resuscitation of a head injured patient as this can easily contribute in lessening the mortality and morbidity to some extent. Use of 5% glucose is contraindicated and even ringer lactate is not good being relatively hyposmolar^{1,14,15}. Isotonic saline is usually recommended but may not take care of raised ICP^{16,17,18}. Concomitant use of cerebral decongestants such as mannitol may be counter productive due to its diuretic effects which may worsen hypotension^{38,39,40}. Colloids and even albumen though theoretically seem appealing but have not demonstrated better outcome³⁴. Hypertonic saline seems ideal as it restores normovolemia, decreases cerebral edema and lowers ICP^{22,23,24}. Experimental studies & recent clinical studies do show promise on its efficacy and safety for resuscitation of head injured patients^{8,19,20,21}. However better designed double blind randomized studies are needed

to recommended its routine use for the treatment of raised intracranial pressure in head injury⁴¹.

REFERENCES

- Kelly DF. Emergency department management. In Marion DW (ed): Traumatic brain injury, New York, Thieme, 1999 pp 67-69.
- Jones PA, Andrew PJD, Midgley S, et al. Measuring the burden of secondary insults in the head injured patients during intensive care. *J Neurosurg Anaesth* 1994, 6: 4-14.
- Pick J, Chestnut RM, Marshall LF, et al. Extra cranial complications of severe head injury. *J Neurosurg* 1997, 77:901-7.
- Miller JD, Becker DP. Secondary insults to injured brain. *JR Coll Surg Edin* 1982, 27:292-8.
- Pietropaoli JA, Rogers FB, Shackford SR, et al. Deleterious effects of intra operative hypotension in patients with severe head injury. *J Trauma* 1992; 33:403-7.
- Feldman JA & Fish S. Resuscitation fluids for a patient with head injury and hypovolemic shock. *J Emer Med* 1991; 9:465-8.
- Manley G and Hemphill C. ICU fluid management in head Trauma: still No consensus. *Neurotrauma & critical care News*. 2005.
- Gunnar W, Merlotti G, Jonasson B, Barnett J. Resuscitation from hemorrhagic shock: Alteration of intracranial pressure after normal saline, 3% saline and dextran 40. *Ann Surg* 1986; 204: 686-92.
- Shenkin HA, Bezier HS, and Bouzarth WF. Restriction of fluid intake. Rational management of neurosurgical patient. *J Neurosurg* 1976; 45:432-9.
- Chestnut RM. Medical management of o intracranial pressure. In Cooper & Golfinos (eds): Head injury, 4th edition, Baltimore; William & Wilkins 2000-pp 229-63.
- Tien R and Chestnut RM. Medical management of the traumatic brain injured patient. In Cooper & Golfinos (eds): Head injury, 4th edition, Baltimore; William & Wilkins 2000, pp 457-482.
- York J, Arrillagh, Graham R and Miller R. Fluid resuscitation of patients with multiple injuries and severe closed head injury: Experience with aggressive fluid resuscitation strategy. *J Trauma Injury, Infection and Critical Care*. 2000, 48:376-80.
- Rosner MJ, Rosner SD, Johnson AH. Cerebral perfusion: management protocol and clinical results. *J Neurosurg* 1995, 83:949-62.
- Sutin KM, Ruskin KJ, Kaufman BS. Intravenous fluid therapy in neurologic injury. *Critical Care Clin* 1992; 8: 367-408.
- Bakey L, Crawford JD and White JC. The effects of intravenous fluids on cerebrospinal fluid pressure. *Surg Gynecol Obstet* 1954; 99:48-52.
- Kaieda R, Todd MM, Cook LN, et al. Acute effects of changing plasma osmolality and colloid oncotic pressure on formation of brain edema after cryogenic injury. *Neurosurgery* 1989; 24: 671-8.
- Tommasino C, Moore S, Todd MM. Cerebral effects of isovolemic haemodilution with crystalloids or colloid solutions. *Critical Care Med* 1988; 16:862-8.
- Reed RL II Resuscitation of the multiply injured patient. In Wilkin and Rengachary (eds) Neurosurgery Vol II second edition. New York; McGraw-Hill 1996, pp 2689-2697.
- Vassar MJ, Perry CA, Gannaway WL, et al. 7.5% sodium chloride/ dextran for resuscitation of trauma patients undergoing helicopter transport. *Arch Surg* 1991; 126:1065-72.
- Nakayama S, Sibley L, Gunther RA. Small volume resuscitation with hypertonic saline (2,300 m Osmol/L) *Circ Shock* 1984; 12:149-59.
- Shackford SR, Norton CM, Todd MM. Renal, Cerebral and pulmonary effects of hypertonic resuscitation in a porcine model of hemorrhagic shock. *Surgery* 1988; 104:553-560.
- Worthley LIG, Cooper DJ, Jones N. treatment of resistant intracranial hypertension with hypertonic saline *J Neurosurg* 1988; 68:478-80.
- Mazzoni MC, Borgstrom P, Anfors KE, et al Dynamic fluid redistribution in hyper osmotic resuscitation of hypo-volemic hemorrhage. *Am J Physiol* 1988, 139:629-37.
- Holcroft JW, Vassar MJ, Turner JE, et al. 3% NaCl/Dextran in resuscitation of severely injured patients. *Ann Surg* 1987; 206: 278-88.
- Wildenthal K, Skelton CL, Collman HN. Cardiac muscle mechanics in hyper osmotic solutions. *Am J Physiol* 1969, 217:302-8.
- Kolsen – Petersen JA. Immune effects of hypertonic saline : Fact or Fiction. *Acta Anaesthesia (Scand)* 2004; 48:667-678.
- Peterson B, Khanna S, Fisher B, Marshall L. Prolonged hyponatremia controls elevated intracranial pressure in head injured pediatric patients. *Critical Care Med* 2000; 28:1136-1143.
- Qureshi A, Saurez IJ, Castro A and Bhardwaj A. Use of hypertonic saline/acetate infusion in the treatment of cerebral edema in patients with head trauma. Experience at a single centre. *Journal of Trauma: Injury, Infection and Critical Care* 1999, 47:659-665.
- Shackfield SR, Bourguignon PR, Wald SL, et al. Hypertonic saline resuscitation of patients with head injury .A prospective randomized clinical trial. *Journal of Trauma :Injury, Infection & Critical Care* 1998; 44:50-58.
- Stern RH, Riggs JE, Schochet SS. Osmotic demyelination syndrome following correction of hyponatremia. *N Engl J Med* 1986, 34; 1535-42.
- Trammer B1, Iacobacci RJ, Kindt GW. Effects of crystalloids and colloids infusions on intracranial pressure and computerized

- electro encephalographic data in dogs with vasogenic brain edema.
Neurosurgery 1989, 25:173-8.
32. Harke H, Thoenics R, Margral I, Momsen W. The influence of different plasma substitutes on blood clotting and platelets functions during and after operations.
Anesthetist 1976, 25:366-73.
 33. Strauss RG, Stump DC, Henriksen RA, Saunders R. Effects of hydroxyl starch on fibrinogen, tissue clot formation and fibrinolysis.
Transfusion 1988; 25:230-4.
 34. Velanovich V. Crystalloid versus colloid fluid resuscitation: A meta – analysis of mortality.
Surgery 1989; 105: 65-71.
 35. Finfer S, Bellomo R, Boyce N, French J, Myburgh J, Norton R. A comparison of albumin and saline for fluid resuscitation in intensive care unit.
New Engl J Med 2004, 350:2247-56.
 36. Weight JA. Resuscitation and initial management.
Critical Care Clin 1993, 9:657-71.
 37. Gopinath SP and Roberston C. Intensive care unit management. In Marion DW (ed). Traumatic Brain injury: New York, Thieme 1999 pp 101-114.
 38. Kofke WA. Mannitol: Potential for rebound intracranial hypertension.
J Neurosurg 1992; 77:584-9.
 39. Mirski, MA, Denchev DI, Schnitzer MS and Hanley D. Comparison between Hypertonic Saline and Mannitol in the reduction of elevated intracranial pressure in a Rodent model of acute cerebral injury.
J Neurosurg Anaesth 2000;12: 334-44.
 40. Gunnar W, Jonasson O, Merlotti G, Stone J, Barret J. Head injury & haemorrhagic shock : studies of the blood brain barrier and intra cranial pressure after resuscitation with normal saline solution, 3% saline solution and dextran-40.
Surgery 1988; 103:398-407.
 41. Chestnut RM. Management of brain and spine injuries.
Critical Care Clinics 2004; 20:25-55.