

locus coeruleus. These data suggest complex MOA of VNS in both acute and chronic phases.

In recent years, we have studied the combination of VNS and corpus callosotomy, and found the combination of both techniques in selected patients achieves better results than both techniques separately. In this paper we would discuss our tentative experience and indications.

<http://dx.doi.org/10.1016/j.ijep.2015.12.003>

Experience with short video EEG in small town (yield and cost effectiveness)



Nashik Anand Diwan

Background: Semiology, type of seizure, true or pseudo-seizure (PNES – psychogenic non epileptic seizures) are often hard to differentiate clinically. Accurate diagnosis is essential for the optimum medical or surgical treatment and outcome for the patient. Most of the times, diagnosis requires inpatient video telemetry, which is both time consuming and expensive. Short video electroencephalography (SV-EEG) has been described previously and was shown to be a useful diagnostic tool in other specialist centres.

Objective: To determine the usefulness of SV-EEG in the diagnosis and management of various seizure types.

Method: After start of SV-EEG facility in Nashik, first 100 cases were selected, 1–55 years done over last 15 months.

Results: SV-EEG done on OPD basis for period of 1–8 h. Age – 1–62 years, M=64:F=36. Abnormal SV-EEG was reported in 75 patients. A positive SV-EEG supporting a diagnosis of true seizures occurred in 62 of patients (generalised epilepsy=15, focal epilepsy=47). PNES was diagnosed in 13 pts. Attacks recorded in these patients were 1–14. No attack or no interictal abnormality was noted in 25% of patients (n=25), resulting in an inconclusive SV-EEG. One patient had undergone anterior temporal lobectomy surgery based on this SV-EEG.

Conclusion: The positive rates of attacks from SV-EEG were comparable and even better to previously published results and show that SV-EEG is easily implemented in small town centres. It is cost effective method with very good diagnostic yield.

No	Inti site	Age	Sex	V EEG Duration (hrs)	Abnormality	No of events	No	Inti site	Age	Sex	V EEG Duration (hrs)	Abnormality	No of events
1 SD	10	MA	2	Normal	Nil	52 VP	11	F	7	PNES	Nil	1	
2 AD	8	MA	2	Normal	Nil	53 AS	34	MA	6	RT Hemisphere dysfunction	Nil	1	
3 SP	1impf	F	6	Left MTLE	Nil	54 JS	38	MA	6.5	RT Post Quadrant Epilepsy	Nil	1	
4 AN	2	MA	6	West syndrome	Nil	55 SA	7.5	MA	4.5	Left Frontal SMA Se	Nil	3	
5 HC	25	MA	6	Normal	Nil	56 SC	31	F	7.5	RT Ant Temp. Epilepsy	Nil	3	
6 AV	18	F	6	Normal	Nil	57 SP	31	F	7	Left MTLE	Nil	3	
7 VS	15	F	6	Normal	Nil	58 SN	56	F	6	Left MTLE	Nil	3	
8 WC	55	MA	6	Left MTLE	Nil	59 UJ	32	F	4	PNES	Nil	3	
9 VS	37	F	6	Normal	Nil	60 VJ	29	MA	4.5	PNES	Nil	4	
10 TD	15	F	8	RT Focal epilepsy	Nil	61 VV	32	MA	4	PNES	Nil	2	
11 AK	27	F	6	Normal	Nil	62 VV	24	MA	7	RT Occital Epilepsy	Nil	1	
12 VV	13	MA	8	PNES	2	63 AP	30	F	7	PNES	Nil	4	
13 SJ	30	F	6	RT MTLE	3	64 KC	18	MA	5.5	Normal	Nil	1	
14 GS	12	F	6	RT MTLE	2	65 AP	34	MA	6	Left Centro-Temporal Epilepsy	Nil	9	
15 GC	14	MA	6	Normal	0	66 DB	28	MA	7	RT Post Quadrant Epilepsy	Nil	1	
16 PK	16	F	7	Normal	Nil	67 DB	28	MA	5	RT Post Quadrant Epilepsy	Nil	1	
17 KD	3	F	4	Normal	5	68 GA	3.5	MA	3.5	Secondary LGS	Nil	22	
18 SP	14	MA	4	Typical absence	5	69 RB	27	MA	6	Normal	Nil	1	
19 AT	20	MA	6	Normal	Nil	70 RB	27	MA	4	Left & Rt MTLE	2	2	
20 SJ	33	F	7	RT Temporal Epilepsy	Nil	71 RA	21	F	7.5	RT Gen epileptisy	Nil	1	
21 HG	55	F	4	PNES	8	72 TS	18	MA	4.5	PNES	3	3	
22 CP	40	F	6	Normal	Nil	73 SC	18	MA	1	Gen Tonic	Nil	14	
23 MK	35	MA	5	PNES	3	74 RH	14	MA	7.3	RT Hemisphere Seizures	3	1	
24 SD	9	F	6	RT Post Quadrant Epilepsy	3	75 RA	21	F	7	PNES	Nil	1	
25 SK	6	MA	5	RT Post Quadrant Epilepsy	Nil	76 AA	12	F	7	RT Hemisphere Seizures	Nil	1	
26 SD	2	MA	6	Post-ictal Generalized	Nil	77 AS	24	MA	2.5	Normal	Nil	1	
27 RMP	15	F	2	PNES	2	78 AD	15	F	6	RT Gen Epilepsy-Absence	Nil	1	
28 VS	34	MA	6	PNES	3	79 AS	24	MA	2.5	Normal	Nil	1	
29 WP	51	MA	6	PNES	2	80 KJ	18	MA	7	Gen Tonic-Symptomatic generalisi	Nil	1	
30 PG	34	MA	6.5	Normal	1	81 BS	38	MA	5	RT MTLE	5	1	
31 NEM	24	MA	7	Left MTLE	1	82 SJ	63	MA	7	Normal	Nil	1	
32 US	4	MA	6	MS-LGS	2	83 SJ	63	MA	7	Normal	Nil	1	
33 MP	23	MA	7	Left Centro-Temporal	3	84 NR	26	MA	6	No localization/lateralization	Nil	1	
34 VS	24	MA	6.25	RT Post Quadrant Epilepsy	Nil	85 NA	21	F	7.5	Left MTLE	Nil	1	
35 MC	44	MA	6.5	Left RT Epilepsy	Nil	86 AK	17	MA	7	Left MTLE	Nil	1	
36 AP	11	F	6	Normal	Nil	87 AK	17	MA	7	Left MTLE	Nil	1	
37 JJ	6	F	6	Left Centro-Temporal	Nil	88 AS	22	MA	7.5	Left Centro-Temporal	Nil	1	
38 RM	40	MA	5	Left MTLE	Nil	89 SN	25	MA	7	Normal	Nil	1	
39 AS	5	MA	6	RT F-C Left Ant Temporal	Nil	90 SA	2	F	6	Self (convuls)	Nil	1	
40 VP	16	MA	5.5	RT Gen epileptisy-IME	3	91 PS	28	F	5.5	PNES (LME)	Nil	1	
41 MK	5	MA	5.5	LGS	Nil	92 RA	62	MA	7.5	Left Frontal	Nil	1	
42 DW	17	F	7	Normal	1	93 AN	3.5	MA	4	Post quadrant-sec generalisatd	Nil	1	
43 CW	12	MA	6	RT Frontopolar epilepsy	1	94 MJ	42	F	7	Left MTLE	Nil	1	
44 SB	17	F	6	Normal	3	95 AJ	42	F	7	Left MTLE	Nil	1	
45 TD	51	F	6	PNES	Multifoc	96 PA	13	MA	4	MTLE	Nil	1	
46 SB	3.5	F	6	Normal	3	97 AM	12	MA	7.5	RT Post Quadrant Epilepsy	Nil	1	
47 AP	1impf	MA	2	Normal	Nil	98 SB	45	F	7.5	Normal	Nil	1	
48 SJ	12	MA	7.5	RT Post-ictal-sec gen	Nil	99 AD	22	MA	6	PNES (Absence)	Nil	1	
49 SD	35	MA	7.5	Normal	Nil	100 MA	17	F	7.5	Left F-C Epilepsy	Nil	1	
50 SD	35	MA	7	Left MTLE	Nil								

<http://dx.doi.org/10.1016/j.ijep.2015.12.004>

Uncommon lesions in the medial temporal lobe presenting with intractable epilepsy



Anant Mehrotra

Dept of Neurosurgery, SGPGI, Lucknow

Introduction: Medial temporal lobe is a major site of seizure origin. Lesions present in the medial temporal lobe might predominantly present with epilepsy which might even be refractory to anti-epileptic drugs. We describe 8 uncommon lesions involving the medial temporal lobe which presented with intractable seizures.

Material and methods: 8 patients were included in the study from July, 2014 to July, 2015 who had presented to a tertiary care centre with seizures which were not controlled on medications. Complete clinical and radiological assessment of these cases was done. Treatment received and the seizure outcome (Engel's grade) were also noted.

Results: 6 cases presented with complex partial seizures out of which 5 had olfactory auras. 5 patients had right sided lesions and remaining 3 had left sided lesions. Among these 8 cases, 2 were tuberculomas and cavernomas each, 1 was epidermoid, 1 was ganglioglioma and 1 was a low grade glioma. All patients had a complete excision of the concerned lesion. Anterior medial temporal lobe resection (including amygdala and hippocampal resection) was performed in all these cases. 7 cases had Engel grade 1 seizure control and 1 had Engel grade 2 seizure control. No significant post-operative complication occurred in any of the patients.

Conclusion: Medial temporal lobe may harbour various pathologies and due to its location, it predisposes the patient for seizures. Lesionectomy when combined with AMTR gives good seizure control.

<http://dx.doi.org/10.1016/j.ijep.2015.12.005>

RNA-Seq analysis of hippocampal tissues reveals novel candidate genes for drug refractory epilepsy in patients with MTLE-HS



Aparna Banerjee Dixit¹, Jyotirmoy Banerjee¹, Arpna Srivastava², Manjari Tripathi³, Chitra Sarkar⁴, Aanchal Kakkar⁴, P. Sarat Chandra²

¹ Center for Excellence in Epilepsy, A Joint NBRC-AIIMS Collaboration, NBRC, Manesar, India

² Department of Neurosurgery, AIIMS, New Delhi, India

³ Department of Neurology, AIIMS, New Delhi, India

⁴ Department of Pathology, AIIMS, New Delhi, India

Array-based profiling studies shows aberrant gene expression patterns during epileptogenesis. We have performed RNAseq analysis of the hippocampal tissues resected from the patients with MTLE-HS to investigate the molecular basis of epileptogenicity and/or pharmacoresistance in MTLE. For non-epileptic control experiments, healthy tissues from tumour margins obtained during tumour surgeries were used. RNA sequencing was performed using standard protocols on Illumina HiSeq 2500 platform. Differential gene expression