


Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.elsevier.com/locate/ijnt

Case Report

Use of autologous comminuted calvarial fragments and pedicled pericranial graft for single stage repair of frontal and cranial base injury

Amit Agrawal^{a,*}, Surya Pratap Singh^b^a Professor of Neurosurgery, Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore 524003, Andhra Pradesh, India^b Resident of Neurosurgery, Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore 524003, Andhra Pradesh, India

ARTICLE INFO

Article history:

Received 4 March 2013

Accepted 7 April 2013

Available online 12 April 2013

Keywords:

Frontal bone

Frontal sinus fracture

Autologous bone

Bone graft

Calvarial bone graft

ABSTRACT

Frontal sinus fractures are usually caused motor vehicular accidents. Because of the anatomic position of the frontal sinus and the enormous amount of force; these injuries are often associated with injuries to skull base, intracranial, ocular, and maxillofacial structures and thus has a large potential of complications. In present case we describe the use of autologous comminuted calvarial fragments and pedicled pericranial graft for single stage repair of frontal and cranial base injury.

Copyright © 2013, Neurotrauma Society of India. All rights reserved.

1. Introduction

Frontal sinus fractures are usually caused motor vehicular accidents and account for 5–12% of all maxillofacial fractures.^{1–6} Because of the anatomic position of the frontal sinus and the enormous amount of force; these injuries are often associated with injuries to skull base, intracranial, ocular and maxillofacial structures¹ and thus has a large potential of complications.² In present case we describe the use of autologous comminuted calvarial fragments and pedicled pericranial graft for single stage repair of frontal and cranial base injury.

2. Case report

A 55-year-old male presented 6 h after road traffic accident. The vehicle in which he was traveling had head on collision with another vehicle. He sustained multiple facial injuries. There was history of loss of consciousness for 30 min, vomiting 3–4 episodes, nasal bleeding and left ear. On examination in emergency room his vital were stable, general and systemic examination was unremarkable except bilateral crepitations on chest auscultation. Neurologically he was in altered sensorium (GCS-E2, V3, M5), moving all four limbs equally and

* Corresponding author. Tel.: +91 8096410032 (mobile).

E-mail addresses: dramitagrawal@gmail.com, dramit_in@yahoo.com (A. Agrawal).

0973-0508/\$ – see front matter Copyright © 2013, Neurotrauma Society of India. All rights reserved.

<http://dx.doi.org/10.1016/j.ijnt.2013.04.005>




Fig. 1 – Pre-operative CT images showing extensive comminuted fractures involving frontal bone, anterior and posterior table of frontal sinus and anterior cranial base with patchy frontal contusions and pneumocephalus.




Fig. 2 – Intra-operative photograph showing (A) extensive comminuted and depressed frontal bone fragments, (B) grossly comminuted anterior, posterior tables of frontal sinus and cribriform plate and (C) grossly intact dura after removal of all loose bone fragments and sinus mucosa.

pupils were equal and reacting to light. Local examination revealed a large laceration on right side of forehead and depressed fragment on frontal bone. Pre-operative CT images showing extensive comminuted fractures involving frontal bone, anterior and posterior table of frontal sinus and anterior cranial base with patchy frontal contusions and pneumocephalus (Fig. 1). The patient was taken for emergency surgery. Bicoronal scalp flap was raised. Frontal bone and fracture were exposed (Fig. 2A). There was a depressed comminuted fracture of frontal bone with multiple loose fragments. All loose fragments and necrotic, unhealthy tissue including shattered cribriform plate was removed (Fig. 2B

and C). Almost all the posterior wall of the frontal sinus involved in comminuted fracture and it needed to be removed (Fig. 2C). The defect was thoroughly irrigated with diluted povidone-iodine and plenty of normal saline. All loose bone fragments were cleaned and washed with povidone-iodine and normal saline. Pedicled pericranial graft was harvested (Fig. 3A). Inner calvarial table fragment retrieved from fractured frontal was used to reconstruct cribriform plate (Fig. 3B). Pedicled pericranial graft was carpeted and secured over the bone graft to isolate the nasal cavity and redundant frontal sinuses from intracranial cavity (Fig. 3C). Utmost care was taken to avoid kinking and tenting of the pedicle of the graft.




Fig. 3 – (A) Harvested well-vascularized pedicled pericranial flap, (B) placement of bone fragment harvested from shattered posterior table of frontal sinus to reconstruct cribriform plate and for separation of the anterior skull base from the nasal cavity and (C) placement of pedicled pericranial graft over the bone fragment and raw surfaces.



Fig. 4 – Follow up photograph (left) showing acceptable cosmetic result and (right) X ray showing well aligned bone fragments and construction of the nasal roof.

Gel foam was placed over the graft. Artificial dural graft was placed over the intact dural to hold the bone fragments and the graft was secured to the intact frontal bone margins. Relatively larger loose fractured frontal bone fragments were placed over the graft. The scalp flap was closed over these fragments. Incision was closed in layers. A dressing with mild pressure was placed to avoid unnecessary movements of the bone fragments. Patient was electively ventilated and could be weaned off over next 48 h. He received intravenous broad spectrum antibiotics for five days and anti-epileptics were continued. The wound healed well with acceptable cosmetic results and no other complication (Figs. 4 and 5).

3. Discussion

Involvement of both outer and inner tables of frontal sinus represent greater severity of impact thus resulting in fractures of both plates, floor with bone fragmentation and derangement.⁷⁻¹³ If left untreated there can be infectious and

other life threatening complications more frequently after multiple fractures.^{1,14,15} To avoid these complications, early intervention not only makes anatomical reconstruction possible¹⁶ but also helps to prevent secondary operations and reduce the risk of infection.¹⁷ Management of these fractures require detailed clinical and radiographic evaluation, prompt surgical intervention (to excise any necrotic tissues inside or outside the cranial cavity), brain isolation by meticulous dural closure, ablation of the frontal air sinuses and repair of the bony defect.^{1,18} Reconstruction and recontouring of these frontal and anterior skull base defects can be technically challenging¹⁹⁻²¹ and the choice of surgical management is usually decided by the site and extent of the damage.¹² The basics of surgical management to repair these defects include (1) Separation of the anterior cranial base from the nasal cavity (preferably with vascularized tissue), (2) Thorough and complete mucosal removal, (3) Use of autologous material preferably cancellous bone) to repair the bony defect and (4) Also to restore the esthetics.^{1,4,19,22-27} Various factors favor the use of autologous bone for reconstruction including a lower incidence of graft loss, the mechanical, immunological, and technical-grafting properties of autologous bone.^{19-21,28-33} Apart from this autogenous local bone obviates second surgical intervention, donor site morbidity^{10,28-30,34} and ideal particularly in our circumstances where the finances may be a limiting factor.^{28-30,34} In summary, the use of autologous comminuted calvarial fragments and pedicled pericranial graft for single stage repair of frontal and cranial base injury in present case resulted in good functional and esthetic outcome.

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Metzinger SE, Metzinger RC. Complications of frontal sinus fractures. *Craniomaxillofac Trauma Reconstr.* 2009;2:27-34.




Fig. 5 – Six months follow up photograph showing well healed wound.

2. Montovani JC, Nogueira EA, Ferreira FD, Lima Neto AC, Nakajima V. Surgery of frontal sinus fractures: epidemiologic study and evaluation of techniques. *Braz J Otorhinolaryngol.* 2006;72:204–209.
3. Manolidis S, Hollier Jr LH. Management of frontal sinus fractures. *Plast Reconstr Surg.* 2007;120:32S–48S.
4. Metzinger SE, Guerra AB, Garcia RE. Frontal sinus fractures: management guidelines. *Facial Plast Surg: FPS.* 2005;21:199–206.
5. Xie C, Mehendale N, Barrett D, Bui CJ, Metzinger SE. 30-year retrospective review of frontal sinus fractures: the charity hospital experience. *J Craniomaxillofac Trauma.* 2000;6:7–15. discussion 16–18.
6. Luce EA. Frontal sinus fractures: guidelines to management. *Plast Reconstr Surg.* 1987;80:500–510.
7. Donald PJ, Bernstein L. Compound frontal sinus injuries with intracranial penetration. *Laryngoscope.* 1978;88:225–232.
8. Hybels RL, Newman MH. Posterior table fractures of the frontal sinus: I. An experimental study. *Laryngoscope.* 1977;87:171–179.
9. Hybels RL. Posterior table fractures of the frontal sinus: II. Clinical aspects. *Laryngoscope.* 1977;87:1740–1745.
10. Aydin S, Kucukyuruk B, Abuzayed B, Aydin S, Sanus GZ. Cranioplasty: review of materials and techniques. *J Neurosci Rural Pract.* 2011;2:162–167.
11. Adkins WY, Cassone RD, Putney FJ. Solitary frontal sinus fracture. *Laryngoscope.* 1979;89:1099–1104.
12. Stanley Jr RB. Fractures of the frontal sinus. *Clin Plast Surg.* 1989;16:115–123.
13. Heller EM, Jacobs JB, Holliday RA. Evaluation of the frontonasal duct in frontal sinus fractures. *Head Neck.* 1989;11:46–50.
14. Agrawal A, Johrapurkar SR. Neglected case of frontal sinus fracture. *Infect Dis Clin Pract.* 2008;16:309–310.
15. Piek J. Surgical treatment of complex traumatic frontobasal lesions: personal experience in 74 patients. *Neurosurg Focus.* 2000;9:e2.
16. Piotrowski WP, Beck-Mannagetta J. Surgical techniques in orbital roof fractures: early treatment and results. Official Publication of the European Association for Cranio-maxillo-facial Surgery. *J Craniomaxillofac Surg.* 1995;23:6–11.
17. Converse JM, Hogan VM. Open-sky approach for reduction of naso-orbital fractures. Case report. *Plast Reconstr Surg.* 1970;46:396–398.
18. EL-Rifaie KM, Taher AA. Frontobasal fractures. Guidelines to management Egypt. *J Plast Reconstr Surg.* 2006;27:113–119.
19. Gil Z, Abergel A, Leider-Trejo L, et al. A comprehensive algorithm for anterior skull base reconstruction after oncological resections. Official Journal of North American Skull Base Society [et al]. *Skull base.* 2007;17:25–37.
20. Ilankovan V, Jackson IT. Experience in the use of calvarial bone grafts in orbital reconstruction. *Br J Oral Maxillofac Surg.* 1992;30:92–96.
21. Ducic Y. Titanium mesh and hydroxyapatite cement cranioplasty: a report of 20 cases. Official Journal of the American Association of Oral and Maxillofacial Surgeons. *J Oral Maxillofac Surg.* 2002;60:272–276.
22. Tabaddor K, LaMorgese J. Complication of a large cranial defect. Case report. *J Neurosurg.* 1976;44:506–508.
23. Artico M, Ferrante L, Pastore FS, et al. Bone autografting of the calvaria and craniofacial skeleton: historical background, surgical results in a series of 15 patients, and review of the literature. *Surg Neurol.* 2003;60:71–79.
24. Neligan PC, Boyd JB. Reconstruction of the cranial base defect. *Clin Plast Surg.* 1995;22:71–77.
25. Disa JJ, Robertson BC, Metzinger SE, Manson PN. Transverse glabellar flap for obliteration/isolation of the nasofrontal duct from the anterior cranial base. *Ann Plast Surg.* 1996;36:453–457.
26. Thaller SR, Donald P. The use of pericranial flaps in frontal sinus fractures. *Ann Plast Surg.* 1994;32:284–287.
27. Yamamoto Y, Minakawa H, Yoshida T, et al. Role of bone graft in reconstruction of skull base defect: is a bone graft necessary. *Skull Base Surg.* 1993;3:223–229.
28. Agrawal A, Badhu B, Wakode P. Management of a case of massive orbital fracture associated with intracranial injury. *Nigerian J Ophthalmol.* 2008;15:13–16.
29. Agrawal A, Borle RM, Bhola N, Daga A, Bora S, Sachdeva S. Multiple fractures involving the orbit and incidental finding of large fourth ventricular epidermoid. *J Craniofac Surg.* 2009;20:261–262.
30. Rajendra PB, Mathew TP, Agrawal A, Sabharwal G. Characteristics of associated craniofacial trauma in patients with head injuries: an experience with 100 cases. *J Emerg Trauma Shock.* 2009;2:89–94.
31. Earley MJ, Green MF, Milling MA. A critical appraisal of the use of free flaps in primary reconstruction of combined scalp and calvarial cancer defects. *Br J Plast Surg.* 1990;43:283–289.
32. Hendus J, Draf W, Bockmuhl U. Reconstruction of the frontoorbital frame using split-thickness calvarial bone grafts. *Laryngorhinootologie.* 2005;84:899–904.
33. Agrawal A, Baisakhiya N, Bhola N. Split calvarial graft to repair the large frontal bone defect. *J Maxillofac Oral Surg.* 2010;9:166–169.
34. Agrawal A, Garg LN. Split calvarial bone graft for the reconstruction of skull defects. *J Surg Tech Case Rep.* 2011;3:13–16.