

Fertility, Pregnancy and Lactation After Bariatric Surgery – a Consensus Statement from the OEGGG

Fertilität, Schwangerschaft und Stillzeit nach bariatrischen Operationen – eine OEGGG-Stellungnahme



Authors

Tina Stopp^{1*}, Veronica Falcone^{1*}, Michael Feichtinger², Christian Göbl¹

Affiliations

- 1 Universitätsklinik für Frauenheilkunde, Klinische Abteilung für Geburtshilfe und feto-maternale Medizin, Medizinische Universität, Wien, Austria
- 2 Wunschbaby Institut Feichtinger, Wien, Austria

Key words

bariatric surgery, pregnancy, dumping syndrome, foetal growth restriction, glucose metabolism

Schlüsselwörter

bariatrische Chirurgie, Schwangerschaft, Dumping-Syndrom, fetale Wachstumsrestriktion, Glukosestoffwechsel

received 12.7.2018

revised 20.8.2018

accepted 20.8.2018

Bibliography

DOI <https://doi.org/10.1055/a-0706-7578>

Geburtsh Frauenheilk 2018; 78: 1207–1211 © Georg Thieme Verlag KG Stuttgart · New York | ISSN 0016-5751

Correspondence

Christian S. Göbl, MD MSc PhD

Universitätsklinik für Frauenheilkunde, Klinische Abteilung für Geburtshilfe und feto-maternale Medizin
Währinger Gürtel 18–20, 1090 Wien, Austria
christian.goebel@meduniwien.ac.at



Deutsche Version unter:

<https://doi.org/10.1055/a-0706-7578>

ABSTRACT

Bariatric surgery is recommended when other weight loss interventions, such as lifestyle modification or medications, have failed. A considerable number of women undergoing bariatric surgery are of childbearing age; hence, it is necessary to be aware of the effects of bariatric surgery on pregnancy for managing these patients. Although bariatric surgery is associated with positive effects on cardiovascular and metabolic parameters, side effects such as anaemia, the risk of developing internal hernia, altered glucose metabolism and the risk of small for gestational age offspring have to be considered. Pregnant women with a history of gastric bypass should not undergo the oral glucose tolerance test (OGTT) due to the high risk of hypoglycaemia. There are no contraindications for vaginal delivery and breastfeeding. This paper has been published as a consensus statement by the Austrian Society of Gynaecology and Obstetrics (OEGGG).

ZUSAMMENFASSUNG

Bariatrisch-chirurgische Maßnahmen werden zunehmend eingesetzt, wenn andere Interventionen wie die Lebensstilmodifikation oder eine Pharmakotherapie versagt haben. Vor allem ist auch eine steigende Anwendung dieser Eingriffe an Frauen im gebärfähigen Alter zu beobachten. Daher ist es notwendig, sich mit den Auswirkungen der bariatrischen Chirurgie auf Schwangerschaft und Geburt auseinanderzusetzen. Neben den positiven Effekten auf kardiovaskuläre und metabolische Parameter, müssen mögliche Nebenwirkungen wie mütterliche Anämie, ein erhöhtes Risiko für intraabdominelle Hernien, ein veränderter Glukosestoffwechsel sowie ein erhöhtes Risiko für eine fetale Wachstumsrestriktion beachtet werden. Der oGTT (oraler Glukosetoleranztest) ist bei Patientinnen nach einer bariatrisch-chirurgischen Intervention kontraindiziert, da dieser ein erhöhtes Risiko für hypoglykämie Episoden birgt. Für die vaginale Geburt und für das Stillen besteht keine Kontraindikation. Dieser Artikel wurde als Stellungnahme durch die Österreichische Gesellschaft für Gynäkologie und Geburtshilfe (OEGGG) veröffentlicht.

* contributed equally

Summary of Recommendations

1. Preconceptional counselling: The patient should be informed by an experienced obstetrician about the possible complications of pregnancy following gastric bypass (increased risk of foetal growth restriction, malnutrition, dumping syndrome, unreliability of the OGTT in diagnosing gestational diabetes, internal hernia, premature delivery). Patients who do not wish to have a child must be apprised of the reduced efficacy of oral contraceptives.
2. First presentation: Measurement of HbA_{1c} and fasting blood glucose, iron status, vitamin status, calcium and parathyroid hormone, nutritional supplementation, if necessary endocrinological assessment, repeating once per trimester (close monitoring as needed)
3. Explanation of the development of dumping symptoms, recommendation of a carbohydrate-modified diet, involvement if necessary of a dietologist
4. Monthly ultrasound checks with biometrics
5. Determination of diurnal blood glucose profile between week 24 and 28 of pregnancy for at least one week, or as of week 14–16 of pregnancy until delivery, depending on patient compliance
6. No contraindication for vaginal delivery
7. No contraindication for breastfeeding (breastfeeding should be recommended)
8. Appropriate nutritional supplementation during lactation

Background

Up to 21.1% of women in Austria aged 30 to 45 years are overweight (BMI 25–29.9 kg/m²) and 9.1% obese (BMI ≥ 30 kg/m²) [1]. Decreased fertility and complications of pregnancy such as diabetes, hypertension, pre-eclampsia and neonatal macrosomia are frequently reported in this population [2, 3].

Weight loss achieved by a change in lifestyle, drug therapy or bariatric surgery is associated with improved fertility and reduced rates of complications during pregnancy [4, 5].

Bariatric surgery is the most effective method for achieving weight loss [6]. Bariatric surgery is indicated above a BMI of ≥ 40 kg/m² or at a BMI of 35–40 kg/m² with associated comorbidities (possibly even at a BMI of > 30 to < 35 kg/m² in case of type 2 diabetes). The procedure may entail a restrictive or malabsorptive technique, or a combination of the two [7, 8].

Bariatric surgery is being offered increasingly to patients of childbearing age. It is therefore necessary to be aware of its effects on pregnancy and childbirth. In addition to the potential positive effects on cardiovascular [9] and metabolic parameters [10], potential adverse effects such as maternal anaemia [11], an increased risk of intra-abdominal hernias [12], impaired glucose metabolism [13] and a higher risk of foetal growth restriction [14] must be afforded closer attention.

Methods

The references for this consensus statement were extracted from the PubMed and MEDLINE databases using the following MeSH keywords: “obesity”, “bariatric surgery”, “pregnancy and bariatric

surgery”, “obesity and fertility”, “obesity and pharmacology”, “obesity and bariatric surgery”, “obesity and diabetes”, “diabetes and pregnancy”, “gestational diabetes and hypertension”, “obesity and hypertension”, “bariatric surgery and hypertension”, “obesity and heart disease”, “bariatric surgery and heart disease”, “gastric bypass and anemia”, “gastric bypass and hyperparathyroidism”, “bariatric surgery and vitamin D”, “dietary supplements and gastric bypass”, “gastric bypass and abdominal hernia”, “fetal macrosomia”, “infant, small for gestational age”, “breastfeeding and bariatric surgery”. Longitudinal studies such as cohort studies and systematic reviews (including meta-analyses) were given preference in the compilation of this consensus statement. In addition, the guidelines of the American College of Obstetricians and Gynaecologists (ACOG) on the care of patients during pregnancy and delivery after bariatric surgery were taken as a reference.

Preconceptional Aspects

Obesity is often associated with hyperandrogenaemia and polycystic ovary syndrome (PCOS). Follicle growth and oocyte maturation are impaired by the compensatory hyperinsulinaemia and often increased insulin resistance secondary to PCOS, resulting in reduced fertility [15, 16]. Hence, even young women who are overweight must often rely on assisted reproductive technologies (ART) to fulfil their desire for a child. Obesity is thus considered a risk factor for a lower number of eggs and embryos of poorer quality, accompanied by decreased rates of pregnancy and live birth [17]. Weight loss surgery appears to have a positive effect on hyperandrogenaemia in most female patients [18], and spontaneous conception has been achieved in up to 58% of infertile women following surgery [19]. Patients having undergone ART before and after bariatric surgery were found to have an increased number of eggs, better egg quality, and higher live birth rates during postoperative treatment cycles [20].

Patients who do not wish to have children must be informed that the efficacy of oral contraceptives could be reduced as a result of the operation (especially in the case of malabsorptive procedures) and that parenteral dosage forms or non-hormonal methods should be considered in the choice of a suitable contraceptive method [21].

Pregnancy after Bariatric Surgery

Glucose metabolism and gestational diabetes

Bariatric surgery reduces the risk of gestational diabetes [11, 22], but malabsorptive procedures in particular can cause fluctuations in blood glucose. In the 2 h OGTT, for instance, lower fasting glucose concentrations and an excessive increase in blood glucose were noted 60 min after glucose delivery followed by hyperinsulinaemic hypoglycaemia after 120 min. Consequently, it can be difficult to diagnose gestational diabetes as the OGTT does not deliver reliable results [13]. Alternatively, subcutaneous glucose could be measured continuously or capillary glucose measured repeatedly in the context of a diurnal blood glucose profile [23, 24]. A diurnal blood glucose profile is recommended as of week 24 to

28 of pregnancy (also earlier, if necessary) [23]. The target values are <95 mg/dl (fasting) and <140 mg/dl (1 h postprandial). No screening test is available as yet for gestational diabetes in pregnancy after weight-loss surgery. To rule out any pre-existing diabetes mellitus, the HbA_{1c} value (and fasting glucose in addition) should be determined at the first gynaecological examination [23].

Attention should also be paid in this patient population to the issue of a dumping syndrome (especially hypoglycaemia after carbohydrate intake). Early dumping syndrome occurs 15 min to 1 h after a carbohydrate-rich meal and is characterised by transient arterial hypotension, reflex tachycardia, flush, hyperhidrosis and even syncope [25,26]. Late dumping syndrome develops two to three hours after a meal and is caused by excessive insulin secretion with reactive hypoglycaemia and corresponding symptoms [25,26]. Standard treatment consists of a modified diet, mainly the avoidance of rapidly absorbed carbohydrates. Pharmacological interventions are not indicated in pregnancy, as insufficient data are available [25]. If a dumping syndrome is suspected, dietary advice from a dietologist or specialist with an additional qualification in nutritional medicine is recommended with a view to introducing a carbohydrate-modified diet.

Nutrient uptake

Bariatric surgery, especially when malabsorptive procedure is used, influences the uptake of micronutrients and macronutrients.

Several studies suggest a higher rate of iron deficiency anaemia following bariatric surgery [5, 27–29]. Therefore, the iron status should be checked even before conception, if possible, and closely monitored during pregnancy. The current recommendations for iron replacement in pregnant women having undergone bariatric surgery range from 40 mg to 600 mg per day [30–32]. The iron status should be checked during each trimester of pregnancy as a minimum [33].

Vitamin B₁₂ deficiency has also been observed in pregnant women who have had bariatric surgery (5,34). A sublingual dose of 350 µg/day or intramuscular administration of 1000 µg every 4–12 weeks is currently recommended with respect to supplementation [30,32]. The vitamin B₁₂ status should likewise be monitored in each trimester [33].

Very obese patients are frequently found to have vitamin D₃ deficiency; of patients in whom bariatric surgery is planned, up to 84% are affected [35]. Elevated parathyroid hormone concentrations may also be noted in preoperative patients due to decreased levels of vitamin D and calcium [35–37]. Bariatric surgery using a malabsorptive procedure, in particular, appears to intensify this problem. It has been reported that up to 73.6% of patients develop secondary hyperparathyroidism five years after surgery despite vitamin D supplementation [36], with long-term effects on bone health [32,38]. As an adequate maternal calcitriol concentration and the resulting increase in calcium absorption are important to foetal bone mineralisation, a sufficient supply of vitamin D₃ and calcium is essential in patients who have had bariatric surgery [30]. There is no global consensus on supplementation; current literature describes doses of 1000–2000 mg calcium citrate [30–32] and 50–150 µg or 1000 IU vitamin D₃ per day [30,

32], including regular laboratory tests and ultrasound checks for assessing foetal growth [30, 33, 39].

At present there is no evidence of an increased risk of folic acid deficiency in pregnant women following bariatric surgery [34, 40], provided this patient population likewise receives supplementation at the standard recommendation of 400 µg folic acid per day [31, 32, 39]. The folic acid concentrations should still be monitored in each trimester [33, 39].

Other deficiencies observed in pregnant women post bariatric surgery involve vitamins A and K. There is no general consensus with respect to supplementation of either vitamin, but several authors suggest that regular checks and, if necessary, supplementation is necessary, whereby in the case of vitamin A in particular attention must be paid to its potential teratogenicity, and daily dose of 5000 IU/day (ideally in the form of β-carotene) should not be exceeded [30,32]. Furthermore, a dose of 15 mg zinc per day and daily supply of at least 60 g of protein are recommended [30, 31, 33].

Pre-eclampsia

The growing incidence of pre-eclampsia worldwide is associated, among others, with the rise in pathological obesity [41–43]. Whereas approximately 2–8% of all pregnant women suffer pre-eclampsia [44], it can affect up to 13% of patients with grade 3 obesity [41]. Numerous studies have confirmed that bariatric surgery prior to conception can reduce the risk by up to 75% compared with an obese control group [45] in whom surgery was not performed [5, 46–50]. The effect appears to be more pronounced during the first two years postoperatively [51].

Foetal development

Several studies suggest that the risk of a foetus being small for gestational age (SGA) is increased in pregnancy after gastric bypass [14, 51, 52]. This complication could be related to the amount of weight lost by the mother and the surgical technique, as the risk of SGA appears to be less pronounced after a purely restrictive intervention [14]. The direct consequence of foetal adaptations in an undernourished mother [53] are metabolic effects reported in SGA children lasting through to adulthood [54]. At present, long-term data on the children of mothers having undergone bariatric surgery are still sparse; hence, so far only a limited assessment can be made regarding the extent to which the post-operative intrauterine conditions could influence the child's later development.

The increased risk of premature delivery (spontaneous or medically induced) must be noted, moreover, especially if the woman becomes pregnant during the first year after bariatric surgery. This could be attributable to increased weight loss during pregnancy, which may result in foetal malnutrition and premature delivery [55]. The data in this respect are not conclusive. In several studies, no significant correlation could be found between the increased risk of premature birth and prior bariatric surgery [51, 52]. The American College of Obstetricians and Gynecologists (ACOG) [33] and other authors [32] recommend delaying pregnancy until at least 12–18 months after bariatric surgery, but this is not supported by all current studies [56]. Obstetricians must bear in mind

that the optimal timing for pregnancy following gastric bypass has not yet been ascertained.

The literature reports that, compared with normal pregnancies, there is a slight but not statistically significant increase in the rates of intrauterine foetal death (IUFT) and perinatal mortality after bariatric surgery [52].

Growth checks at intervals of several weeks and registration with a perinatal centre are advisable with a view to preventing or minimising long-term consequences.

Surgical complications

The incidence of herniation after gastric bypass can reach 5%. Pregnant women have an even higher risk after bariatric surgery, possibly due to increased intra-abdominal pressure exerted by the pregnant uterus on the intestine [57]. In the event of acute abdomen, the obstetrician must also consider internal hernia and refer the patient with a view to surgery. If internal hernia is suspected, the patient should fast and be admitted for pain management and monitoring. If the pain recurs after building up the diet, subacute surgery should be performed. If the pain symptoms persist despite treatment and fasting, acute surgery will be necessary to prevent intestinal necrosis and foetal complications [58]. Timely diagnosis and treatment of an intra-abdominal hernia are crucial to maternal and foetal health, as deaths of both mother and child have been reported in the literature [59].

Lactation

Malabsorption and subsequent malnutrition are widespread consequences following bariatric surgery. Current studies are examining whether this nutrient deficiency affects the composition of breast milk; the data available so far provide no indication that the composition of the breast milk is inadequate after bariatric surgery [60]. In the light of current data, breastfeeding is certainly to be recommended [32, 60].

Conclusion

Bariatric surgery is associated with maternal and foetal effects on pregnancy [5]. Preconceptional counselling from the obstetrician is urgently recommended to ensure that the woman is aware of the potential risks of pregnancy following bariatric surgery. At first presentation during pregnancy, blood should be taken to determine the iron and vitamin status and intervene, where necessary. The patient should be informed about the symptoms and the avoidance of a dumping syndrome, and a diurnal blood glucose profile should be taken between week 24 and 28 of pregnancy [23]. The OGTT is contraindicated in pregnancy after bariatric surgery due to the increased risk of hypoglycaemia [13]. Regular ultrasound checks are recommended for monitoring foetal growth. Vaginal delivery and breastfeeding are not contraindicated.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Statistik Austria. BMI (Body Mass Index). Online: https://www.statistik.at/web_de/statistiken/menschen_und_gesellschaft/gesundheit/gesundheitsdeterminanten/bmi_body_mass_index/105608.html; last access: 07.03.2018
- [2] Catalano PM, Shankar K. Obesity and pregnancy: mechanisms of short term and long term adverse consequences for mother and child. *BMJ* 2017; 356: j1
- [3] Marchi J, Berg M, Dencker A et al. Risks associated with obesity in pregnancy, for the mother and baby: a systematic review of reviews. *Obes Rev* 2015; 16: 621–638
- [4] Best D, Avenell A, Bhattacharya S. How effective are weight-loss interventions for improving fertility in women and men who are overweight or obese? A systematic review and meta-analysis of the evidence. *Hum Reprod Update* 2017; 23: 681–705
- [5] Guelinckx I, Devlieger R, Vansant G. Reproductive outcome after bariatric surgery: a critical review. *Hum Reprod Update* 2009; 15: 189–201
- [6] Colquitt JL, Pickett K, Loveman E et al. Surgery for weight loss in adults. *Cochrane Database Syst Rev* 2014; (8): CD003641
- [7] Fried M, Yumuk V, Oppert JM et al. Interdisciplinary European Guidelines on Metabolic and Bariatric Surgery. *Obes Surg* 2014; 24: 42–55
- [8] Deutsche Gesellschaft für Allgemein- und Viszeralchirurgie. S3-Leitlinie: Chirurgie der Adipositas und metabolischer Erkrankungen. Online: <https://www.awmf.org/leitlinien/detail/II/088-001.html>; last access: 29.08.2018
- [9] Vest AR, Heneghan HM, Agarwal S et al. Bariatric surgery and cardiovascular outcomes: a systematic review. *Heart* 2012; 98: 1763–1777
- [10] Schauer PR, Nor Hanipah Z, Rubino F. Metabolic surgery for treating type 2 diabetes mellitus: Now supported by the world's leading diabetes organizations. *Cleve Clin J Med* 2017; 84 (7 Suppl. 1): S47–S56
- [11] Galazis N, Docheva N, Simillis C et al. Maternal and neonatal outcomes in women undergoing bariatric surgery: a systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2014; 181: 45–53
- [12] Vannevel V, Jans G, Bialecka M et al. Internal Herniation in Pregnancy After Gastric Bypass: A Systematic Review. *Obstet Gynecol* 2016; 127: 1013–1020
- [13] Feichtinger M, Stopp T, Hofmann S et al. Altered glucose profiles and risk for hypoglycaemia during oral glucose tolerance testing in pregnancies after gastric bypass surgery. *Diabetologia* 2017; 60: 153–157
- [14] Chevrot A, Kayem G, Coupaye M et al. Impact of bariatric surgery on fetal growth restriction: experience of a perinatal and bariatric surgery center. *Am J Obstet Gynecol* 2016; 214: 655.e1–655.e7
- [15] Gougeon A. Regulation of ovarian follicular development in primates: facts and hypotheses. *Endocr Rev* 1996; 17: 121–155
- [16] Nestler JE. Metformin for the Treatment of the Polycystic Ovary Syndrome. *N Engl J Med* 2008; 358: 47–54
- [17] Provost MP, Acharya KS, Acharya CR et al. Pregnancy outcomes decline with increasing body mass index: analysis of 239,127 fresh autologous in vitro fertilization cycles from the 2008–2010 Society for Assisted Reproductive Technology registry. *Fertil Steril* 2016; 105: 663–669
- [18] Escobar-Morreale HF, Santacruz E, Luque-Ramírez M et al. Prevalence of 'obesity-associated gonadal dysfunction' in severely obese men and women and its resolution after bariatric surgery: a systematic review and meta-analysis. *Hum Reprod Update* 2017; 23: 390–408
- [19] Milone M, De Placido G, Musella M et al. Incidence of Successful Pregnancy After Weight Loss Interventions in Infertile Women: a Systematic Review and Meta-Analysis of the Literature. *Obes Surg* 2016; 26: 443–451
- [20] Milone M, Sosa Fernandez LM, Sosa Fernandez LV et al. Does Bariatric Surgery Improve Assisted Reproductive Technology Outcomes in Obese Infertile Women? *Obes Surg* 2017; 27: 2106–2112

- [21] Yska JP, van der Linde S, Tapper VV et al. Influence of Bariatric Surgery on the Use and Pharmacokinetics of Some Major Drug Classes. *Obes Surg* 2013; 23: 819–825
- [22] Willis K, Alexander C, Sheiner E. Bariatric Surgery and the Pregnancy Complicated by Gestational Diabetes. *Curr Diab Rep* 2016; 16: 21
- [23] Adam S, Ammori B, Soran H et al. Pregnancy after bariatric surgery: screening for gestational diabetes. *BMJ* 2017; 356: j533
- [24] Bonis C, Lorenzini F, Bertrand M et al. Glucose Profiles in Pregnant Women After a Gastric Bypass: Findings from Continuous Glucose Monitoring. *Obes Surg* 2016; 26: 2150–2155
- [25] Narayanan RP, Syed AA. Pregnancy Following Bariatric Surgery – Medical Complications and Management. *Obes Surg* 2016; 26: 2523–2529
- [26] Rariy CM, Rometo D, Korytkowski M. Post-Gastric Bypass Hypoglycemia. *Curr Diab Rep* 2016; 16: 19
- [27] Devlieger R, Guelinckx I, Jans G et al. Micronutrient levels and supplement intake in pregnancy after bariatric surgery: a prospective cohort study. *PLoS One* 2014; 9: e114192
- [28] Nomura RMY, Dias MCG, Igai AMK et al. Anemia During Pregnancy after Silastic Ring Roux-en-Y Gastric Bypass: Influence of Time to Conception. *Obes Surg* 2011; 21: 479–484
- [29] Jans G, Matthys C, Bogaerts A et al. Maternal micronutrient deficiencies and related adverse neonatal outcomes after bariatric surgery: a systematic review. *Adv Nutr* 2015; 6: 420–429
- [30] Kaska L, Kobiela J, Abacjew-Chmylko A et al. Nutrition and Pregnancy after Bariatric Surgery. *ISRN Obes* 2013; 2013: 492060
- [31] Beard JH, Bell RL, Duffy AJ. Reproductive Considerations and Pregnancy after Bariatric Surgery: Current Evidence and Recommendations. *Obes Surg* 2008; 18: 1023–1027
- [32] Busetto L, Dicker D, Azran C et al. Practical Recommendations of the Obesity Management Task Force of the European Association for the Study of Obesity for the Post-Bariatric Surgery Medical Management. *Obes Facts* 2018; 10: 597–632
- [33] American College of Obstetricians and Gynecologists. ACOG practice bulletin no. 105: bariatric surgery and pregnancy. *Obstet Gynecol* 2009; 113: 1405–1413
- [34] Weng TC, Chang CH, Dong YH et al. Anaemia and related nutrient deficiencies after Roux-en-Y gastric bypass surgery: a systematic review and meta-analysis. *BMJ Open* 2015; 5: e006964
- [35] Fish E, Beverstein G, Olson D et al. Vitamin D Status of Morbidly Obese Bariatric Surgery Patients. *J Surg Res* 2010; 164: 198–202
- [36] Wei JH, Lee WJ, Chong K et al. High Incidence of Secondary Hyperparathyroidism in Bariatric Patients: Comparing Different Procedures. *Obes Surg* 2018; 28: 798–804
- [37] Chakhtoura MT, Nakhoul NN, Shawwa K et al. Hypovitaminosis D in bariatric surgery: A systematic review of observational studies. *Metabolism* 2016; 65: 574–585
- [38] Zhang Q, Chen Y, Li J et al. A meta-analysis of the effects of bariatric surgery on fracture risk. *Obes Rev* 2018; 19: 728–736
- [39] Mechanick JL, Youdim A, Jones DB et al. Clinical Practice Guidelines for the Perioperative Nutritional, Metabolic, and Nonsurgical Support of the Bariatric Surgery Patient – 2013 Update: Cosponsored by American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Surg Obes Relat Dis* 2013; 9: 159–191
- [40] Gascoin G, Gerard M, Sallé A et al. Risk of low birth weight and micronutrient deficiencies in neonates from mothers after gastric bypass: a case control study. *Surg Obes Relat Dis* 2017; 13: 1384–1391
- [41] Mbah A, Kornosky J, Kristensen S et al. Super-obesity and risk for early and late pre-eclampsia. *BJOG* 2010; 117: 997–1004
- [42] Spradley FT, Palei AC, Granger JP. Increased risk for the development of preeclampsia in obese pregnancies: weighing in on the mechanisms. *Am J Physiol Regul Integr Comp Physiol* 2015; 309: R1326–R1343
- [43] Spradley FT. Metabolic abnormalities and obesity's impact on the risk for developing preeclampsia. *Am J Physiol Regul Integr Comp Physiol* 2017; 312: R5–R12
- [44] World Health Organization. WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia. 2011. Online: http://whqlibdoc.who.int/publications/2011/9789241548335_eng.pdf; last access: 17.10.2017
- [45] Bennett WL, Gilson MM, Jamshidi R et al. Impact of bariatric surgery on hypertensive disorders in pregnancy: retrospective analysis of insurance claims data. *BMJ* 2010; 340: c1662
- [46] Maggard MA, Yermilov I, Li Z et al. Pregnancy and fertility following bariatric surgery: a systematic review. *JAMA* 2008; 300: 2286–2296
- [47] Magdaleno R, Pereira BG, Chaim EA et al. Pregnancy after bariatric surgery: a current view of maternal, obstetrical and perinatal challenges. *Arch Gynecol Obstet* 2012; 285: 559–566
- [48] Vrebosch L, Bel S, Vansant G et al. Maternal and Neonatal Outcome After Laparoscopic Adjustable Gastric Banding: a Systematic Review. *Obes Surg* 2012; 22: 1568–1579
- [49] Sheiner E, Willis K, Yogev Y. Bariatric Surgery: Impact on Pregnancy Outcomes. *Curr Diab Rep* 2013; 13: 19–26
- [50] González I, Lecube A, Rubio MÁ et al. Pregnancy after bariatric surgery: improving outcomes for mother and child. *Int J Womens Health* 2016; 8: 721–729
- [51] Yi X, Li Q, Zhang J et al. A meta-analysis of maternal and fetal outcomes of pregnancy after bariatric surgery. *Int J Gynaecol Obstet* 2015; 130: 3–9
- [52] Johansson K, Cnattingius S, Näslund I et al. Outcomes of pregnancy after bariatric surgery. *N Engl J Med* 2015; 372: 814–824
- [53] Fernandez-Twinn DS, Ozanne SE. Mechanisms by which poor early growth programs type-2 diabetes, obesity and the metabolic syndrome. *Physiol Behav* 2006; 88: 234–243
- [54] Levy-Marchal C, Jaquet D. Long-term metabolic consequences of being born small for gestational age. *Pediatr Diabetes* 2004; 5: 147–153
- [55] Stephansson O, Johansson K, Näslund I et al. Bariatric Surgery and Preterm Birth. *N Engl J Med* 2016; 375: 805–806
- [56] Kjær MM, Nilas L. Timing of pregnancy after gastric bypass – a national register-based cohort study. *Obes Surg* 2013; 23: 1281–1285
- [57] Leal-González R, De la Garza-Ramos R, Guajardo-Pérez H et al. Internal hernias in pregnant women with history of gastric bypass surgery: Case series and review of literature. *Int J Surg Case Rep* 2013; 4: 44–47
- [58] Gudbrand C, Andreasen LA, Boilesen AE. Internal Hernia in Pregnant Women After Gastric Bypass: a Retrospective Register-Based Cohort Study. *Obes Surg* 2015; 25: 2257–2262
- [59] Moore KA, Ouyang DW, Whang EE. Maternal and fetal deaths after gastric bypass surgery for morbid obesity. *N Engl J Med* 2004; 351: 721–722
- [60] Jans G, Devlieger R, De Preter V et al. Bariatric Surgery Does Not Appear to Affect Women's Breast-Milk Composition. *J Nutr* 2018; 148: 1096–1102