




Endoscopic treatment of complicated bile duct stricture after surgery for traumatic bile duct injury




► **Fig. 1** Occlusive percutaneous cholangiography using a 12-mm Fogarty balloon to dilate the common bile duct and make it visible under endoscopic ultrasound.




► **Fig. 2** To safely open the lumen-apposing metal stent in a small-caliber common bile duct, the stent was placed as deep as possible in the biliary tract through a guidewire.



► **Fig. 3** Biflanged stent and lumen-apposing metal stent at 1-month follow-up.



► **Fig. 4** “Hand-tailored,” fully covered, self-expandable metal stents placed through the choledochoduodenostomy crosswise to avoid excessive tension of the anastomotic area.



► **Fig. 5** Final outcome on the hepaticojejunostomy.

Hepatic lesions are common in abdominal traumas [1] and generally involve liver parenchyma.

Nevertheless, bile duct lesions are rare and challenging to treat [2, 3].

Biliodigestive anastomoses are the most frequently performed surgeries [3], but the risk of long-term adverse events is high, especially for biliary anastomotic strictures [4].

Here we present a case of traumatic biliary disconnection previously treated by

biliodigestive latero-lateral anastomosis on Roux-en-Y-loop, complicated by anastomotic stricture.

A 42-year-old man without relevant comorbidities was unseated from his horse and reported a severe bile leak due to intrapancreatic biliary tract disconnection. For this reason, he underwent hepaticojejunostomy latero-lateral anastomosis on Roux-en-Y-loop, with previous ligation of the distal choledocus.


After six months, a biliary anastomotic stricture developed and percutaneous

transhepatic biliary drainage was performed as the primary treatment.

Because of the very low quality of life due to poor tolerance/compliance with percutaneous transhepatic biliary drainage, the patient was referred to our endoscopy center.

At first, a standard approach to the major duodenal papilla was attempted with a duodenoscope (ED-127; Pentax Medical), but it failed because the intrapancreatic biliary tract was completely disconnected.

At this point, an occlusive cholangiography was performed (using a Fogarty balloon pushed into the bile duct through the percutaneous access) (► **Fig. 1**) in order to make the common bile duct visible on endoscopic ultrasound. The Hot AXIOS electrocautery-enhanced delivery system (Boston Scientific) was chosen to change the approach, and a fully covered, 8×8-mm lumen-apposing metal stent (LAMS) was released between the duodenum and the choledocal stump (► **Fig. 2**). Then a fully covered, 16-mm×2-cm biflanged metal stent (Nagi Stent; Euro-Medical Corp.) was placed through the hepaticojejunostomy using the percutaneous access to achieve stable dilation of the anastomotic stricture (► **Fig. 3**).



▶ Video 1 A case of traumatic choledochal disconnection previously treated by biliodigestive latero-lateral anastomosis on Roux-en-Y-loop complicated by anastomotic stricture and solved by “tailored” endoscopic treatment.

At the six-month follow-up, both stents were removed, and two “hand-tailored,” fully covered, 6-mm×3-cm self-expandable metal stents (WallFlex; Boston Scientific) were placed through the new choledocoduodenostomy to consolidate the anastomosis (▶ **Video 1**). The stents were positioned crosswise in order to avoid excessive tightening on the walls of the biliary ducts (▶ **Fig. 4**).

These stents were removed after six months and, at the cholangiography, the choledocoduodenostomy was patent (▶ **Fig. 5**). The patient remained asymptomatic and no recurrence of symptoms was reported.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests

None

The authors

Valerio Pontecorvi¹, Lorenzo Dioscoridi¹, Edoardo Forti¹, Marcello Cintolo¹, Aurora Giannetti¹, Simone Vagnarelli², Massimiliano Mutignani¹

- 1 Digestive and Interventional Endoscopy Unit, Azienda Socio-Sanitaria Territoriale Grande Ospedale Metropolitan Niguarda, Milano, Italy
- 2 Interventional Radiology, San Giovanni Hospital, Rome, Italy

Corresponding author

Massimiliano Mutignani, MD
Endoscopy Unit Niguarda Ca-Granda Hospital, Piazza dell’Ospedale Maggiore 3, 20162 Milano, Italy
Fax: +39-02-64442911
massimiliano.mutignani@ospedaleniguarda.it

References

- [1] Navsaria PH, Nicol AJ, Edu S et al. Selective nonoperative management in 1106 patients with abdominal gunshot wounds: conclusions on safety, efficacy, and the role of selective CT imaging in a prospective single center study. *Ann Surg* 2015; 261: 760–764
- [2] Tiwari C, Shah H, Waghmare M et al. Management of traumatic liver and bile duct laceration. *Euroasian J Hepato Gastroenterol* 2017; 7: 188–190
- [3] Wong VW, Gee A, Hansen P et al. Isolated avulsion of the common hepatic duct from blunt abdominal trauma. *Case Rep Surg* 2012; 2012: 254563
- [4] Dimou FM, Adhikari D, Mehta HB et al. Incidence of hepaticojejunostomy stricture after hepaticojejunostomy. *Surgery* 2016; 160: 691–698

Bibliography

DOI <https://doi.org/10.1055/a-1045-4246>
Published online: 2.12.2019
Endoscopy 2020; 52: E172–E173
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS
<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>