



## Peroral endoscopic myotomy and simultaneous endoscopic diverticuloseptotomy in a case of achalasia with diverticula




► **Fig. 1** Contrast esophagogram before peroral endoscopic myotomy shows a large epiphrenic diverticulum (black triangle) on the right anterolateral esophageal wall. A thin streak of contrast medium is visible across the lower esophageal sphincter (black star).

A 58-year-old man with symptoms of dysphagia and regurgitation was admitted for treatment. Evaluation with endoscopy and contrast esophagogram revealed achalasia cardia with a small mid-esophageal diverticulum and a large epiphrenic diverticulum (► **Fig. 1**, ► **Fig. 2**). Peroral endoscopic myotomy (POEM) and a simultaneous endoscopic diverticuloseptotomy were performed in this case (► **Video 1**). Firstly, submucosal injection and mucosal incision were made at about 5 cm above the epiphrenic diverticulum. Secondly, a submucosal tunnel was created pointing toward the diverticular septum and extended on both sides of the septum, i.e., the diverticular and esophageal lumen side (► **Fig. 3**). Thirdly, complete myotomies of the muscle layer of the diverticular septum and the esophagus were performed separately. Due to technical difficulty, a small mucosal perforation occurred at the most narrow, twisted, and spasmodic part of the distal




► **Fig. 2** Endoscopic images show a small mid-esophageal diverticulum (black arrow) and a large epiphrenic diverticulum (black triangle). White arrows show the entrance of the narrow lumen of distal esophagus.




► **Video 1** Peroral endoscopic myotomy and simultaneous endoscopic diverticuloseptotomy in a case of achalasia with diverticula.

esophagus (► **Fig. 4**). In the final step, the small mucosal injury and the entry of the tunnel were closed with endoclips. An X-ray contrast study 5 days after POEM revealed a free flow of contrast medium across the gastroesophageal junction and a collapsed epiphrenic diverticulum (► **Fig. 5**). There was substantial improvement in the patient's clinical symptoms as well. At the 3-month follow-up, the

patient reported complete resolution of dysphagia. POEM has been introduced for achalasia treatment as a less invasive alternative to laparoscopic Heller myotomy [1]. However, a few patients with achalasia have a co-existing large epiphrenic diverticulum, which may cause technical difficulties and increase the rates of procedure-related adverse events [2, 3]. In our



► **Fig. 3** A submucosal tunnel was created pointing toward the diverticular septum and extended on both sides of the septum.



► **Fig. 4** A small mucosal injury occurred at the most narrow, twisted, and spasmodic part of the distal esophagus during peroral endoscopic myotomy (black arrow).

case, a small esophageal mucosal injury occurred during POEM, which was completely sealed by an endoclip without any postoperative complications. Classically, a diverticulum of the middle esophagus is classified as a Rokitansky diverticulum, and it rarely attains an appreciable size or produces any symptoms [4]. We did not treat the small mid-esophageal diverticulum in this case.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AM



► **Fig. 5** Contrast esophagogram 5 days after peroral endoscopic myotomy shows quick passage of contrast medium across the gastroesophageal junction into the stomach; the previously large diverticulum has collapsed.

### Competing interests

None

### The authors

Wen Li<sup>1</sup>, Fang Liu<sup>1</sup>, Liang Wu<sup>2</sup>

- 1 Department of Gastroenterology and Hepatology, The First Medical Center of Chinese PLA General Hospital, Beijing, China
- 2 International Center for Diagnosis and Treatment of Liver Diseases, The Fifth Medical Center of Chinese PLA General Hospital, Beijing, China

### Corresponding author

**Liang Wu**

International Center for Diagnosis and Treatment of Liver Diseases, The Fifth Medical Center of Chinese PLA General Hospital, Beijing, 100039, China  
 Fax: +86-10-55499107  
 wuliangdoc@163.com

### References

- [1] Zaninotto G, Leusink A, Markar SR. Management of achalasia in 2019. *Curr Opin Gastroenterol* 2019. doi:10.1097/MOG.0000000000000544
- [2] Bechara R, Woo M, Hookey L et al. Peroral endoscopic myotomy (POEM) for complex achalasia and the POEM difficulty score. *Dig Endosc* 2019; 31: 148–155
- [3] Li HK, Linghu EQ. New endoscopic classification of achalasia for selection of candidates for peroral endoscopic myotomy. *World J Gastroenterol* 2013; 19: 556–560
- [4] Sato H, Takeuchi M, Hashimoto S et al. Esophageal diverticulum: New perspectives in the era of minimally invasive endoscopic treatment. *World J Gastroenterol* 2019; 25: 1457–1464

### Bibliography

DOI <https://doi.org/10.1055/a-1059-9322>

Published online: 2.12.2019

Endoscopy 2020; 52: E168–E169

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>