

## Single-session bridge-to-surgery choledochooduodenostomy and duodenal stenting in patient with malignant biliary and duodenal obstruction



► **Fig. 1** Computed tomography scan confirmed correct positioning of both stents.

Endoscopic ultrasound (EUS)-guided biliary drainage using lumen-apposing metal stent (LAMS) is considered an effective alternative approach when endoscopic retrograde cholangiopancreatography fails in patients with malignant obstructive jaundice. Duodenal neoplastic stenosis may coexist and patient management becomes challenging. Sequential placement of biliary and duodenal stents has been described previously in patients with unresectable pancreatic cancer [1]. We report the case of a 73-year-old man affected by obstructive jaundice, cholangitis, and vomiting due to resectable cancer of the pancreatic head. Secondary duodenal infiltration prevented access to the papilla of Vater, and therefore EUS-guided choledochooduodenostomy (EUS-CD) was performed.

From the duodenal bulb, an 8×8 mm LAMS (Hot Axios; Boston Scientific, Marlborough, Massachusetts, USA) was directly deployed, and good biliary drainage was obtained. During the same session, an uncovered 60×10 mm self-expandable metal stent (Wallflex; Boston Scientific) was deployed across the 30 mm length of the duodenal stenosis, taking care not to dislocate the

**Single-session bridge-to-surgery  
choledocho-duodenostomy and duodenal  
stenting in patient with malignant biliary and duodenal  
obstruction**

Manno M MD<sup>1</sup>, Scopelliti F MD<sup>2</sup>, Gabbani T MD<sup>1</sup>, Deiana S MD<sup>1</sup>, Ottaviani L MD<sup>1</sup>, Vavassori S MD<sup>1</sup>, Soriani P MD<sup>1</sup>  
Gastroenterology and Digestive Endoscopy Unit, Azienda USL Modena, Carpi (MO), Italy  
Department of Hepato-Pancreato-Biliary Surgery, Pederzoli Hospital, Peschiera del Garda (VR), Italy

► **Video 1** Single-session bridge-to-surgery choledochooduodenostomy and duodenal stenting in patient with malignant biliary and duodenal obstruction.

LAMS (► **Video 1**). The proximal flange was positioned within the duodenal bulb, adjacent to the LAMS. Subsequent computed tomography scan confirmed the correct position of both stents (► **Fig. 1**). Jaundice progressively resolved, and the patient restarted oral feeding and was referred to surgery.

The LAMS distal flange was positioned within the common bile duct; therefore, it was possible to easily perform the usual common hepatic duct jejunostomy. The proximal flange of both stents (LAMS and duodenal stent) was located within the duodenal bulb (and not transpylorically). A pylorus-preserving pancreaticoduodenostomy, rather than pylorus-resecting pancreaticoduodenostomy, was therefore performed [2].

In conclusion, even in challenging patients with malignant distal biliary and duodenal obstruction by resectable pancreatic cancer, bridge-to-surgery single-session EUS-CD and duodenal stenting is feasible and effective. However, attention is required when placing the duodenal stent to avoid LAMS dislocation or compromise further surgical treatment.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AZ

### Competing interests

The authors declare that they have no conflict of interest.

### The authors

**Mauro Manno<sup>1</sup>, Filippo Scopelliti<sup>2</sup>, Tommaso Gabbani<sup>1</sup>, Simona Deiana<sup>1</sup>, Laura Ottaviani<sup>1</sup>, Sara Vavassori<sup>1</sup>, Paola Soriani<sup>1</sup>**

- 1 Gastroenterology and Digestive Endoscopy Unit, Azienda USL Modena, Carpi, Italy
- 2 Department of Hepato-Pancreato-Biliary Surgery, Pederzoli Hospital, Peschiera del Garda, Italy

### Corresponding author

**Mauro Manno, MD**

Gastroenterology and Digestive Endoscopy Unit, Azienda USL Modena, Via Guido Molinari 2, 41012 Carpi (MO), Italy  
Fax: +39-059-659250  
m.manno@ausl.mo.it

## References

- [1] Anderloni A, Buda A, Carrara S et al. Single-session double-stent placement in concomitant malignant biliary and duodenal obstruction with a cautery-tipped lumen apposing metal stent. *Endoscopy* 2016; 48: E321–E322
- [2] Klaiber U, Probst P, Buchler MW et al. Pylorus preservation pancreatectomy or not. *Transl Gastroenterol Hepatol* 2017; 2: 100

## Bibliography

DOI <https://doi.org/10.1055/a-11119-0960>  
Published online: 27.2.2020  
*Endoscopy* 2020; 52: E318–E319  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

### ENDOSCOPY E-VIDEOS <https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>