


Colitis cystica profunda mimicking mucinous adenocarcinoma of the rectum diagnosed by endoscopic submucosal dissection


A 48-year-old man presented to our hospital with complaints of fresh rectal bleeding. He was intellectually disabled and often inserted his fingers into the anus after defecation. Laboratory investigations demonstrated lowered hemoglobin level (9.3 g/dL; normal range 13.5–17.5 g/dL). Colonoscopy showed a polypoid submucosal tumor-like elevated lesion with redness and rough surface, along with erosion in the lower rectum (► Fig. 1 a, b). Endoscopic ultrasonography (EUS) showed hypoechoic and several anechoic cystic lesions in the submucosa, without invasion of the muscularis propria (► Fig. 1 c). Magnetic resonance imaging (MRI) showed peripheral contrast-enhanced cystic lesions in the lower rectum. Regional lymphadenitis in the lower rectum was not seen on MRI (► Fig. 2). Boring biopsy was suggestive of mucinous adenocarcinoma.

After obtaining informed consent, we performed endoscopic submucosal dissection (ESD) for complete en bloc resection to confirm the preoperative diagnosis (► Fig. 1 d, ► Fig. 3 a, ► Video 1). Histological examination revealed colitis cystica profunda (CCP) with no evidence of dysplasia or malignancy (► Fig. 3 b, c). The patient progressed well after ESD and was discharged 7 days postoperatively. The patient was treated conservatively to improve bowel habits.

CCP is a rare benign disease characterized by mucin-filled cysts beneath the muscularis mucosa [1]. Knowledge of this pathological entity is important as it can mimic a well-differentiated adenocarcinoma of the rectum and therefore could lead to unnecessary surgical resection [2]. EUS, computed tomography, and MRI are useful for diagnosis of CCP [3, 4]; however, the definitive diagnosis is based on endoscopic biopsies, which should be large and deep enough to include the submucosal layer, or resected surgical



► Fig. 1 Examination and imaging of the lesion. a, b Colonoscopy showed a polypoid submucosal tumor-like elevated lesion with redness and rough surface, and erosion in the lower rectum, approximately 7 cm in size. c Endoscopic ultrasonography showed hypoechoic and several anechoic cystic lesions located in the submucosa without invasion of the muscularis propria (red arrows). d The mucosa immediately after endoscopic submucosal dissection.



► Fig. 2 Abdominal magnetic resonance imaging (MRI). a, b Homogenous and variable hypointense nodules on T1-weighted MRI (a) and hyperintense nodules on T2-weighted fat-suppressed MRI (b) showed no infiltrative signs and regional swollen lymph nodes in the lower rectum (red arrow). c Contrast-enhanced images on T2-weighted fat-suppressed MRI showed peripheral enhancement (red arrow).

specimens, as small endoscopic biopsies are frequently unrevealing [5]. ESD may provide an endoscopic technique for diagnosis and treatment of CCP.

Endoscopy_UCTN_Code_CCL_1AD_2AC

Competing interests

The authors declare that they have no conflict of interest.


The authors

Takashi Abe¹, Marina Hamamoto¹, Takayuki Nagai¹, Takehiko Nariyasu¹, Masato Hanzawa¹, Yasuhisa Hiroshima¹, Kazunari Murakami²


- 1 Department of Gastroenterology, Oita Kouseiren Tsurumi Hospital, Beppu, Japan
- 2 Department of Gastroenterology, Faculty of Medicine, Oita University, Yufu, Japan

Corresponding author

Takashi Abe, MD, PhD
Department of Gastroenterology, Oita Kouseiren Tsurumi Hospital, Tsurumi 4333, Beppu City, Oita 874-8585, Japan
Fax: +81-977-237884
takashi0315@oita-u.ac.jp



► **Fig. 3** The resected specimen. **a** Photograph of the resected specimen, approximately 9×6 cm in size. **b** Macroscopic view of the resected specimen, showing several cystic lesions containing mucin. **c** Histological images of the resected specimen stained with hematoxylin and eosin, showing colitis cystica profunda without findings of dysplasia or malignancy (× 12.5).



► **Video 1** Endoscopic submucosal dissection of colitis cystica profunda in the lower rectum along with resection of the tumor using the pocket-creation and clip-with-line methods.

References

- [1] Kornprat P, Langner C, Pfeifer J et al. Colitis cystica profunda associated with rectal prolapse: report of a case. *Int J Colorectal Dis* 2007; 22: 1555–1556
- [2] Kayaçetin E, Kayaçetin S. Colitis cystica profunda simulating rectal carcinoma. *Acta Chir Belg* 2005; 105: 306–308
- [3] Valenzuela M, Martín-Ruiz JL, Alvarez-Cienfuegos E et al. Colitis cystica profunda: imaging diagnosis and conservative treatment: report of two cases. *Dis Colon Rectum* 1996; 39: 587–590
- [4] Rumi N, Cilla S, De Ninno M et al. Colitis cystica profunda of the rectum with adenomatous dysplastic features: radiologic-

pathologic correlation. *Radiol Case Rep* 2019; 14: 740–745

- [5] Spicakova K, Pueyo BA, de la Piscina PR et al. Colitis cystica profunda: a report of 2 cases with a 15-year follow-up. *Gastroenterol Hepatol* 2017; 40: 406–408

Bibliography

Endoscopy 2021; 53: E157–E159

DOI 10.1055/a-1224-3347

ISSN 0013-726X

published online 20.8.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>