



## Combination of endoscopic-ultrasound guided choledochoduodenostomy and gastrojejunostomy resolving combined distal biliary and duodenal obstruction




► **Fig. 1** Computed tomography revealed a 3.5×3.5×3.2-cm heterogeneous enhancing periampullary mass with adjacent bowel wall invasion at the second part duodenum, causing luminal narrowing of the second part duodenum and upstream dilatation of the common bile duct.




► **Fig. 2** A large friable ampullary mass causing supra-ampullary duodenal obstruction.




► **Fig. 3** Endoscopic ultrasound revealed a periampullary mass (mass) with a dilated common bile duct (CBD).



► **Video 1** Endoscopic-ultrasound guided choledochoduodenostomy and gastrojejunostomy resolving combined distal biliary and duodenal obstruction in patient with periampullary cancer.



► **Fig. 4** An 8×12-mm lumen-apposing metal stent (LAMS) was successfully placed transduodenally into the distal common bile duct.



► **Fig. 5** A 16×20-mm lumen-apposing metal stent (LAMS) was successfully placed transgastrically into the lumen of the jejunum.

A 68-year-old man presented with abdominal pain, jaundice, and weight loss for 1 month. Abdominal computed tomography revealed a periampullary mass measuring 3.5×3.5×3.2 cm with dilated bile duct (► **Fig. 1**).

An endoscopic retrograde cholangiopancreatography (ERCP) procedure was not possible owing to a large friable ampullary mass causing supra-ampullary duodenal obstruction (► **Fig. 2**). An endo-

scopic ultrasound-guided choledochoduodenostomy (EUS-CDS) was consequently performed (► **Video 1**) with a linear echoendoscope (GF-UCT180; Olympus, Aizu, Japan). A dilated distal common bile duct (CBD) from an ampullary mass was shown (► **Fig. 3**). A 19-gauge endoscopic ultrasound-guided fine-needle aspiration (EUS-FNA) needle (Echotip Ultra; Cook Medical Ltd., Limerick, Ireland) with an angled 0.025-inch guide-

wire (Visiglide 2, Olympus) was used for puncturing. A 6-Fr cystotome (Endo-Flex, Voerde, Germany) and a 4-mm balloon dilatation catheter (Hurricane RX; Boston Scientific, Cork, Ireland) were used for dilation. An 8×12-mm lumen-apposing metal stent (LAMS) (Niti-S Spaxus; Taewoong Medical Co., Ilsan, Korea) was successfully placed transduodenally into the distal CBD (► Fig. 4). Subsequently, an EUS-guided gastrojejunostomy was performed. A 10-Fr nasobiliary catheter (Flexima; Boston Scientific, Marlborough, Massachusetts, USA) was placed into the jejunum to flush a mix of diluted contrast, saline, and methylene blue into the lumen of the jejunum in order to distend the small bowel loop. A 16×20-mm LAMS with an electrocautery delivery system (Niti-S Spaxus; Taewoong Medical Co.) was successfully placed transgastrically into the lumen of the jejunum (► Fig. 5). The patient resumed diet with a decline of bilirubin level at 48 hours after the procedure without adverse events.

This case reported the feasibility of a combination of EUS-guided choledochoduodenostomy and EUS-guided gastrojejunostomy to resolve a problem of bile duct and duodenal obstruction type II [1]. Previously, most literature used a combination of EUS-guided biliary drainage and duodenal stents with a technical and clinical success rate of 71.4% to 100% and 94.1% to 100%, respectively [2]. Future study to compare the efficacy of a combined EUS-guided biliary drainage with EUS-guided gastrojejunostomy versus EUS-guided biliary drainage with a duodenal stent is warranted.

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AD

## Competing interests

The authors declare that they have no conflict of interest.

## The authors

**Pradermchai Kongkam<sup>1,2</sup>, Thanawat Luangsukrer<sup>1</sup>, Kamin Harinwan<sup>3</sup>, Kunvadee Vanduangden<sup>1</sup>, Suppawatsa Plaidum<sup>1</sup>, Rungsun Rerknimitr<sup>1</sup>, Pinit Kullavanijaya<sup>1</sup>**

- 1 Gastrointestinal Endoscopy Excellence Center, Department of Medicine, Faculty of Medicine, Chulalongkorn University and King Chulalongkorn Memorial Hospital, Thai Red Cross Society, Bangkok, Thailand
- 2 Pancreas Research Unit, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand
- 3 Division of Gastroenterology, Department of Medicine, Faculty of Medicine, Phramongkutklao Hospital, Phramongkutklao College of Medicine, Bangkok, Thailand

## Corresponding author

**Thanawat Luangsukrer, MD**  
Gastrointestinal Endoscopy Excellence Center, Chulalongkorn University and King Chulalongkorn Memorial Hospital, Thai Red Cross Society, 1873, Rama 4 Road, Patumwan, Bangkok, Thailand 10500  
Fax: +66-2-652-4219  
drthanawat@gmail.com

## References

- [1] Mutignani M, Tringali A, Shah SG et al. Combined endoscopic stent insertion in malignant biliary and duodenal obstruction. *Endoscopy* 2007; 39: 440–447
- [2] Mangiavillano B, Khashab MA, Tarantino I et al. Success and safety of endoscopic treatments for concomitant biliary and duodenal malignant stenosis: a review of the literature. *World J Gastrointest Surg* 2019; 11: 53–611

## Bibliography

*Endoscopy* 2021; 53: E355–E356  
DOI 10.1055/a-1294-9399  
ISSN 0013-726X  
published online 19.11.2020  
© 2020. Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at  
<https://mc.manuscriptcentral.com/e-videos>