


Intracavity rendezvous procedure




► **Fig. 1** Abdominal computed tomography demonstrates the intra-abdominal collection.




► **Fig. 3** Percutaneous transhepatic cholangiography shows contrast leakage at the biliary anastomosis with no delineation of the recipient duct.



► **Fig. 5** Fully covered self-expanding metal stent deployed across the anastomosis and contrast passed through the stent into the small bowel with no leakage.



► **Fig. 2** Endoscopic retrograde cholangiopancreatography shows contrast leakage at the biliary anastomosis.



► **Fig. 4** Guidewire passed successfully across the anastomosis.

Biliary complications are common after liver transplantation, and the vast majority of these can be managed endoscopically [1,2]. We describe a case of a spontaneous bile leak after liver transplantation which eventually required an extra-ductal rendezvous to re-establish recipient and donor bile duct continuity.

A 47-year-old woman underwent liver transplantation (donation after brain death, duct-to-duct anastomosis) for end-stage liver cirrhosis. Two weeks after transplantation she was diagnosed with a bile leak (► **Fig. 1**). The patient initially had two endoscopic retrograde cholangiopancreatograms performed and both demonstrated a bile leak at the level of the biliary anastomosis. On both occasions it proved impossible to pass a guidewire into the donor duct (► **Fig. 2**). A percutaneous transhepatic cholangio-

gram was performed, but the recipient duct could not be accessed (► **Fig. 3**). A rendezvous procedure was undertaken. Endoscopically, a guidewire (450 cm, 0.018 inch; Terumo, Tokyo, Japan) was passed into the biloma. The guidewire was then snared with an Amplatz Goose Neck loop snare (6Fr, 15 mm; ev3 Inc. Plymouth, Maine, USA) to establish access across the leak (► **Fig. 4**; ► **Video 1**). The Terumo wire was then exchanged for a VisiGlide wire (450 cm, 0.025 inch; Olympus, USA). The donor:recipient duct anastomotic site was dilated with a dilatation balloon (6 mm, 4 cm; Hurricane RX, Boston Scientific, USA) before a fully covered self-expanding metal stent (8 mm, 4 cm; Kaffes stent, Taewoong Medical, Japan) was inserted over the guidewire across the anastomosis. Further contrast injection through the percutaneous route did not demonstrate a bile leak (► **Fig. 5**). The patient was discharged home 2 days later without any complications.



Video 1 Rendezvous procedure in the biloma to re-establish donor:recipient duct continuity following spontaneous bile leak after liver transplantation.

An intracavity rendezvous procedure is a viable management solution in cases where endoscopic retrograde cholangiopancreatography and the percutaneous transhepatic approach have failed to resolve a bile leak after liver transplantation. A novel fully covered self-expanding metal stent (Kaffes stent) can be used to bridge the anastomotic area between the donor and the recipient duct.

Endoscopy_UCTN_Code_TTT_1AR_2AJ

Competing interests

The authors declare that they have no conflict of interest.

The authors

Muhammad Farman¹, Pauline Kane², Stephen Gregory², Andreas Prchalias¹, Deepak Joshi¹

- 1 Institute of Liver Studies, King's College Hospital NHS Foundation Trust, London, UK
- 2 Department of Radiology, King's College Hospital NHS Foundation Trust, London, UK

Corresponding author

Muhammad Farman, MBBS, MRCPI
Institute of Liver Studies, King's College Hospital NHS Foundation Trust, Denmark Hill, London, SE5 9RS, UK
mfarman83@gmail.com

References

- [1] Kochhar G, Parungao JM, Hanouneh IA et al. Biliary complications following liver transplantation. *World J Gastroenterol* 2013; 19: 2841–2846
- [2] Thuluvath PJ, Pfau PR, Kimmey MB et al. Biliary complications after liver transplantation: the role of endoscopy. *Endoscopy* 2005; 37: 857–863

Bibliography

Endoscopy 2021; 53: E466–E467

DOI 10.1055/a-1346-7555

ISSN 0013-726X

published online 4.2.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>