

## Endoscopic septotomy for fistula after bariatric surgery

Gastric fistula following bariatric surgery is a complication with considerable morbidity and mortality [1,2]. After Roux-en-Y gastric bypass (RYGB), the gastric pouch may develop a chronic fistula and the formation of a perigastric cavity bounded by a septum. The altered anatomy chronically elevates the intraluminal pressure, impairing emptying of the perigastric cavity and perpetuating the fistula [3].

Endoscopic septotomy is a minimally invasive technique for cutting the septum. The goals are ensuring adequate patency of the perigastric cavity, decreasing its pressure, and draining the fistula [1, 4, 5]. A 67-year-old woman underwent a RYGB (body mass index: 48 kg/m<sup>2</sup> before, 25 kg/m<sup>2</sup> after). She developed a gastrocutaneous fistula, which was unsuccessfully treated with a long-term nasoenteric tube. Eight months after RYGB, she was referred for endoscopic assessment (► **Video 1**). Sutures were identified in the greater curvature of the gastric pouch. They were removed with endoscopic scissors and a perigastric cavity (bounded by a septum) with a fistula orifice in it was identified. The fistula orifice was initially treated with argon plasma coagulation and a guidewire was externalized through the fistula's cutaneous orifice (► **Fig. 1**), with placement of a 7-Fr double-pigtail stent in the fistula tract. After 3 months, the gastrocutaneous leak was reduced but not resolved. We removed the pigtail, performed a septotomy with an IT knife, and placed an esophageal fully covered (28 × 160 mm) self-expandable metal stent. The proximal end of the stent was fixed by endosuture to avoid migration. After 1 week, the stent was removed and the patient was able to take a soft diet. Four weeks later, the cutaneous fistula orifice was closed (► **Fig. 2**) and esophagogastro-duodenoscopy confirmed closure of the



► **Video 1** Complex fistulas after bariatric surgery require challenging endoscopic management. We report a case of bariatric surgery complicated by gastrocutaneous fistula, which was successfully treated with endoscopic septotomy.



► **Fig. 1** Guidewire externalized through the cutaneous orifice of the fistula.




► **Fig. 2** The cutaneous fistula orifice is closed.

fistula's gastric orifice (► **Fig. 3**). At 3 weeks' follow-up, the patient was asymptomatic and doing well on a regular diet.

Endoscopy\_UCTN\_Code\_CPL\_1AH\_2AG

### Competing interests

The authors declare that they have no conflict of interest.



**Fig. 3** Endoscopic appearance of the perigastric cavity: **a** after suture removal, **b** after 7-Fr double-pigtail insertion through the fistula orifice, and **c** after treatment with argon plasma coagulation and 7-Fr double-pigtail stent. **d** Closed fistula after septotomy (final appearance).

## Bibliography

Endoscopy 2022; 54: E38–E39

DOI 10.1055/a-1375-0159

ISSN 0013-726X

published online 19.2.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,

70469 Stuttgart, Germany

### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>

## The authors

**Mateus Bond Boghossian**  **Mateus Pereira Funari**  **Epifanio Silvino do Monte Junior**  **Rodrigo Silva de Paula Rocha**, **Diogo Turiani Hourneaux de Moura**  **Thiago Ferreira de Souza**, **Eduardo Guimarães Hourneaux de Moura** 

Gastrointestinal Endoscopy Unit, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil

## Corresponding author

**Mateus Bond Boghossian, MD**

Av. Dr Enéas de Carvalho Aguiar 225, 6º andar, bloco 3, Cerqueira César, 05403-010 – São Paulo, SP, Brazil  
mateus.boghossian@hc.fm.usp.br

## References

- [1] Baretta G, Campos J, Correia S et al. Bariatric postoperative fistula: a life-saving endoscopic procedure. *Surg Endosc* 2015; 29: 1714–1720
- [2] Campos JM, Pereira EF, Evangelista LF et al. Gastrobronchial fistula after sleeve gastrectomy and gastric bypass: endoscopic management and prevention. *Obes Surg* 2011; 21: 1520–1529
- [3] Angrisani L, Hasani A, Santonicola A et al. Endoscopic septotomy for the treatment of sleeve gastrectomy fistula: timing and indications. *Obes Surg* 2018; 3: 846–847
- [4] Rodrigues-Pinto E, Repici A, Donatelli G et al. International multicenter expert survey on endoscopic treatment of upper gastrointestinal anastomotic leaks. *Endosc Int Open* 2019; 7: E1671–E1682
- [5] Mahadev S, Kumbhari V, Campos JM et al. Endoscopic septotomy: an effective approach for internal drainage of sleeve gastrectomy-associated collections. *Endoscopy* 2017; 49: 504–508